



# wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 08:16 pm GMT

PDB ID : 1VEE  
Title : NMR structure of the hypothetical rhodanese domain At4g01050 from *Arabidopsis thaliana*  
Authors : Pantoja-Uceda, D.; Lopez-Mendez, B.; Koshiba, S.; Inoue, M.; Kigawa, T.; Terada, T.; Shirouzu, M.; Tanaka, A.; Seki, M.; Shinozaki, K.; Yokoyama, S.; Guntert, P.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2004-03-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)  
NmrClust : Kelley et al. (1996)  
MolProbity : 4.02b-467  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
ShiftChecker : trunk28760  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	134	

## 2 Ensemble composition and analysis

This entry contains 20 models. Model 19 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:6-A:125 (120)	0.33	19

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 7, 8, 11, 14, 16, 17, 18, 19, 20
2	3, 5, 6, 10, 13
3	4, 15
4	1, 9
Single-model clusters	12

### 3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2002 atoms, of which 1001 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called proline-rich protein family.

Mol	Chain	Residues	Atoms					Trace
1	A	134	Total	C	H	N	O	0
			2002	631	1001	170	200	

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q9M158
A	2	SER	-	CLONING ARTIFACT	UNP Q9M158
A	3	SER	-	CLONING ARTIFACT	UNP Q9M158
A	4	GLY	-	CLONING ARTIFACT	UNP Q9M158
A	5	SER	-	CLONING ARTIFACT	UNP Q9M158
A	6	SER	-	CLONING ARTIFACT	UNP Q9M158
A	7	GLY	-	CLONING ARTIFACT	UNP Q9M158
A	129	SER	-	CLONING ARTIFACT	UNP Q9M158
A	130	GLY	-	CLONING ARTIFACT	UNP Q9M158
A	131	PRO	-	CLONING ARTIFACT	UNP Q9M158
A	132	SER	-	CLONING ARTIFACT	UNP Q9M158
A	133	SER	-	CLONING ARTIFACT	UNP Q9M158
A	134	GLY	-	CLONING ARTIFACT	UNP Q9M158

## 4 Residue-property plots [i](#)

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: proline-rich protein family



### 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 19. Colouring as in section 4.1 above.

- Molecule 1: proline-rich protein family



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORSION ANGLE DYNAMICS*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *STRUCTURES WITH THE LEAST RESTRAINT VIOLATIONS*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
OPALP	refinement	1.3
CYANA	structure solution	2.0.29

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5929
Number of chemical shift lists	1
Total number of shifts	1609
Number of shifts mapped to atoms	1609
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%

No validations of the models with respect to experimental NMR restraints is performed at this time.

## 6 Model quality i

### 6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.59±0.01	0±0/936 (0.0±0.0%)	1.03±0.02	1±1/1266 (0.1±0.1%)
All	All	0.59	0/18720 (0.0%)	1.03	14/25320 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.1±0.9
All	All	0	21

There are no bond-length outliers.

5 of 7 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	78	TYR	CB-CG-CD2	-6.74	116.95	121.00	20	1
1	A	28	ARG	NE-CZ-NH2	-6.57	117.01	120.30	14	4
1	A	28	ARG	CD-NE-CZ	6.09	132.13	123.60	6	1
1	A	113	ARG	NE-CZ-NH2	-6.04	117.28	120.30	1	5
1	A	113	ARG	N-CA-CB	-5.77	100.21	110.60	5	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	78	TYR	Sidechain	9
1	A	113	ARG	Sidechain	6
1	A	28	ARG	Sidechain	4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group	Models (Total)
1	A	103	TYR	Sidechain	2

## 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	917	919	919	2±1
All	All	18340	18380	18380	39

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

5 of 30 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:80:LEU:HD21	1:A:109:ALA:HB2	0.63	1.71	1	1
1:A:69:PHE:CD1	1:A:75:THR:HG21	0.61	2.30	11	5
1:A:22:ALA:HB1	1:A:78:TYR:CE1	0.53	2.39	10	1
1:A:69:PHE:CE1	1:A:75:THR:HG21	0.53	2.39	11	1
1:A:40:ASN:HB3	1:A:123:ILE:CD1	0.52	2.35	15	1

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/134 (90%)	107±3 (89±2%)	11±3 (9±2%)	2±1 (1±1%)	18	63
All	All	2400/2680 (90%)	2144 (89%)	222 (9%)	34 (1%)	18	63

5 of 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.



Mol	Chain	Res	Type	Models (Total)
1	A	85	GLY	14
1	A	84	ASP	4
1	A	108	GLY	3
1	A	98	GLY	3
1	A	7	GLY	2

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/107 (91%)	88±2 (90±3%)	9±2 (10±3%)	14	59
All	All	1940/2140 (91%)	1751 (90%)	189 (10%)	14	59

5 of 45 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	68	LYS	20
1	A	40	ASN	16
1	A	65	LEU	12
1	A	70	LYS	12
1	A	89	LEU	8

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.6 Ligand geometry

There are no ligands in this entry.

## 6.7 Other polymers

There are no such molecules in this entry.

## 6.8 Polymer linkage issues

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 93% for the entire structure.

### 7.1 Chemical shift list 1

File name: BMRB entry 5929

Chemical shift list name: *assigned\_chem\_shift\_list\_1*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1609
Number of shifts mapped to atoms	1609
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	6

#### 7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	134	$2.48 \pm 0.16$	Should be applied
$^{13}\text{C}_\beta$	118	$2.84 \pm 0.10$	Should be applied
$^{13}\text{C}'$	133	$2.70 \pm 0.13$	Should be applied
$^{15}\text{N}$	127	$-0.16 \pm 0.21$	None needed ( $< 0.5$ ppm)

#### 7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 1331 atoms were assigned a chemical shift out of a possible 1427. 19 out of 19 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	588/588 (100%)	234/234 (100%)	240/240 (100%)	114/114 (100%)
Sidechain	665/738 (90%)	414/432 (96%)	238/274 (87%)	13/32 (41%)

*Continued on next page...*

Continued from previous page...

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Aromatic	78/101 (77%)	49/53 (92%)	27/46 (59%)	2/2 (100%)
Overall	1331/1427 (93%)	697/719 (97%)	505/560 (90%)	129/148 (87%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	28	ARG	NE	117.20	92.63 – 76.73	20.5
1	A	128	THR	CB	58.90	78.10 – 61.30	-6.4
1	A	28	ARG	HH22	9.21	8.76 – 4.76	6.1
1	A	38	SER	C	165.00	183.48 – 165.88	-5.5
1	A	82	LYS	HG2	-0.02	2.67 – 0.07	-5.3
1	A	28	ARG	HH21	9.21	9.06 – 4.56	5.3

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

