



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 08:09 am GMT

PDB ID : 2VES  
Title : CRYSTAL STRUCTURE OF LPXC FROM PSEUDOMONAS AERUGINOSA COMPLEXED WITH THE POTENT BB-78485 INHIBITOR  
Authors : Mochalkin, I.; Knafels, J.D.  
Deposited on : 2007-10-26  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>4%</div> <div> <div></div> <div>93%</div> <div>6%</div> <div></div> </div> </div>
1	B	299	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>6%</div> <div></div> </div> </div>
1	C	299	<div> <div>2%</div> <div> <div></div> <div>95%</div> <div>5%</div> <div></div> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-3-O-[3-HYDROXYMYRISTOYL] N-ACETYLGLUCOSAMINE DEACETYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	4	0
			2305	1461	396	443	5			
1	B	288	Total	C	N	O	S	0	7	0
			2270	1439	391	435	5			
1	C	299	Total	C	N	O	S	0	3	0
			2339	1486	398	449	6			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	SER	CYS	ENGINEERED MUTATION	UNP P47205
B	40	SER	CYS	ENGINEERED MUTATION	UNP P47205
C	40	SER	CYS	ENGINEERED MUTATION	UNP P47205

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Zn	0	0
			2	2		
2	A	2	Total	Zn	0	0
			2	2		
2	C	3	Total	Zn	0	0
			3	3		

- Molecule 3 is (2R)-N-HYDROXY-3-NAPHTHALEN-2-YL-2-[(NAPHTHALEN-2-YLSULFONYL)AMINO]PROPANAMIDE (three-letter code: GVR) (formula: C<sub>23</sub>H<sub>20</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			30	23	2	4	1		
3	B	1	Total	C	N	O	S	0	0
			30	23	2	4	1		
3	C	1	Total	C	N	O	S	0	0
			30	23	2	4	1		

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	O	S	0	0
			5	4	1		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	412	Total	O		0	0
			412	412			
5	B	402	Total	O		0	0
			402	402			
5	C	409	Total	O		0	0
			409	409			


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-3-O-[3-HYDROXYMYRISTOYL] N-ACETYLGLUCOSAMINE DEACETYLASE



- Molecule 1: UDP-3-O-[3-HYDROXYMYRISTOYL] N-ACETYLGLUCOSAMINE DEACETYLASE



- Molecule 1: UDP-3-O-[3-HYDROXYMYRISTOYL] N-ACETYLGLUCOSAMINE DEACETYLASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.54Å 103.33Å 105.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.72 – 1.90 34.15 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (73.72-1.90) 99.6 (34.15-1.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.06 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.177 , 0.221 0.193 , 0.236	Depositor DCC
$R_{free}$ test set	3927 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.0	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8244	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GVR, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2361	0.57	1/3189 (0.0%)
1	B	0.39	0/2341	0.61	2/3161 (0.1%)
1	C	0.39	0/2390	0.59	0/3230
All	All	0.38	0/7092	0.59	3/9580 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	33[A]	VAL	CG1-CB-CG2	5.23	119.26	110.90
1	B	33[B]	VAL	CG1-CB-CG2	5.23	119.26	110.90
1	A	230	TYR	CA-CB-CG	5.08	123.05	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2305	0	2318	11	0
1	B	2270	0	2281	9	0
1	C	2339	0	2350	10	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	3	0	0	0	0
3	A	30	0	19	0	0
3	B	30	0	19	1	0
3	C	30	0	19	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
5	A	412	0	0	1	0
5	B	402	0	0	1	0
5	C	409	0	0	1	0
All	All	8244	0	7006	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242:ALA:O	1:B:246:LEU:HG	2.02	0.60
1:A:10:ILE:HD12	1:A:29:LYS:HG2	1.84	0.60
1:A:242:ALA:O	1:A:246:LEU:HG	2.06	0.55
1:A:10:ILE:HD13	1:A:30:PRO:HD2	1.91	0.53
1:B:249:LEU:HG	1:B:271:LEU:HD22	1.92	0.51
1:C:10:ILE:HD12	1:C:29:LYS:HG2	1.92	0.51
1:B:279:ASP:OD1	5:B:2387:HOH:O	2.19	0.51
1:C:242:ALA:O	1:C:246:LEU:HG	2.13	0.49
1:A:163:PRO:HA	1:A:166:ARG:HD3	1.94	0.49
1:A:179:THR:CG2	1:A:184:GLU:HG3	2.43	0.48
1:C:133:ARG:HG2	1:C:280:ALA:HA	1.95	0.48
1:C:126:LYS:NZ	5:C:2225:HOH:O	2.47	0.47
1:B:163:PRO:HA	1:B:166:ARG:HD3	1.96	0.47
1:C:202:SER:O	1:C:203:GLN:HG2	2.14	0.47
1:A:5:ARG:HD3	1:A:119:LEU:HD22	1.97	0.45
1:C:5:ARG:HD3	1:C:119:LEU:HD23	1.98	0.45
1:C:86:LEU:HD22	1:C:297:MET:CE	2.46	0.45
1:A:177:SER:HB2	1:A:294:ILE:HG22	1.99	0.45
1:C:5:ARG:HD3	1:C:119:LEU:CD2	2.49	0.43
1:B:166:ARG:HD2	1:C:166:ARG:NH1	2.35	0.42
1:A:10:ILE:HD11	5:A:2024:HOH:O	2.20	0.41
1:A:10:ILE:HD12	1:A:29:LYS:CG	2.50	0.41
1:A:10:ILE:HD13	1:A:30:PRO:CD	2.50	0.41
1:A:88:ILE:HD11	1:A:119:LEU:CD2	2.50	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:86:LEU:HD22	1:C:297:MET:HE1	2.02	0.41
1:B:240:LEU:HD23	1:B:240:LEU:C	2.41	0.40
1:B:3:LYS:HB3	1:B:122:GLN:O	2.21	0.40
1:B:18:LEU:HD21	3:B:1290:GVR:C3	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/299 (99%)	289 (98%)	7 (2%)	0	100	100
1	B	293/299 (98%)	290 (99%)	3 (1%)	0	100	100
1	C	300/299 (100%)	292 (97%)	8 (3%)	0	100	100
All	All	889/897 (99%)	871 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/253 (100%)	249 (99%)	3 (1%)	75	75
1	B	251/253 (99%)	245 (98%)	6 (2%)	54	47
1	C	255/253 (101%)	253 (99%)	2 (1%)	85	85

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	758/759 (100%)	747 (98%)	11 (2%)	75	67

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	198[A]	GLU
1	A	198[B]	GLU
1	A	223	LEU
1	B	132	LYS
1	B	171	GLN
1	B	173	SER
1	B	180	SER
1	B	269[A]	GLN
1	B	269[B]	GLN
1	C	183	LYS
1	C	203	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	GLN
1	A	220	ASN
1	B	115	GLN
1	B	171	GLN
1	B	220	ASN
1	C	203	GLN
1	C	220	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GVR	A	1296	2	33,33,33	2.17	5 (15%)	46,47,47	1.29	5 (10%)
3	GVR	B	1290	2	33,33,33	2.06	3 (9%)	46,47,47	1.25	2 (4%)
4	SO4	B	1292	-	4,4,4	0.12	0	6,6,6	0.10	0
3	GVR	C	1301	2	33,33,33	2.14	4 (12%)	46,47,47	1.29	1 (2%)
4	SO4	C	1304	-	4,4,4	0.24	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GVR	A	1296	2	-	0/21/21/21	0/4/4/4
3	GVR	B	1290	2	-	0/21/21/21	0/4/4/4
4	SO4	B	1292	-	-	0/0/0/0	0/0/0/0
3	GVR	C	1301	2	-	0/21/21/21	0/4/4/4
4	SO4	C	1304	-	-	0/0/0/0	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1296	GVR	C3-S2	-10.05	1.62	1.76
3	C	1301	GVR	C3-S2	-9.95	1.62	1.76
3	B	1290	GVR	C3-S2	-9.42	1.63	1.76
3	A	1296	GVR	S2-N1	-2.26	1.58	1.61
3	C	1301	GVR	C6-C3	2.14	1.40	1.36
3	A	1296	GVR	C6-C3	2.42	1.41	1.36
3	A	1296	GVR	C19-C20	2.60	1.48	1.42
3	C	1301	GVR	C7-C8	2.63	1.48	1.42
3	C	1301	GVR	C19-C20	2.72	1.48	1.42
3	A	1296	GVR	C7-C8	2.75	1.48	1.42

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1290	GVR	C7-C8	2.76	1.48	1.42
3	B	1290	GVR	C19-C20	2.84	1.48	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1301	GVR	O14-S2-O13	-6.93	110.68	119.55
3	B	1290	GVR	O14-S2-O13	-6.25	111.55	119.55
3	A	1296	GVR	O14-S2-O13	-5.63	112.35	119.55
3	A	1296	GVR	C25-C26-N1	-2.71	107.22	110.58
3	A	1296	GVR	O28-C27-N29	-2.64	119.60	123.08
3	B	1290	GVR	O28-C27-N29	-2.38	119.94	123.08
3	A	1296	GVR	O13-S2-C3	2.45	110.98	107.95
3	A	1296	GVR	C6-C3-S2	2.64	122.22	120.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1290	GVR	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	294/299 (98%)	0.23	11 (3%) 42 46	20, 26, 34, 41	14 (4%)
1	B	288/299 (96%)	0.30	14 (4%) 30 34	20, 26, 36, 44	13 (4%)
1	C	299/299 (100%)	0.16	6 (2%) 65 69	20, 25, 35, 41	7 (2%)
All	All	881/897 (98%)	0.23	31 (3%) 44 48	20, 26, 35, 44	34 (3%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	THR	5.3
1	B	178	SER	4.1
1	B	287	GLU	3.9
1	B	179	THR	3.7
1	B	1	MET	3.6
1	C	61	THR	3.5
1	A	294	ILE	3.5
1	A	178	SER	3.4
1	B	61	THR	3.4
1	B	177	SER	3.2
1	A	59	GLU	3.2
1	A	60	THR	3.0
1	A	230	TYR	3.0
1	A	290	ARG	2.9
1	B	288	ASP	2.8
1	B	59	GLU	2.7
1	C	79[A]	LEU	2.6
1	A	166	ARG	2.5
1	B	76	VAL	2.4
1	A	167	GLY	2.3
1	C	179[A]	THR	2.3
1	B	243	ILE	2.2
1	A	240	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	203	GLN	2.2
1	B	60	THR	2.1
1	B	80	LEU	2.1
1	C	19	HIS	2.1
1	C	297	MET	2.1
1	B	81[A]	SER	2.0
1	A	243	ILE	2.0
1	B	123	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GVR	B	1290	30/30	0.97	0.08	-1.18	17,18,20,20	0
2	ZN	B	1289	1/1	1.00	0.10	-1.53	21,21,21,21	0
2	ZN	A	1295	1/1	1.00	0.09	-1.64	22,22,22,22	0
3	GVR	A	1296	30/30	0.98	0.07	-1.71	17,19,19,20	0
3	GVR	C	1301	30/30	0.98	0.08	-1.82	18,20,23,23	0
2	ZN	C	1300	1/1	1.00	0.09	-1.97	20,20,20,20	0
4	SO4	B	1292	5/5	0.98	0.07	-2.95	37,38,38,39	0
2	ZN	A	1297	1/1	1.00	0.02	-6.09	17,17,17,17	0
2	ZN	C	1302	1/1	1.00	0.01	-6.13	18,18,18,18	0
2	ZN	B	1291	1/1	1.00	0.02	-13.22	19,19,19,19	0
2	ZN	C	1303	1/1	0.99	0.11	-	45,45,45,45	0
4	SO4	C	1304	5/5	0.95	0.15	-	33,34,35,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.