



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 1, 2018 – 05:13 PM EST

PDB ID : 5VEU
Title : Human Cytochrome P450 3A5 (CYP3A5)
Authors : Hsu, M.-H.; Johnson, E.F.
Deposited on : 2017-04-05
Resolution : 2.91 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1813 (2.94-2.90)
Clashscore	112137	2045 (2.94-2.90)
Ramachandran outliers	110173	1997 (2.94-2.90)
Sidechain outliers	110143	1999 (2.94-2.90)
RSRZ outliers	101464	1825 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	
1	B	480	
1	C	480	
1	D	480	
1	E	480	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	480	
1	G	480	
1	H	480	
1	I	480	
1	J	480	
1	K	480	
1	L	480	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	RIT	A	602	-	-	-	X
3	RIT	B	602	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 44740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 3A5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	462	3714	2413	616	668	17	0	0	0
1	B	458	3679	2391	609	662	17	0	0	0
1	C	469	3773	2448	627	681	17	0	0	0
1	D	459	3687	2395	610	665	17	0	0	0
1	E	441	3540	2310	577	636	17	0	0	0
1	F	456	3661	2381	604	659	17	0	0	0
1	G	455	3651	2374	602	658	17	0	0	0
1	H	459	3687	2395	610	665	17	0	0	0
1	I	464	3732	2423	619	673	17	0	0	0
1	J	458	3679	2391	609	662	17	0	0	0
1	K	458	3681	2392	609	663	17	0	0	0
1	L	448	3590	2339	588	646	17	0	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	MET	-	initiating methionine	UNP P20815
A	23	ALA	-	expression tag	UNP P20815
A	498	HIS	-	expression tag	UNP P20815
A	499	HIS	-	expression tag	UNP P20815
A	500	HIS	-	expression tag	UNP P20815

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	501	HIS	-	expression tag	UNP P20815
B	22	MET	-	initiating methionine	UNP P20815
B	23	ALA	-	expression tag	UNP P20815
B	498	HIS	-	expression tag	UNP P20815
B	499	HIS	-	expression tag	UNP P20815
B	500	HIS	-	expression tag	UNP P20815
B	501	HIS	-	expression tag	UNP P20815
C	22	MET	-	initiating methionine	UNP P20815
C	23	ALA	-	expression tag	UNP P20815
C	498	HIS	-	expression tag	UNP P20815
C	499	HIS	-	expression tag	UNP P20815
C	500	HIS	-	expression tag	UNP P20815
C	501	HIS	-	expression tag	UNP P20815
D	22	MET	-	initiating methionine	UNP P20815
D	23	ALA	-	expression tag	UNP P20815
D	498	HIS	-	expression tag	UNP P20815
D	499	HIS	-	expression tag	UNP P20815
D	500	HIS	-	expression tag	UNP P20815
D	501	HIS	-	expression tag	UNP P20815
E	22	MET	-	initiating methionine	UNP P20815
E	23	ALA	-	expression tag	UNP P20815
E	498	HIS	-	expression tag	UNP P20815
E	499	HIS	-	expression tag	UNP P20815
E	500	HIS	-	expression tag	UNP P20815
E	501	HIS	-	expression tag	UNP P20815
F	22	MET	-	initiating methionine	UNP P20815
F	23	ALA	-	expression tag	UNP P20815
F	498	HIS	-	expression tag	UNP P20815
F	499	HIS	-	expression tag	UNP P20815
F	500	HIS	-	expression tag	UNP P20815
F	501	HIS	-	expression tag	UNP P20815
G	22	MET	-	initiating methionine	UNP P20815
G	23	ALA	-	expression tag	UNP P20815
G	498	HIS	-	expression tag	UNP P20815
G	499	HIS	-	expression tag	UNP P20815
G	500	HIS	-	expression tag	UNP P20815
G	501	HIS	-	expression tag	UNP P20815
H	22	MET	-	initiating methionine	UNP P20815
H	23	ALA	-	expression tag	UNP P20815
H	498	HIS	-	expression tag	UNP P20815
H	499	HIS	-	expression tag	UNP P20815
H	500	HIS	-	expression tag	UNP P20815

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
H	501	HIS	-	expression tag	UNP P20815
I	22	MET	-	initiating methionine	UNP P20815
I	23	ALA	-	expression tag	UNP P20815
I	498	HIS	-	expression tag	UNP P20815
I	499	HIS	-	expression tag	UNP P20815
I	500	HIS	-	expression tag	UNP P20815
I	501	HIS	-	expression tag	UNP P20815
J	22	MET	-	initiating methionine	UNP P20815
J	23	ALA	-	expression tag	UNP P20815
J	498	HIS	-	expression tag	UNP P20815
J	499	HIS	-	expression tag	UNP P20815
J	500	HIS	-	expression tag	UNP P20815
J	501	HIS	-	expression tag	UNP P20815
K	22	MET	-	initiating methionine	UNP P20815
K	23	ALA	-	expression tag	UNP P20815
K	498	HIS	-	expression tag	UNP P20815
K	499	HIS	-	expression tag	UNP P20815
K	500	HIS	-	expression tag	UNP P20815
K	501	HIS	-	expression tag	UNP P20815
L	22	MET	-	initiating methionine	UNP P20815
L	23	ALA	-	expression tag	UNP P20815
L	498	HIS	-	expression tag	UNP P20815
L	499	HIS	-	expression tag	UNP P20815
L	500	HIS	-	expression tag	UNP P20815
L	501	HIS	-	expression tag	UNP P20815

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0
2	C	1	43	34	1	4	4	0	0
2	D	1	43	34	1	4	4	0	0
2	E	1	43	34	1	4	4	0	0
2	F	1	43	34	1	4	4	0	0
2	G	1	43	34	1	4	4	0	0
2	H	1	43	34	1	4	4	0	0
2	I	1	43	34	1	4	4	0	0
2	J	1	43	34	1	4	4	0	0
2	K	1	43	34	1	4	4	0	0
2	L	1	43	34	1	4	4	0	0


- Molecule 3 is RITONAVIR (three-letter code: RIT) (formula: $C_{37}H_{48}N_6O_5S_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	S			
3	A	1	Total	50	37	6	5	2	0	0
3	B	1	Total	50	37	6	5	2	0	0
3	H	1	Total	50	37	6	5	2	0	0






- Molecule 1: Cytochrome P450 3A5




- Molecule 1: Cytochrome P450 3A5




- Molecule 1: Cytochrome P450 3A5





- Molecule 1: Cytochrome P450 3A5







● Molecule 1: Cytochrome P450 3A5



● Molecule 1: Cytochrome P450 3A5



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	148.99Å 198.38Å 234.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.16 – 2.91 39.16 – 2.91	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.16-2.91) 99.7 (39.16-2.91)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.90Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.214 , 0.258 0.214 , 0.258	Depositor DCC
R_{free} test set	7543 reflections (4.96%)	DCC
Wilson B-factor (Å ²)	70.7	Xtrriage
Anisotropy	0.364	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	44740	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, RIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/3803	0.42	0/5148
1	B	0.26	0/3767	0.42	0/5101
1	C	0.26	0/3864	0.43	0/5230
1	D	0.25	0/3775	0.42	0/5112
1	E	0.25	0/3625	0.42	0/4912
1	F	0.25	0/3749	0.41	0/5077
1	G	0.26	0/3737	0.41	0/5060
1	H	0.25	0/3775	0.41	0/5112
1	I	0.26	0/3822	0.42	0/5175
1	J	0.25	0/3767	0.41	0/5101
1	K	0.25	0/3769	0.41	0/5104
1	L	0.24	0/3676	0.41	0/4980
All	All	0.25	0/45129	0.42	0/61112

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3714	0	3810	32	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3679	0	3766	56	0
1	C	3773	0	3868	33	0
1	D	3687	0	3770	41	0
1	E	3540	0	3618	36	0
1	F	3661	0	3746	42	0
1	G	3651	0	3737	35	0
1	H	3687	0	3770	41	0
1	I	3732	0	3819	34	0
1	J	3679	0	3766	42	0
1	K	3681	0	3765	34	0
1	L	3590	0	3674	36	0
2	A	43	0	30	3	0
2	B	43	0	30	3	0
2	C	43	0	30	2	0
2	D	43	0	30	3	0
2	E	43	0	30	5	0
2	F	43	0	30	3	0
2	G	43	0	30	2	0
2	H	43	0	30	6	0
2	I	43	0	30	2	0
2	J	43	0	30	2	0
2	K	43	0	30	2	0
2	L	43	0	30	3	0
3	A	50	0	48	5	0
3	B	50	0	48	16	0
3	H	50	0	48	11	0
All	All	44740	0	45613	485	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 485 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ARG:NH1	3:B:602:RIT:H683	1.96	0.81
3:H:602:RIT:H141	3:H:602:RIT:H48	1.61	0.81
1:B:106:ARG:NH1	3:B:602:RIT:C68	2.46	0.79
1:A:38:ILE:HD11	1:A:386:VAL:HG21	1.64	0.79
1:B:117:ALA:HB1	1:B:301:ILE:HG13	1.67	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	458/480 (95%)	441 (96%)	17 (4%)	0	100	100
1	B	452/480 (94%)	430 (95%)	21 (5%)	1 (0%)	51	81
1	C	467/480 (97%)	449 (96%)	17 (4%)	1 (0%)	51	81
1	D	453/480 (94%)	432 (95%)	21 (5%)	0	100	100
1	E	431/480 (90%)	412 (96%)	18 (4%)	1 (0%)	51	81
1	F	450/480 (94%)	431 (96%)	19 (4%)	0	100	100
1	G	447/480 (93%)	432 (97%)	15 (3%)	0	100	100
1	H	453/480 (94%)	436 (96%)	17 (4%)	0	100	100
1	I	460/480 (96%)	441 (96%)	19 (4%)	0	100	100
1	J	452/480 (94%)	434 (96%)	18 (4%)	0	100	100
1	K	452/480 (94%)	432 (96%)	20 (4%)	0	100	100
1	L	440/480 (92%)	424 (96%)	16 (4%)	0	100	100
All	All	5415/5760 (94%)	5194 (96%)	218 (4%)	3 (0%)	55	85

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	424	ASP
1	B	383	ILE
1	E	353	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	419/434 (96%)	415 (99%)	4 (1%)	80	94
1	B	415/434 (96%)	412 (99%)	3 (1%)	87	96
1	C	426/434 (98%)	420 (99%)	6 (1%)	71	91
1	D	416/434 (96%)	412 (99%)	4 (1%)	80	94
1	E	399/434 (92%)	391 (98%)	8 (2%)	60	86
1	F	413/434 (95%)	412 (100%)	1 (0%)	94	98
1	G	412/434 (95%)	411 (100%)	1 (0%)	94	98
1	H	416/434 (96%)	412 (99%)	4 (1%)	80	94
1	I	421/434 (97%)	419 (100%)	2 (0%)	91	97
1	J	415/434 (96%)	412 (99%)	3 (1%)	87	96
1	K	415/434 (96%)	414 (100%)	1 (0%)	94	98
1	L	405/434 (93%)	405 (100%)	0	100	100
All	All	4972/5208 (96%)	4935 (99%)	37 (1%)	87	96

5 of 37 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	482	LEU
1	E	105	ARG
1	J	146	PHE
1	E	54	ARG
1	E	57	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	460	GLN
1	H	265	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	601	1,3	28,50,50	1.78	4 (14%)	17,82,82	1.51	3 (17%)
3	RIT	A	602	2	48,53,53	0.50	1 (2%)	50,71,71	0.82	2 (4%)
2	HEM	B	601	1,3	28,50,50	1.80	4 (14%)	17,82,82	1.51	4 (23%)
3	RIT	B	602	2	48,53,53	0.50	1 (2%)	50,71,71	0.57	0
2	HEM	C	601	1	28,50,50	1.77	4 (14%)	17,82,82	1.44	1 (5%)
2	HEM	D	601	1	28,50,50	1.79	4 (14%)	17,82,82	1.53	5 (29%)
2	HEM	E	601	1	28,50,50	1.76	4 (14%)	17,82,82	1.54	3 (17%)
2	HEM	F	601	1	28,50,50	1.76	4 (14%)	17,82,82	1.51	4 (23%)
2	HEM	G	601	1	28,50,50	1.77	4 (14%)	17,82,82	1.48	2 (11%)
2	HEM	H	601	1,3	28,50,50	1.83	4 (14%)	17,82,82	1.99	6 (35%)
3	RIT	H	602	2	48,53,53	0.54	1 (2%)	50,71,71	0.98	3 (6%)
2	HEM	I	601	1	28,50,50	1.77	4 (14%)	17,82,82	1.46	2 (11%)
2	HEM	J	601	1	28,50,50	1.76	4 (14%)	17,82,82	1.51	2 (11%)
2	HEM	K	601	1	28,50,50	1.75	4 (14%)	17,82,82	1.54	3 (17%)
2	HEM	L	601	1	28,50,50	1.75	4 (14%)	17,82,82	1.53	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	601	1,3	-	0/6/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RIT	A	602	2	-	0/49/53/53	0/4/4/4
2	HEM	B	601	1,3	-	0/6/54/54	0/0/8/8
3	RIT	B	602	2	-	0/49/53/53	0/4/4/4
2	HEM	C	601	1	-	0/6/54/54	0/0/8/8
2	HEM	D	601	1	-	0/6/54/54	0/0/8/8
2	HEM	E	601	1	-	0/6/54/54	0/0/8/8
2	HEM	F	601	1	-	0/6/54/54	0/0/8/8
2	HEM	G	601	1	-	0/6/54/54	0/0/8/8
2	HEM	H	601	1,3	-	0/6/54/54	0/0/8/8
3	RIT	H	602	2	-	0/49/53/53	0/4/4/4
2	HEM	I	601	1	-	0/6/54/54	0/0/8/8
2	HEM	J	601	1	-	0/6/54/54	0/0/8/8
2	HEM	K	601	1	-	0/6/54/54	0/0/8/8
2	HEM	L	601	1	-	0/6/54/54	0/0/8/8

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	601	HEM	C3B-C2B	-4.22	1.34	1.40
2	H	601	HEM	C3C-C2C	-4.05	1.35	1.40
2	A	601	HEM	C3B-C2B	-3.96	1.35	1.40
2	B	601	HEM	C3B-C2B	-3.96	1.35	1.40
2	C	601	HEM	C3C-C2C	-3.95	1.35	1.40

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	601	HEM	CAA-CBA-CGA	-5.64	103.02	112.66
3	A	602	RIT	C77-C75-N74	-3.68	107.33	113.72
3	H	602	RIT	C77-C75-N74	-3.45	107.72	113.72
3	H	602	RIT	C26-C12-C13	-3.29	105.81	111.74
2	B	601	HEM	CAA-CBA-CGA	-2.50	108.39	112.66

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

15 monomers are involved in 63 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	HEM	3	0
3	A	602	RIT	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	HEM	3	0
3	B	602	RIT	16	0
2	C	601	HEM	2	0
2	D	601	HEM	3	0
2	E	601	HEM	5	0
2	F	601	HEM	3	0
2	G	601	HEM	2	0
2	H	601	HEM	6	0
3	H	602	RIT	11	0
2	I	601	HEM	2	0
2	J	601	HEM	2	0
2	K	601	HEM	2	0
2	L	601	HEM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	462/480 (96%)	-0.03	12 (2%) 56 52	31, 51, 83, 109	0
1	B	458/480 (95%)	0.26	18 (3%) 40 36	37, 63, 94, 129	0
1	C	469/480 (97%)	-0.03	11 (2%) 61 58	27, 45, 77, 111	0
1	D	459/480 (95%)	0.10	11 (2%) 59 56	35, 57, 85, 123	0
1	E	441/480 (91%)	0.96	67 (15%) 2 2	45, 100, 129, 141	0
1	F	456/480 (95%)	0.58	37 (8%) 13 9	55, 79, 111, 128	0
1	G	455/480 (94%)	0.02	5 (1%) 80 80	33, 56, 86, 109	0
1	H	459/480 (95%)	0.08	9 (1%) 65 63	40, 60, 84, 104	0
1	I	464/480 (96%)	0.09	11 (2%) 59 56	36, 54, 82, 123	0
1	J	458/480 (95%)	0.30	24 (5%) 28 24	31, 65, 94, 140	0
1	K	458/480 (95%)	0.09	12 (2%) 56 52	39, 60, 88, 133	0
1	L	448/480 (93%)	1.11	93 (20%) 1 1	64, 104, 135, 143	0
All	All	5487/5760 (95%)	0.29	310 (5%) 25 21	27, 63, 117, 143	0

The worst 5 of 310 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	257	LYS	6.2
1	L	377	CYS	6.2
1	E	338	VAL	6.1
1	J	265	GLN	6.0
1	E	489	VAL	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	RIT	B	602	50/50	0.84	0.32	2.50	55,90,108,129	0
3	RIT	A	602	50/50	0.85	0.33	2.19	39,79,107,111	0
3	RIT	H	602	50/50	0.84	0.31	1.89	53,83,106,113	0
2	HEM	D	601	43/43	0.97	0.28	1.18	34,44,50,56	0
2	HEM	G	601	43/43	0.97	0.24	1.15	26,36,41,68	0
2	HEM	F	601	43/43	0.95	0.26	1.13	37,46,63,65	0
2	HEM	I	601	43/43	0.97	0.26	0.88	30,37,42,48	0
2	HEM	J	601	43/43	0.97	0.28	0.80	38,46,49,56	0
2	HEM	C	601	43/43	0.98	0.23	0.70	22,31,37,39	0
2	HEM	H	601	43/43	0.97	0.24	0.54	42,50,54,56	0
2	HEM	A	601	43/43	0.98	0.22	0.40	30,36,41,44	0
2	HEM	E	601	43/43	0.93	0.31	0.38	66,79,92,100	0
2	HEM	B	601	43/43	0.97	0.24	0.31	37,46,55,60	0
2	HEM	L	601	43/43	0.94	0.28	0.31	69,78,92,97	0
2	HEM	K	601	43/43	0.97	0.22	0.29	37,44,50,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.