



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:03 pm GMT

PDB ID : 2VGO  
Title : CRYSTAL STRUCTURE OF AURORA B KINASE IN COMPLEX WITH REVERSINE INHIBITOR  
Authors : D'Alise, A.M.; Amabile, G.; Iovino, M.; Di Giorgio, F.P.; Bartiromo, M.; Sessa, F.; Villa, F.; Musacchio, A.; Cortese, R.  
Deposited on : 2007-11-15  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	<div> <div>6%</div> <div>84%</div> <div>10%</div> <div>5%</div> </div>
1	B	284	<div> <div>7%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>
2	C	44	<div> <div>11%</div> <div>86%</div> <div>5%</div> <div>7%</div> </div>
2	D	44	<div> <div>36%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	AD5	A	1357	-	-	-	X
3	AD5	B	1359	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5956 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE 12-A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	P	S	0	2	0
			2258	1449	404	391	1	13			
1	B	278	Total	C	N	O	P	S	0	2	0
			2324	1488	418	403	1	14			

- Molecule 2 is a protein called INNER CENTROMERE PROTEIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	41	Total	C	N	O	S	0	1	0
			340	220	55	64	1			
2	D	43	Total	C	N	O	S	0	0	0
			350	225	58	66	1			

- Molecule 3 is N 6 -CYCLOHEXYL-N 2 -(4-MORPHOLIN-4-YLPHENYL)-9H-PURINE-2, 6-DIAMINE (three-letter code: AD5) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			29	21	7	1		
3	B	1	Total	C	N	O	0	0
			29	21	7	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	305	Total	O	0	0
			305	305		
4	B	261	Total	O	0	0
			261	261		
4	C	28	Total	O	0	0
			28	28		
4	D	32	Total	O	0	0
			32	32		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SERINE/THREONINE-PROTEIN KINASE 12-A




- Molecule 1: SERINE/THREONINE-PROTEIN KINASE 12-A



- Molecule 2: INNER CENTROMERE PROTEIN A



- Molecule 2: INNER CENTROMERE PROTEIN A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.74Å 67.60Å 116.63Å 90.00° 96.59° 90.00°	Depositor
Resolution (Å)	116.25 – 1.70 24.11 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (116.25-1.70) 96.1 (24.11-1.70)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.31 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.197 , 0.229 0.196 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	17.7	Xtriage
Anisotropy	0.854	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5956	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, AD5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/2308	0.54	0/3106
1	B	0.37	0/2374	0.52	0/3194
2	C	0.32	0/349	0.54	0/474
2	D	0.35	0/358	0.50	0/485
All	All	0.37	0/5389	0.53	0/7259

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	0	2261	24	0
1	B	2324	0	2334	18	0
2	C	340	0	335	5	0
2	D	350	0	347	6	0
3	A	29	0	27	1	0
3	B	29	0	27	2	0
4	A	305	0	0	4	0
4	B	261	0	0	5	0
4	C	28	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	32	0	0	1	0
All	All	5956	0	5331	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:823:ARG:HG3	2:C:823:ARG:HH11	0.94	1.09
1:A:112:GLU:HG3	4:A:2022:HOH:O	1.68	0.93
2:C:823:ARG:HG3	2:C:823:ARG:NH1	1.73	0.87
1:A:211:LYS:HD3	1:A:242:PRO:HA	1.58	0.86
1:A:150:HIS:HD2	1:A:152:ASN:H	1.23	0.84
3:B:1359:AD5:N1	3:B:1359:AD5:HAA	2.00	0.77
1:A:143:GLU:HG2	2:D:833:LEU:CD1	2.15	0.77
1:B:150:HIS:HD2	1:B:152:ASN:H	1.34	0.76
1:A:150:HIS:CD2	1:A:152:ASN:H	2.03	0.76
1:B:258:PRO:HG2	4:B:2233:HOH:O	1.86	0.74
1:A:87:LYS:HB3	4:A:2004:HOH:O	1.90	0.71
1:B:122:LYS:HE2	1:B:168:LEU:HD12	1.74	0.70
1:B:326:TYR:HA	4:B:2233:HOH:O	1.92	0.69
1:A:152:ASN:HD21	1:A:349:ARG:HH21	1.41	0.67
1:A:143:GLU:HG2	2:D:833:LEU:HD12	1.76	0.66
1:B:150:HIS:CD2	1:B:152:ASN:H	2.13	0.65
1:A:143:GLU:OE1	1:A:143:GLU:HA	1.97	0.65
3:A:1357:AD5:N1	3:A:1357:AD5:HAA	2.14	0.62
1:A:237:TRP:CD2	1:A:250:CYS:HB2	2.36	0.61
1:A:139:ARG:HG2	2:D:833:LEU:HD13	1.83	0.60
1:A:143:GLU:O	1:A:147:HIS:HD2	1.87	0.57
2:C:823:ARG:CG	2:C:823:ARG:HH11	1.88	0.55
1:A:193:SER:HB3	4:A:2133:HOH:O	2.05	0.55
1:A:152:ASN:ND2	1:A:349:ARG:HH21	2.04	0.54
2:D:817:LYS:HG3	4:D:2014:HOH:O	2.06	0.54
1:A:105:GLY:HA3	1:A:123:VAL:O	2.07	0.53
1:B:126:LYS:NZ	1:B:161:HIS:HD2	2.06	0.53
1:B:230:LEU:HD22	4:B:2106:HOH:O	2.10	0.51
1:B:272:LEU:HD21	1:B:334:LEU:HG	1.91	0.51
1:B:152:ASN:HD21	1:B:349:ARG:HH21	1.59	0.51
3:B:1359:AD5:N1	3:B:1359:AD5:CAA	2.64	0.50
2:C:823:ARG:CG	2:C:823:ARG:NH1	2.58	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:GLU:HB2	4:A:2030:HOH:O	2.11	0.50
1:B:89:THR:HG23	1:B:91:ASP:H	1.78	0.49
2:D:823:ARG:HA	2:D:827:THR:HG23	1.95	0.49
1:A:143:GLU:HG2	2:D:833:LEU:HD11	1.95	0.47
1:B:304:VAL:HG22	4:B:2233:HOH:O	2.14	0.47
1:A:277:VAL:HG13	1:A:288:PRO:HD2	1.99	0.45
1:A:90:ILE:HD11	1:A:95:ILE:HD11	2.00	0.44
1:B:126:LYS:HZ3	1:B:161:HIS:HD2	1.64	0.44
1:B:277:VAL:HG13	1:B:288:PRO:HD2	2.00	0.44
1:A:246:ARG:HD2	1:A:248:TPO:O2P	2.17	0.43
1:A:177[B]:GLU:OE1	1:A:179:TYR:HE2	2.00	0.43
1:B:101:LYS:HG2	1:B:101:LYS:O	2.18	0.43
1:A:143:GLU:O	1:A:147:HIS:CD2	2.70	0.43
1:B:306:LEU:HA	4:B:2186:HOH:O	2.17	0.43
1:B:309:PRO:HA	1:B:310:PRO:HD3	1.92	0.41
1:A:309:PRO:HA	1:A:310:PRO:HD3	1.87	0.41
1:A:150:HIS:HD2	1:A:152:ASN:N	2.03	0.41
1:B:246:ARG:HB3	1:B:248:TPO:O2P	2.21	0.40
1:B:135:GLU:HG3	2:C:837:PHE:CD2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/284 (94%)	259 (97%)	9 (3%)	0	100	100
1	B	275/284 (97%)	266 (97%)	9 (3%)	0	100	100
2	C	40/44 (91%)	39 (98%)	1 (2%)	0	100	100
2	D	41/44 (93%)	38 (93%)	3 (7%)	0	100	100
All	All	624/656 (95%)	602 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/255 (96%)	238 (98%)	6 (2%)	53	33
1	B	252/255 (99%)	246 (98%)	6 (2%)	54	35
2	C	37/39 (95%)	35 (95%)	2 (5%)	26	9
2	D	38/39 (97%)	36 (95%)	2 (5%)	26	9
All	All	571/588 (97%)	555 (97%)	16 (3%)	49	28

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	LYS
1	A	112	GLU
1	A	143	GLU
1	A	294	HIS
1	A	298	HIS
1	A	299	ARG
1	B	86	ARG
1	B	101	LYS
1	B	103	LYS
1	B	128	GLN
1	B	138	LEU
1	B	163	ARG
2	C	806	LEU
2	C	823	ARG
2	D	814	GLN
2	D	824	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	115	ASN
1	A	128	GLN
1	A	136	HIS
1	A	147	HIS
1	A	150	HIS
1	A	152	ASN
1	B	128	GLN
1	B	137	GLN
1	B	147	HIS
1	B	150	HIS
1	B	152	ASN
1	B	161	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	A	248	1	9,10,11	0.67	0	10,14,16	1.09	0
1	TPO	B	248	1	9,10,11	0.59	0	10,14,16	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	248	1	-	0/8/11/13	0/0/0/0
1	TPO	B	248	1	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	248	TPO	1	0
1	B	248	TPO	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	AD5	A	1357	-	30,33,33	1.14	3 (10%)	36,45,45	1.60	3 (8%)
3	AD5	B	1359	-	30,33,33	1.33	6 (20%)	36,45,45	1.65	5 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AD5	A	1357	-	-	0/11/28/28	0/5/5/5
3	AD5	B	1359	-	-	1/11/28/28	0/5/5/5

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1357	AD5	C5-C4	-2.97	1.33	1.40
3	B	1359	AD5	C5-C4	-2.89	1.34	1.40
3	A	1357	AD5	CAV-N2	-2.32	1.35	1.40
3	B	1359	AD5	CAV-N2	-2.24	1.36	1.40
3	A	1357	AD5	CAN-NBC	2.06	1.49	1.46
3	B	1359	AD5	CAM-NBC	2.13	1.50	1.46
3	B	1359	AD5	CAN-NBC	2.30	1.50	1.46
3	B	1359	AD5	CAN-CAJ	2.44	1.60	1.50
3	B	1359	AD5	CAM-CAI	2.58	1.60	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1357	AD5	N3-C2-N1	-4.91	118.81	126.23
3	B	1359	AD5	N3-C2-N1	-4.79	118.99	126.23
3	B	1359	AD5	OAU-CAJ-CAN	-2.63	105.94	111.83
3	B	1359	AD5	CAV-N2-C2	-2.39	122.96	129.17
3	B	1359	AD5	CAN-NBC-CAM	3.76	119.53	111.57
3	A	1357	AD5	CAN-NBC-CAM	4.25	120.57	111.57
3	B	1359	AD5	C2-N3-C4	4.59	120.35	115.11
3	A	1357	AD5	C2-N3-C4	4.82	120.61	115.11

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1359	AD5	N1-C6-N6-CBB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1357	AD5	1	0
3	B	1359	AD5	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/284 (94%)	0.54	17 (6%) 21 24	10, 17, 31, 39	0
1	B	277/284 (97%)	0.41	21 (7%) 15 17	10, 19, 37, 48	0
2	C	41/44 (93%)	0.73	5 (12%) 5 6	24, 27, 37, 41	0
2	D	43/44 (97%)	1.61	16 (37%) 0 0	21, 37, 42, 43	0
All	All	629/656 (95%)	0.57	59 (9%) 9 11	10, 20, 39, 48	0

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	104	PHE	7.3
1	A	244	LEU	7.1
1	A	103	LYS	5.4
1	B	136	HIS	5.3
1	A	243	SER	5.2
1	B	357	SER	5.1
1	B	163	ARG	4.6
2	D	804	GLY	4.5
1	A	102	GLY	4.4
1	A	104	PHE	4.2
2	D	827	THR	4.0
1	A	355	TYR	3.9
1	B	78	THR	3.9
2	D	803	SER	3.8
1	B	355	TYR	3.6
2	D	822	ASP	3.6
1	A	105	GLY	3.5
2	C	837	PHE	3.5
1	B	179	TYR	3.5
1	B	129	LEU	3.3
1	B	134	VAL	3.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	D	800	ALA	3.1
2	D	805	ASN	3.1
1	B	130	GLU	3.1
2	C	834	GLU	3.1
1	A	114	GLN	3.0
1	B	127	SER	3.0
2	D	823	ARG	2.9
1	A	136	HIS	2.9
1	B	183	GLN	2.9
1	B	128	GLN	2.9
1	B	79	ALA	2.9
1	A	278	LEU	2.8
1	A	87	LYS	2.7
2	D	837	PHE	2.7
2	D	834	GLU	2.6
1	B	164	LYS	2.6
2	D	828	ILE	2.5
2	D	798	ILE	2.5
2	D	832	LYS	2.4
2	D	838	ASN	2.4
1	A	101	LYS	2.4
1	A	299	ARG	2.3
2	C	797	PRO	2.3
1	B	139	ARG	2.3
2	C	798	ILE	2.3
1	B	103	LYS	2.3
1	B	131	LYS	2.3
1	A	245	ARG	2.2
1	B	356	GLN	2.2
1	B	102	GLY	2.2
1	B	101	LYS	2.2
2	D	824	MET	2.1
2	D	826	GLY	2.1
2	D	829	ASP	2.1
1	A	166	ILE	2.1
1	A	118	ILE	2.0
2	C	813	GLN	2.0
1	A	179	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	B	248	11/12	0.98	0.05	-	18,19,21,21	0
1	TPO	A	248	11/12	0.98	0.07	-	18,20,21,21	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	AD5	B	1359	29/29	0.57	0.28	2.81	48,49,52,52	0
3	AD5	A	1357	29/29	0.68	0.26	2.64	36,39,42,42	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.