



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:56 pm GMT

PDB ID : 1VMA
Title : Crystal structure of Cell division protein ftsY (TM0570) from *Thermotoga maritima* at 1.60 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2004-09-15
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>12%</div> <div>88%</div> <div>8%</div> <div>.</div> </div>
1	B	306	<div> <div>17%</div> <div>83%</div> <div>13%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	295	-	-	-	X
3	CIT	B	295	-	-	X	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cell division protein FtsY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	294	Total	C	N	O	S	0	7	0
			2235	1435	366	429	5			
1	B	294	Total	C	N	O	S	0	5	0
			2223	1428	368	421	6			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP Q9WZ40
A	-10	GLY	-	EXPRESSION TAG	UNP Q9WZ40
A	-9	SER	-	EXPRESSION TAG	UNP Q9WZ40
A	-8	ASP	-	EXPRESSION TAG	UNP Q9WZ40
A	-7	LYS	-	EXPRESSION TAG	UNP Q9WZ40
A	-6	ILE	-	EXPRESSION TAG	UNP Q9WZ40
A	-5	HIS	-	EXPRESSION TAG	UNP Q9WZ40
A	-4	HIS	-	EXPRESSION TAG	UNP Q9WZ40
A	-3	HIS	-	EXPRESSION TAG	UNP Q9WZ40
A	-2	HIS	-	EXPRESSION TAG	UNP Q9WZ40
A	-1	HIS	-	EXPRESSION TAG	UNP Q9WZ40
A	0	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	-11	MET	-	EXPRESSION TAG	UNP Q9WZ40
B	-10	GLY	-	EXPRESSION TAG	UNP Q9WZ40
B	-9	SER	-	EXPRESSION TAG	UNP Q9WZ40
B	-8	ASP	-	EXPRESSION TAG	UNP Q9WZ40
B	-7	LYS	-	EXPRESSION TAG	UNP Q9WZ40
B	-6	ILE	-	EXPRESSION TAG	UNP Q9WZ40
B	-5	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	-4	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	-3	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	-2	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	-1	HIS	-	EXPRESSION TAG	UNP Q9WZ40
B	0	HIS	-	EXPRESSION TAG	UNP Q9WZ40

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			13	6	7		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	219	Total	O	0	0
			219	219		
4	B	175	Total	O	0	0
			175	175		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cell division protein FtsY



- Molecule 1: cell division protein FtsY



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.45Å 48.10Å 90.79Å 90.00° 107.94° 90.00°	Depositor
Resolution (Å)	24.88 – 1.60 24.88 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.5 (24.88-1.60) 98.5 (24.88-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.2.0001	Depositor
R, R_{free}	0.207 , 0.253 0.220 , 0.263	Depositor DCC
R_{free} test set	4218 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	19.0	Xtriage
Anisotropy	0.253	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4866	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.93 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2171e-05.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2294 (0.0%)	0.92	3/3101 (0.1%)
1	B	0.71	0/2272	0.89	8/3074 (0.3%)
All	All	0.79	1/4566 (0.0%)	0.90	11/6175 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	285	GLU	CD-OE1	5.10	1.31	1.25

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ASP	CB-CG-OD1	7.98	125.48	118.30
1	B	32	ASP	CB-CG-OD1	6.25	123.93	118.30
1	B	47	ASP	CB-CG-OD2	5.89	123.60	118.30
1	B	82	ASP	CB-CG-OD2	5.59	123.33	118.30
1	B	176	ASP	CB-CG-OD2	5.58	123.33	118.30
1	B	201	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	B	118	ASP	CB-CG-OD2	5.40	123.16	118.30
1	B	5	ASP	CB-CG-OD2	5.28	123.05	118.30
1	B	21	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	66	ASP	CB-CG-OD1	5.13	122.92	118.30
1	A	132	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2235	0	2278	23	0
1	B	2223	0	2275	27	0
2	A	1	0	0	0	0
3	B	13	0	5	6	0
4	A	219	0	0	1	0
4	B	175	0	0	4	0
All	All	4866	0	4558	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:PHE:CE1	1:A:272[A]:VAL:HG22	2.10	0.86
1:B:105:LYS:N	3:B:295:CIT:H21	1.96	0.81
1:B:75:ILE:HG23	1:B:257:LEU:HD21	1.66	0.78
1:B:189:LYS:HE2	1:B:232:ILE:HG23	1.66	0.78
1:B:105:LYS:H	3:B:295:CIT:H21	1.53	0.73
1:A:124:VAL:HG23	1:A:175[A]:LYS:HG2	1.77	0.65
1:B:12:GLN:O	1:B:16:GLU:HG3	1.99	0.63
1:A:124:VAL:HG23	1:A:175[B]:LYS:HG2	1.83	0.61
1:A:146:VAL:CG1	1:A:146:VAL:O	2.49	0.61
1:B:278:ASP:C	1:B:278:ASP:OD1	2.41	0.59
1:A:86:ASN:CG	1:A:213[B]:GLU:OE2	2.40	0.59
1:A:257:LEU:HD12	1:A:257:LEU:N	2.19	0.58
1:A:100:VAL:HG23	1:A:103:THR:HG23	1.86	0.56
1:B:253:GLY:O	1:B:256[A]:THR:HG22	2.05	0.55
1:B:105:LYS:HB2	3:B:295:CIT:C1	2.36	0.55
1:A:114:LYS:HA	1:A:146:VAL:HG13	1.89	0.54
1:A:146:VAL:HG12	1:A:146:VAL:O	2.07	0.53
1:B:256[A]:THR:HG21	1:B:287:PHE:CZ	2.45	0.51
1:B:189:LYS:HE2	1:B:232:ILE:HD12	1.93	0.51
1:B:72:LYS:HG2	1:B:292:LEU:HD12	1.92	0.50
1:A:114:LYS:HA	1:A:146:VAL:CG1	2.42	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:272:VAL:O	1:B:278:ASP:HB2	2.12	0.49
1:A:122:SER:HB2	1:A:175[A]:LYS:CE	2.44	0.48
1:B:275:LYS:O	1:B:278:ASP:CG	2.53	0.48
1:A:122:SER:HB2	1:A:175[A]:LYS:HE2	1.96	0.47
1:A:86:ASN:N	1:A:213[B]:GLU:OE2	2.43	0.46
1:B:189:LYS:CE	1:B:232:ILE:HD12	2.45	0.46
1:B:140:LYS:O	1:B:144:GLU:HG3	2.15	0.46
1:B:104:GLY:HA2	3:B:295:CIT:O6	2.15	0.46
1:A:280:ARG:HD2	4:A:385:HOH:O	2.15	0.45
1:B:105:LYS:HB2	3:B:295:CIT:O1	2.17	0.45
1:A:6:PHE:CE1	1:A:272[A]:VAL:CG2	2.93	0.44
1:B:132:ARG:HD2	4:B:323:HOH:O	2.16	0.44
1:A:229:GLN:O	1:A:232:ILE:HB	2.17	0.44
1:B:185:ARG:HD2	4:B:460:HOH:O	2.16	0.44
1:B:284:PRO:HD2	4:B:359:HOH:O	2.18	0.44
1:A:257:LEU:N	1:A:257:LEU:CD1	2.81	0.43
1:B:139:LEU:HD12	1:B:139:LEU:HA	1.90	0.42
1:A:151:ILE:HG13	1:A:166:ALA:HA	2.01	0.42
1:B:12:GLN:NE2	1:B:16:GLU:OE2	2.53	0.42
1:A:257:LEU:HD12	1:A:257:LEU:H	1.85	0.41
1:B:86:ASN:CG	1:B:213:GLU:OE2	2.59	0.41
1:B:252:LYS:O	1:B:255:ILE:HG22	2.20	0.41
1:B:105:LYS:H	3:B:295:CIT:C2	2.27	0.41
1:B:105:LYS:HG2	1:B:217:VAL:HG21	2.02	0.41
1:A:257:LEU:H	1:A:257:LEU:CD1	2.34	0.41
1:A:76:LEU:HA	1:A:76:LEU:HD12	1.93	0.41
1:B:77:GLU:HB2	4:B:353:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/306 (98%)	290 (97%)	9 (3%)	0	100	100
1	B	297/306 (97%)	293 (99%)	4 (1%)	0	100	100
All	All	596/612 (97%)	583 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/253 (92%)	232 (99%)	2 (1%)	82	69
1	B	231/253 (91%)	223 (96%)	8 (4%)	41	15
All	All	465/506 (92%)	455 (98%)	10 (2%)	60	29

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LYS
1	A	81	PHE
1	B	27	LYS
1	B	81	PHE
1	B	153	HIS
1	B	189	LYS
1	B	234	LYS
1	B	261[A]	ARG
1	B	261[B]	ARG
1	B	278	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	CIT	B	295	-	3,12,12	1.64	0	3,17,17	3.70	3 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CIT	B	295	-	-	0/6/16/16	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	295	CIT	C3-C4-C5	-3.78	109.05	114.95
3	B	295	CIT	C4-C3-C2	2.75	116.60	109.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	295	CIT	C3-C2-C1	4.37	121.78	114.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	295	CIT	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/306 (96%)	0.91	38 (12%) 4 4	15, 22, 34, 53	0
1	B	294/306 (96%)	1.04	51 (17%) 2 1	16, 23, 38, 47	0
All	All	588/612 (96%)	0.97	89 (15%) 3 2	15, 23, 37, 53	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	186	LEU	9.1
1	B	186	LEU	8.6
1	B	64	ASP	8.5
1	A	187	HIS	7.9
1	B	276	ALA	6.8
1	A	64	ASP	6.2
1	A	188	THR	4.9
1	B	188	THR	4.6
1	B	187	HIS	4.5
1	B	81	PHE	4.0
1	B	179	ILE	3.9
1	B	31	LEU	3.8
1	B	146	VAL	3.8
1	A	80[A]	ASN	3.7
1	A	43	LEU	3.4
1	B	82	ASP	3.4
1	B	144	GLU	3.3
1	B	71	LEU	3.3
1	B	180	ILE	3.3
1	B	191	ASN	3.3
1	A	205	LYS	3.3
1	B	89	PRO	3.2
1	B	35	THR	3.2
1	B	57	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	172	ALA	3.1
1	A	179	ILE	3.1
1	A	180	ILE	3.1
1	B	95	ILE	3.0
1	A	131	PHE	3.0
1	B	97	VAL	3.0
1	A	189	LYS	2.9
1	B	142	TRP	2.9
1	A	215	LEU	2.8
1	B	79	LEU	2.8
1	A	112	LEU	2.8
1	B	215	LEU	2.8
1	B	216	LEU	2.8
1	A	109	CYS	2.8
1	A	203	VAL	2.7
1	B	28	GLY	2.7
1	A	97	VAL	2.7
1	B	147	GLY	2.7
1	B	277	GLU	2.7
1	A	141	ILE	2.7
1	B	172	ALA	2.7
1	B	189	LYS	2.7
1	B	242	ILE	2.6
1	B	291	LEU	2.6
1	A	276	ALA	2.6
1	B	171	LEU	2.6
1	A	144	GLU	2.6
1	A	46	ALA	2.5
1	B	273	GLY	2.5
1	A	243	ILE	2.5
1	B	98	VAL	2.5
1	A	95	ILE	2.5
1	B	133	ALA	2.4
1	B	278	ASP	2.4
1	B	109	CYS	2.4
1	A	146	VAL	2.4
1	B	190	LYS	2.4
1	B	65	GLY	2.4
1	B	34	GLU	2.4
1	B	24	LYS	2.3
1	B	32	ASP	2.3
1	A	292	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	257	LEU	2.3
1	A	42	LEU	2.2
1	A	278	ASP	2.2
1	A	190	LYS	2.2
1	A	48	VAL	2.1
1	A	178	VAL	2.1
1	B	139	LEU	2.1
1	A	56	ILE	2.1
1	A	217	VAL	2.1
1	A	31	LEU	2.1
1	A	169	HIS	2.1
1	A	170	ALA	2.1
1	B	26	LEU	2.1
1	B	2	GLY	2.1
1	B	75	ILE	2.1
1	B	267	ILE	2.1
1	B	182	THR	2.1
1	A	287	PHE	2.1
1	B	96	MET	2.1
1	A	277	GLU	2.0
1	B	275	LYS	2.0
1	A	237	VAL	2.0
1	B	48	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	CIT	B	295	13/13	0.76	0.27	5.63	40,46,53,56	0
2	MG	A	295	1/1	0.95	0.18	2.45	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.