



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:21 pm GMT

PDB ID : 3VQV  
Title : Crystal structure of the catalytic domain of pyrrolysyl-tRNA synthetase in complex with AMPPNP (re-refined)  
Authors : Yanagisawa, T.; Sumida, T.; Ishii, R.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2012-04-01  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	291	<div> <div>5%</div> <div> <div></div> <div>70%</div> <div>15%</div> <div>•</div> <div>11%</div> </div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2377 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyrrolysine-tRNA ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	1	0
			2098	1333	362	394	9			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	246	Total 246	O 246	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyrrolysine-tRNA ligase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.88Å 104.88Å 70.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.07 – 1.90 42.06 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.07-1.90) 99.8 (42.06-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.187 , 0.219 0.187 , 0.219	Depositor DCC
$R_{free}$ test set	1750 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 53.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.042 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2377	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	0/2142	0.87	7/2877 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	298	ARG	NE-CZ-NH1	10.98	125.79	120.30
1	A	298	ARG	NE-CZ-NH2	-8.62	115.99	120.30
1	A	227	ARG	NE-CZ-NH2	-8.49	116.05	120.30
1	A	227	ARG	NE-CZ-NH1	5.77	123.18	120.30
1	A	344	MET	CG-SD-CE	-5.21	91.86	100.20
1	A	227	ARG	CG-CD-NE	-5.18	100.92	111.80
1	A	397	LEU	CB-CG-CD1	5.10	119.67	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2098	0	2103	38	0
2	A	31	0	13	0	0
3	A	2	0	0	0	0
4	A	246	0	0	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2377	0	2116	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:414:ASP:N	4:A:845:HOH:O	1.77	1.00
1:A:279:ASP:O	1:A:282:THR:HG22	1.64	0.97
1:A:436:ASN:HD22	1:A:438:LYS:H	1.32	0.78
1:A:217:ARG:HG2	4:A:757:HOH:O	1.89	0.71
1:A:413:ILE:O	4:A:709:HOH:O	2.09	0.70
1:A:206:ASP:OD2	1:A:227:ARG:NH2	2.26	0.69
1:A:298:ARG:HD2	1:A:327:PRO:O	1.97	0.65
1:A:195:THR:O	1:A:199:GLU:HG3	1.98	0.63
1:A:373:ASP:O	1:A:392:HIS:HD2	1.82	0.62
1:A:252:ARG:HH11	1:A:252:ARG:HG3	1.64	0.62
1:A:212:SER:HB3	4:A:837:HOH:O	1.99	0.61
1:A:337:GLU:HG3	1:A:395:LEU:HD13	1.83	0.60
1:A:436:ASN:ND2	1:A:438:LYS:H	2.00	0.60
1:A:337:GLU:CG	1:A:395:LEU:HD13	2.34	0.58
1:A:413:ILE:CA	4:A:845:HOH:O	2.51	0.58
1:A:204:PRO:HD3	4:A:827:HOH:O	2.05	0.56
1:A:387:THR:HG22	4:A:798:HOH:O	2.09	0.53
1:A:313:ASP:HA	1:A:320:ILE:HD11	1.91	0.52
1:A:436:ASN:HD21	1:A:438:LYS:HG3	1.74	0.52
1:A:278:ILE:HG23	1:A:282:THR:HG21	1.92	0.51
1:A:337:GLU:HG3	1:A:395:LEU:CD1	2.40	0.51
1:A:275:ARG:HG3	1:A:410:GLU:HG2	1.97	0.47
1:A:284:LEU:HA	1:A:287:GLN:HE21	1.80	0.47
1:A:227:ARG:HA	1:A:227:ARG:HD3	1.66	0.46
1:A:301:LEU:HD13	1:A:328:CYS:SG	2.55	0.46
1:A:214:LYS:HD3	1:A:218:GLU:OE1	2.17	0.45
1:A:436:ASN:ND2	1:A:438:LYS:HG3	2.31	0.45
1:A:304:ASN:ND2	4:A:640:HOH:O	2.44	0.44
1:A:275:ARG:HG3	1:A:410:GLU:CG	2.48	0.44
1:A:271:GLU:H	1:A:271:GLU:CD	2.20	0.44
1:A:392:HIS:HE1	4:A:841:HOH:O	2.00	0.43
1:A:252:ARG:CG	1:A:252:ARG:HH11	2.30	0.43
1:A:345:LEU:O	1:A:421:GLY:HA2	2.19	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271:GLU:OE2	1:A:271:GLU:N	2.51	0.42
1:A:280:ASN:OD1	1:A:280:ASN:N	2.52	0.42
1:A:217:ARG:HA	1:A:217:ARG:HD2	1.97	0.41
1:A:238:GLU:HG3	4:A:630:HOH:O	2.19	0.41
1:A:313:ASP:HA	1:A:320:ILE:CD1	2.50	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	253/291 (87%)	245 (97%)	8 (3%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	232/259 (90%)	210 (90%)	22 (10%)	<b>10</b> <b>3</b>

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	190	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	219	LEU
1	A	229	LYS
1	A	231	LEU
1	A	252	ARG
1	A	257	ARG
1	A	270	LEU
1	A	271	GLU
1	A	273	ILE
1	A	275	ARG
1	A	280	ASN
1	A	293	LYS
1	A	297	LEU
1	A	298	ARG
1	A	301	LEU
1	A	337	GLU
1	A	386	ASP
1	A	387	THR
1	A	388	LEU
1	A	394	ASP
1	A	397	LEU
1	A	435	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	287	GLN
1	A	349	GLN
1	A	358	ASN
1	A	368	ASN
1	A	392	HIS
1	A	436	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ANP	A	501	3	29,33,33	3.45	10 (34%)	28,52,52	3.10	6 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	501	3	-	0/13/38/38	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	ANP	C2'-C1'	-4.49	1.46	1.53
2	A	501	ANP	PB-O2B	-2.96	1.48	1.56
2	A	501	ANP	C5-C4	-2.08	1.35	1.40
2	A	501	ANP	PG-O3G	2.66	1.64	1.56
2	A	501	ANP	PB-N3B	3.10	1.71	1.63
2	A	501	ANP	C2-N1	3.69	1.40	1.33
2	A	501	ANP	C2-N3	5.55	1.41	1.32
2	A	501	ANP	O4'-C1'	5.71	1.49	1.41
2	A	501	ANP	PG-N3B	6.25	1.79	1.63
2	A	501	ANP	PG-O1G	12.51	1.60	1.46

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ANP	N3-C2-N1	-13.99	116.68	128.86
2	A	501	ANP	O3G-PG-O1G	-3.66	104.12	113.41
2	A	501	ANP	O3A-PB-N3B	-2.65	99.25	106.59
2	A	501	ANP	C4'-O4'-C1'	-2.24	107.39	109.77
2	A	501	ANP	O1G-PG-N3B	-2.23	108.45	111.79
2	A	501	ANP	O2B-PB-O1B	4.08	118.36	109.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/291 (88%)	-0.08	14 (5%) 26 30	22, 35, 63, 83	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	ASP	4.8
1	A	379	ASP	3.8
1	A	213	GLY	3.7
1	A	282	THR	3.5
1	A	188	PRO	3.3
1	A	212	SER	3.2
1	A	280	ASN	3.1
1	A	205	LYS	2.9
1	A	207	GLU	2.8
1	A	293	LYS	2.6
1	A	286	LYS	2.5
1	A	206	ASP	2.1
1	A	378	GLY	2.1
1	A	385	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	ANP	A	501	31/31	0.99	0.06	-0.92	29,31,37,38	0
3	MG	A	502	1/1	0.98	0.03	-	36,36,36,36	0
3	MG	A	503	1/1	0.96	0.05	-	45,45,45,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.