



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 12, 2017 – 08:18 pm GMT

PDB ID : 1VRY
Title : Second and Third Transmembrane Domains of the Alpha-1 Subunit of Human Glycine Receptor
Authors : Ma, D.; Liu, Z.; Li, L.; Tang, P.; Xu, Y.
Deposited on : 2005-07-20

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	trunk28760
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	76	

2 Ensemble composition and analysis

This entry contains 20 models. Model 14 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:-1-A:16, A:27-A:55 (47)	0.20	14

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 5 clusters and 2 single-model clusters were found.

Cluster number	Models
1	11, 12, 14, 15, 16, 19
2	5, 7, 9, 18
3	4, 10, 13
4	8, 17, 20
5	2, 6
Single-model clusters	1; 3

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 954 atoms, of which 494 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Glycine receptor alpha-1 chain.

Mol	Chain	Residues	Atoms						Trace
1	A	61	Total	C	H	N	O	S	0
			954	303	494	75	81	1	

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	LEU	-	CLONING ARTIFACT	UNP P23415
A	11	LEU	MET	ENGINEERED	UNP P23415
A	35	LEU	MET	ENGINEERED	UNP P23415
A	58	LYS	-	CLONING ARTIFACT	UNP P23415
A	59	LYS	-	CLONING ARTIFACT	UNP P23415
A	60	LYS	-	CLONING ARTIFACT	UNP P23415
A	61	LYS	-	CLONING ARTIFACT	UNP P23415
A	62	HIS	-	CLONING ARTIFACT	UNP P23415
A	63	ARG	-	CLONING ARTIFACT	UNP P23415
A	64	LEU	-	CLONING ARTIFACT	UNP P23415
A	65	LEU	-	CLONING ARTIFACT	UNP P23415
A	66	GLU	-	CLONING ARTIFACT	UNP P23415
A	67	HIS	-	CLONING ARTIFACT	UNP P23415
A	68	HIS	-	CLONING ARTIFACT	UNP P23415
A	69	HIS	-	CLONING ARTIFACT	UNP P23415
A	70	HIS	-	CLONING ARTIFACT	UNP P23415
A	71	HIS	-	CLONING ARTIFACT	UNP P23415
A	72	HIS	-	CLONING ARTIFACT	UNP P23415

5 Refinement protocol and experimental data overview

The models were refined using the following method: *SIMULATED ANNEALING*.

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *TARGET FUNCTION, STRUCTURES WITH THE LOWEST ENERGY, STRUCTURES WITH THE LEAST RESTRAINT VIOLATIONS*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
XPLOR-NIH	refinement	2.9.6
XWINNMR	structure solution	2.6
NMRPIPE SGI6X	structure solution	
XPLOR-NIH	structure solution	2.9.6

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	359	380	380	50±6
All	All	7180	7600	7600	990

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 67.

5 of 141 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:THR:HA	1:A:9:LEU:HD12	0.89	1.43	5	18
1:A:9:LEU:HD13	1:A:10:THR:N	0.86	1.86	13	18
1:A:3:LEU:HD23	1:A:4:GLY:N	0.85	1.86	17	2
1:A:39:LEU:HD12	1:A:40:LEU:N	0.84	1.88	18	10
1:A:8:VAL:HG11	1:A:41:PHE:CZ	0.80	2.11	19	20

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	47/76 (62%)	44±1 (93±2%)	2±1 (3±2%)	2±0 (4±1%)	6	32
All	All	940/1520 (62%)	871 (93%)	30 (3%)	39 (4%)	6	32

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	32	ASP	20
1	A	1	VAL	17
1	A	37	VAL	2

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	39/66 (59%)	31±1 (80±3%)	8±1 (20±3%)	4	34
All	All	780/1320 (59%)	623 (80%)	157 (20%)	4	34

5 of 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	52	VAL	20
1	A	10	THR	20
1	A	48	GLU	20
1	A	7	THR	20
1	A	34	TRP	20

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided