



# Full wwPDB X-ray Structure Validation Report i

Feb 14, 2017 – 02:23 pm GMT

PDB ID : 2VRO  
Title : CRYSTAL STRUCTURE OF ALDEHYDE DEHYDROGENASE FROM BURKHOLDERIA XENOVORANS LB400  
Authors : Bains, J.; Boulanger, M.J.  
Deposited on : 2008-04-09  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NDP	A	1523	-	-	-	X
2	NDP	B	1531	-	-	-	X
4	P6G	B	1532	-	-	-	X

## 2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 8588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALDEHYDE DEHYDROGENASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	521	3838	2408	692	728	10	0	0	0
1	B	527	3885	2440	702	733	10	0	0	0

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0
2	B	1	48	21	7	17	3	0	0

- Molecule 3 is 2-{2-[2-2-(METHOXY-ETHOXY)-ETHOXY]-ETHOXY}-ETHANOL (three-letter code: ETE) (formula: C<sub>9</sub>H<sub>20</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total 14    9    5	0	0

- Molecule 4 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: C<sub>12</sub>H<sub>26</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total 19    12    7	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	329	Total O 329 329	0	0
5	B	407	Total O 407 407	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ALDEHYDE DEHYDROGENASE



- Molecule 1: ALDEHYDE DEHYDROGENASE



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.98 Å    67.69 Å    77.62 Å 111.18°    90.48°    113.27°	Depositor
Resolution (Å)	37.82 – 1.60 37.82 – 1.60	Depositor EDS
% Data completeness (in resolution range)	92.6 (37.82-1.60) 79.0 (37.82-1.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.48 (at 1.60 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
$R$ , $R_{free}$	0.207 , 0.238 0.205 , 0.237	Depositor DCC
$R_{free}$ test set	6128 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.4	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 37.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8588	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, P6G, ETE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	1/3908 (0.0%)	0.60	0/5321
1	B	0.55	0/3959	0.63	0/5393
All	All	0.54	1/7867 (0.0%)	0.61	0/10714

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	167	GLU	CB-CG	-5.59	1.41	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3838	0	3847	37	0
1	B	3885	0	3890	49	0
2	A	48	0	26	3	0
2	B	48	0	26	4	0
3	A	14	0	20	3	0
4	B	19	0	26	6	0
5	A	329	0	0	6	0
5	B	407	0	0	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8588	0	7835	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:189:TRP:CZ2	5:B:2005:HOH:O	2.01	1.12
1:B:189:TRP:HD1	5:B:2186:HOH:O	1.37	1.07
1:B:189:TRP:CD1	5:B:2186:HOH:O	2.12	1.01
1:A:163:TRP:O	1:A:167:GLU:HB2	1.62	0.98
1:B:463:HIS:CE1	1:B:465:ILE:HD11	2.07	0.89
1:B:126:ARG:NH2	5:B:2145:HOH:O	2.09	0.86
1:A:255:ASN:HD21	1:A:495:GLU:H	1.25	0.84
1:B:189:TRP:CH2	5:B:2005:HOH:O	2.20	0.83
2:A:1523:NDP:H52N	5:A:2322:HOH:O	1.78	0.81
1:B:189:TRP:HH2	5:B:2187:HOH:O	1.62	0.81
1:B:189:TRP:HZ3	1:B:192:GLN:OE1	1.63	0.80
1:B:189:TRP:CH2	5:B:2187:HOH:O	2.37	0.78
1:B:255:ASN:HD21	1:B:495:GLU:H	1.29	0.77
1:B:189:TRP:CE3	1:B:189:TRP:O	2.40	0.75
1:B:479:VAL:H	4:B:1532:P6G:H52	1.51	0.74
1:B:420:LEU:HD11	1:B:454:GLU:HG3	1.70	0.74
1:B:383:HIS:HE1	5:B:2303:HOH:O	1.71	0.73
1:B:420:LEU:CD1	1:B:454:GLU:HG3	2.23	0.68
1:B:527:HIS:HE1	5:B:2140:HOH:O	1.77	0.68
1:B:163:TRP:CZ2	1:B:167:GLU:HG3	2.28	0.68
1:B:381:ALA:O	1:B:383:HIS:HD2	1.78	0.66
1:A:294:GLN:HE22	1:A:339:VAL:H	1.45	0.65
1:A:479:VAL:H	3:A:1524:ETE:H252	1.62	0.64
1:B:158:PHE:CE2	2:B:1531:NDP:H4D	2.33	0.64
1:A:232:THR:HG23	1:A:257:GLU:HG3	1.79	0.64
1:A:508:ARG:CZ	5:A:2317:HOH:O	2.46	0.62
4:B:1532:P6G:H142	5:B:2407:HOH:O	1.99	0.61
1:B:356:GLU:HG3	5:B:2292:HOH:O	1.99	0.61
1:B:460:GLY:HA3	1:B:486:GLY:O	2.01	0.59
1:B:479:VAL:H	4:B:1532:P6G:C5	2.14	0.59
1:A:421:PRO:O	5:A:2275:HOH:O	2.16	0.58
1:A:381:ALA:O	1:A:383:HIS:HD2	1.87	0.58
1:B:189:TRP:CZ3	1:B:192:GLN:OE1	2.52	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:GLN:HE22	1:B:339:VAL:H	1.48	0.58
1:A:295:LYS:HD2	3:A:1524:ETE:H262	1.85	0.57
2:B:1531:NDP:O2A	2:B:1531:NDP:H8A	2.04	0.57
1:B:158:PHE:HE2	2:B:1531:NDP:H4D	1.70	0.56
1:A:439:SER:OG	1:A:476:HIS:HD2	1.88	0.56
1:A:224:ARG:HH11	1:A:224:ARG:HG3	1.70	0.56
1:A:233:GLY:O	1:A:258:ALA:HA	2.05	0.56
1:B:439:SER:OG	1:B:476:HIS:HD2	1.90	0.55
1:B:463:HIS:HE1	1:B:465:ILE:HD11	1.67	0.55
1:B:297:THR:O	1:B:476:HIS:HE1	1.90	0.54
1:B:261:LEU:HD13	1:B:490:ARG:HH11	1.71	0.54
1:A:248:VAL:HG23	1:A:249:GLN:HG3	1.89	0.54
1:A:508:ARG:NE	5:A:2317:HOH:O	2.41	0.53
1:A:224:ARG:HH11	1:A:224:ARG:CG	2.22	0.53
1:A:297:THR:O	1:A:476:HIS:HE1	1.92	0.52
1:A:522:LEU:HD13	5:A:2066:HOH:O	2.09	0.52
1:B:189:TRP:HA	1:B:189:TRP:HE3	1.74	0.52
1:B:381:ALA:O	1:B:383:HIS:CD2	2.61	0.52
1:B:158:PHE:CD1	1:B:339:VAL:HG21	2.45	0.51
1:A:255:ASN:HD21	1:A:495:GLU:N	2.03	0.51
1:A:460:GLY:HA3	1:A:486:GLY:O	2.11	0.51
1:B:189:TRP:CE3	1:B:189:TRP:HA	2.46	0.51
1:B:439:SER:CB	1:B:476:HIS:HD2	2.24	0.50
1:B:232:THR:HG23	1:B:257:GLU:HG3	1.94	0.50
4:B:1532:P6G:H82	5:B:2407:HOH:O	2.10	0.49
1:B:295:LYS:HD2	4:B:1532:P6G:H22	1.94	0.49
1:A:463:HIS:HE1	1:A:473:GLN:OE1	1.94	0.49
1:A:297:THR:O	1:A:476:HIS:CE1	2.65	0.49
1:B:463:HIS:HE1	1:B:473:GLN:OE1	1.96	0.49
1:B:297:THR:O	1:B:476:HIS:CE1	2.66	0.49
1:A:104:GLY:HA2	3:A:1524:ETE:H132	1.94	0.49
1:B:356:GLU:CG	5:B:2292:HOH:O	2.61	0.48
1:B:233:GLY:O	1:B:258:ALA:HA	2.13	0.48
1:A:439:SER:CB	1:A:476:HIS:HD2	2.27	0.47
1:A:234:SER:HB3	2:A:1523:NDP:H6N	1.96	0.47
1:A:467:PRO:O	1:A:470:GLN:HG2	2.14	0.47
1:A:391:ASP:OD2	1:A:412:ARG:NH2	2.38	0.47
1:A:286:ARG:NH2	1:A:478:ASN:HD21	2.13	0.47
1:B:512:ILE:HG21	1:B:522:LEU:HD22	1.95	0.47
1:B:189:TRP:O	1:B:189:TRP:HE3	1.96	0.46
2:B:1531:NDP:C8A	2:B:1531:NDP:O2A	2.64	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:479:VAL:N	4:B:1532:P6G:H52	2.25	0.45
1:B:189:TRP:CE3	1:B:189:TRP:CA	3.00	0.45
1:B:438:ALA:HB2	1:B:459:HIS:CD2	2.52	0.45
1:A:361:ALA:HB2	1:A:386:VAL:HG23	1.99	0.45
1:B:517:ALA:O	1:B:521:THR:HG23	2.17	0.45
1:A:0:HIS:C	1:A:0:HIS:ND1	2.71	0.44
1:A:448:LEU:HD13	1:A:464:ALA:HB1	2.00	0.44
1:A:285:VAL:HG21	1:A:319:LYS:HB3	2.00	0.43
1:A:438:ALA:HB2	1:A:459:HIS:CD2	2.53	0.43
1:A:182:LYS:HE3	1:A:216:SER:OG	2.18	0.43
1:B:304:VAL:O	1:B:410:PRO:HA	2.17	0.43
1:A:158:PHE:CD1	1:A:339:VAL:HG21	2.54	0.42
2:A:1523:NDP:H2N	5:A:2238:HOH:O	2.20	0.42
1:B:100:VAL:O	1:B:104:GLY:HA3	2.20	0.42
1:A:220:LEU:HD11	1:A:241:LEU:HG	2.02	0.41
1:A:439:SER:OG	1:A:476:HIS:CD2	2.72	0.41
1:A:285:VAL:HG13	1:A:320:LEU:HD23	2.03	0.41
1:B:255:ASN:HD21	1:B:495:GLU:N	2.05	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	517/532 (97%)	506 (98%)	11 (2%)	0	100 100
1	B	523/532 (98%)	514 (98%)	9 (2%)	0	100 100
All	All	1040/1064 (98%)	1020 (98%)	20 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/398 (98%)	385 (98%)	6 (2%)	70 48
1	B	394/398 (99%)	392 (100%)	2 (0%)	91 84
All	All	785/796 (99%)	777 (99%)	8 (1%)	80 65

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	167	GLU
1	A	216	SER
1	A	224	ARG
1	A	416	ASP
1	A	417	THR
1	A	482	MET
1	B	189	TRP
1	B	482	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	ASN
1	A	255	ASN
1	A	294	GLN
1	A	343	GLN
1	A	346	ASN
1	A	383	HIS
1	A	392	ASN
1	A	433	GLN
1	A	463	HIS
1	A	476	HIS
1	A	478	ASN
1	A	513	GLN
1	B	123	HIS
1	B	255	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	294	GLN
1	B	383	HIS
1	B	392	ASN
1	B	463	HIS
1	B	476	HIS
1	B	513	GLN
1	B	527	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NDP	A	1523	-	43,52,52	1.48	6 (13%)	49,80,80	1.90	6 (12%)
3	ETE	A	1524	-	13,13,13	0.94	0	12,12,12	0.87	0
2	NDP	B	1531	-	43,52,52	1.81	9 (20%)	49,80,80	2.84	13 (26%)
4	P6G	B	1532	-	18,18,18	0.55	0	17,17,17	1.00	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1523	-	-	0/30/77/77	0/5/5/5
3	ETE	A	1524	-	-	0/11/11/11	0/0/0/0
2	NDP	B	1531	-	-	0/30/77/77	0/5/5/5
4	P6G	B	1532	-	-	0/16/16/16	0/0/0/0

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1523	NDP	C4N-C5N	-3.99	1.40	1.49
2	B	1531	NDP	C4N-C5N	-3.94	1.40	1.49
2	B	1531	NDP	P2B-O2X	2.12	1.63	1.54
2	A	1523	NDP	P2B-O1X	2.22	1.58	1.50
2	A	1523	NDP	PA-O1A	2.38	1.59	1.50
2	B	1531	NDP	O4B-C1B	2.46	1.44	1.41
2	B	1531	NDP	PA-O1A	2.60	1.60	1.50
2	A	1523	NDP	P2B-O2B	2.66	1.64	1.59
2	B	1531	NDP	P2B-O1X	3.53	1.62	1.50
2	B	1531	NDP	PN-O5D	3.73	1.75	1.59
2	A	1523	NDP	C6N-C5N	3.76	1.40	1.33
2	B	1531	NDP	C6N-C5N	3.76	1.40	1.33
2	A	1523	NDP	PN-O1N	3.80	1.65	1.50
2	B	1531	NDP	P2B-O2B	3.84	1.66	1.59
2	B	1531	NDP	PN-O1N	4.33	1.67	1.50

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1531	NDP	N3A-C2A-N1A	-9.17	120.87	128.86
2	A	1523	NDP	N3A-C2A-N1A	-8.71	121.27	128.86
2	B	1531	NDP	O5D-PN-O1N	-8.31	75.72	109.25
2	B	1531	NDP	O2N-PN-O1N	-6.66	77.80	112.28
2	B	1531	NDP	O5B-PA-O1A	-6.48	83.11	109.25
2	A	1523	NDP	O5B-PA-O1A	-6.10	84.64	109.25
2	B	1531	NDP	O2A-PA-O5B	-4.97	84.69	108.14
2	B	1531	NDP	C5D-C4D-C3D	-3.68	101.27	115.29
2	B	1531	NDP	C4D-O4D-C1D	-3.49	101.68	109.47
2	A	1523	NDP	O2A-PA-O5B	-2.85	94.68	108.14
2	A	1523	NDP	C4D-O4D-C1D	-2.78	103.28	109.47
2	B	1531	NDP	O4D-C1D-C2D	-2.44	101.24	106.64

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1532	P6G	C8-O7-C6	-2.33	103.20	113.30
2	B	1531	NDP	C5B-C4B-C3B	-2.27	106.62	115.29
2	A	1523	NDP	O2N-PN-O5D	2.15	118.31	108.14
2	B	1531	NDP	O2A-PA-O1A	2.23	123.84	112.28
2	A	1523	NDP	O5D-PN-O1N	2.26	118.35	109.25
2	B	1531	NDP	O5D-C5D-C4D	2.27	117.06	109.00
2	B	1531	NDP	O4D-C1D-N1N	2.45	113.01	108.07
2	B	1531	NDP	O2N-PN-O5D	6.47	138.68	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1523	NDP	3	0
3	A	1524	ETE	3	0
2	B	1531	NDP	4	0
4	B	1532	P6G	6	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	521/532 (97%)	0.29	14 (2%) 55 54	12, 17, 27, 48	0
1	B	527/532 (99%)	0.14	11 (2%) 64 64	11, 15, 23, 31	0
All	All	1048/1064 (98%)	0.22	25 (2%) 59 59	11, 16, 26, 48	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	417	THR	5.3
1	A	415	THR	5.1
1	A	521	THR	5.0
1	B	189	TRP	5.0
1	A	0	HIS	4.9
1	B	0	HIS	4.8
1	A	416	ASP	4.6
1	A	520	GLY	4.6
1	A	517	ALA	4.1
1	B	471	HIS	3.5
1	A	394	THR	3.4
1	B	450	ARG	3.0
1	B	375	ASN	2.9
1	A	522	LEU	2.9
1	A	278	ASP	2.9
1	A	44	ARG	2.9
1	B	273	ASP	2.5
1	B	258	ALA	2.4
1	A	414	THR	2.4
1	A	135	LYS	2.4
1	A	357	GLU	2.3
1	B	135	LYS	2.2
1	B	448	LEU	2.1
1	B	331	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NDP	B	1531	48/48	0.71	0.27	6.08	28,50,67,68	0
2	NDP	A	1523	48/48	0.90	0.19	3.90	18,32,56,57	0
4	P6G	B	1532	19/19	0.83	0.15	2.05	26,29,35,36	0
3	ETE	A	1524	14/14	0.80	0.16	1.84	22,26,29,30	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.