



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 15, 2017 – 12:51 PM EDT

PDB ID : 5VTY
Title : Crystal structure of the A/Hong Kong/1/1968 (H3N2) influenza virus hemagglutinin G225Q/L226A mutant in complex with 3'-SLN
Authors : Wu, N.C.; Wilson, I.A.
Deposited on : unknown
Resolution : 2.36 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	<div> <div>0.2%</div> <div>92%</div> <div>5%</div> <div>..</div> </div>
1	C	323	<div> <div>0.2%</div> <div>92%</div> <div>5%</div> <div>..</div> </div>
1	E	323	<div> <div>4%</div> <div>93%</div> <div>5%</div> <div>..</div> </div>
2	B	174	<div> <div>2%</div> <div>93%</div> <div>5%</div> <div>.</div> </div>
2	D	174	<div> <div>4%</div> <div>94%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	174	 11% 94%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	407	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12540 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2438	1526	429	470	13			
1	C	317	Total	C	N	O	S	0	0	0
			2445	1531	430	471	13			
1	E	321	Total	C	N	O	S	0	0	0
			2481	1552	439	477	13			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ALA	-	expression tag	UNP Q91MA7
A	8	ASP	-	expression tag	UNP Q91MA7
A	9	PRO	-	expression tag	UNP Q91MA7
A	10	GLY	-	expression tag	UNP Q91MA7
A	225	GLN	GLY	engineered mutation	UNP Q91MA7
A	226	ALA	LEU	engineered mutation	UNP Q91MA7
C	7	ALA	-	expression tag	UNP Q91MA7
C	8	ASP	-	expression tag	UNP Q91MA7
C	9	PRO	-	expression tag	UNP Q91MA7
C	10	GLY	-	expression tag	UNP Q91MA7
C	225	GLN	GLY	engineered mutation	UNP Q91MA7
C	226	ALA	LEU	engineered mutation	UNP Q91MA7
E	7	ALA	-	expression tag	UNP Q91MA7
E	8	ASP	-	expression tag	UNP Q91MA7
E	9	PRO	-	expression tag	UNP Q91MA7
E	10	GLY	-	expression tag	UNP Q91MA7
E	225	GLN	GLY	engineered mutation	UNP Q91MA7
E	226	ALA	LEU	engineered mutation	UNP Q91MA7


- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	171	Total	C	N	O	S	0	0	0
			1382	858	241	277	6			
2	D	171	Total	C	N	O	S	0	0	0
			1382	858	241	277	6			
2	F	171	Total	C	N	O	S	0	0	0
			1382	858	241	277	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	123	GLY	ARG	conflict	UNP Q91MA7
D	123	GLY	ARG	conflict	UNP Q91MA7
F	123	GLY	ARG	conflict	UNP Q91MA7

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O		0	0
			14	8	1	5			
3	A	1	Total	C	N	O		0	0
			14	8	1	5			
3	A	1	Total	C	N	O		0	0
			14	8	1	5			
3	A	1	Total	C	N	O		0	0
			14	8	1	5			
3	A	1	Total	C	N	O		0	0
			14	8	1	5			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			15	8	1	6		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			14	8	1	5		
3	E	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		
4	C	1	Total	C	O	0	0
			11	6	5		
4	E	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	C	1	Total	C	O	0	0
			11	6	5		
5	C	1	Total	C	O	0	0
			11	6	5		
5	E	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is O-SIALIC ACID (three-letter code: SIA) (formula: C₁₁H₁₉NO₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			20	11	1	8		
6	C	1	Total	C	N	O	0	0
			20	11	1	8		
6	E	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 7 is BETA-D-GALACTOSE (three-letter code: GAL) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			11	6	5		
7	E	1	Total	C	O	0	0
			11	6	5		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	116	Total	O	0	0
			116	116		
8	B	67	Total	O	0	0
			67	67		
8	C	176	Total	O	0	0
			176	176		
8	D	78	Total	O	0	0
			78	78		
8	E	122	Total	O	0	0
			122	122		
8	F	60	Total	O	0	0
			60	60		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Hemagglutinin HA1 chain



- Molecule 1: Hemagglutinin HA1 chain



- Molecule 1: Hemagglutinin HA1 chain




- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



- Molecule 2: Hemagglutinin HA2 chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	210.08 Å 130.83 Å 72.37 Å 90.00° 98.95° 90.00°	Depositor
Resolution (Å)	110.67 – 2.36 49.99 – 2.36	Depositor EDS
% Data completeness (in resolution range)	98.0 (110.67-2.36) 98.0 (49.99-2.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.37 Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.180 , 0.210 0.184 , 0.212	Depositor DCC
R_{free} test set	3998 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12540	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, SIA, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2494	0.71	2/3399 (0.1%)
1	C	0.50	0/2502	0.75	4/3410 (0.1%)
1	E	0.47	0/2538	0.71	0/3457
2	B	0.45	0/1406	0.67	0/1890
2	D	0.44	0/1406	0.66	0/1890
2	F	0.46	0/1406	0.66	0/1890
All	All	0.47	0/11752	0.70	6/15936 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	167	THR	CB-CA-C	-6.45	94.18	111.60
1	A	201	ARG	NE-CZ-NH1	6.39	123.50	120.30
1	A	201	ARG	NE-CZ-NH2	-5.93	117.33	120.30
1	C	208	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	C	220	ARG	NE-CZ-NH2	-5.12	117.74	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2382	11	0
1	C	2445	0	2391	12	1
1	E	2481	0	2430	11	0
2	B	1382	0	1299	2	0
2	D	1382	0	1299	1	0
2	F	1382	0	1299	2	0
3	A	70	0	62	1	0
3	C	71	0	63	1	0
3	E	99	0	89	0	0
4	A	11	0	9	0	0
4	C	11	0	8	0	0
4	E	11	0	9	0	0
5	A	11	0	10	0	0
5	C	22	0	20	0	0
5	E	11	0	10	0	0
6	A	20	0	17	0	0
6	C	20	0	17	0	0
6	E	20	0	17	0	0
7	A	12	0	11	0	0
7	C	11	0	9	0	0
7	E	11	0	9	0	0
8	A	116	0	0	0	0
8	B	67	0	0	2	0
8	C	176	0	0	0	0
8	D	78	0	0	0	0
8	E	122	0	0	1	0
8	F	60	0	0	0	0
All	All	12540	0	11460	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:203:THR:HG23	1:E:212:THR:HG22	1.75	0.69
1:C:212:THR:HG21	1:E:216:ASN:CB	2.23	0.67
1:C:203:THR:OG1	1:C:212:THR:HG22	1.94	0.66
1:C:167:THR:HG23	1:C:244:VAL:HG22	1.80	0.63
1:C:53:ASN:HD21	1:C:276:THR:HG22	1.63	0.63

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:119:GLU:OE1	1:C:119:GLU:OE1[2_558]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/323 (97%)	306 (98%)	7 (2%)	1 (0%)	44	53
1	C	315/323 (98%)	308 (98%)	6 (2%)	1 (0%)	44	53
1	E	319/323 (99%)	311 (98%)	7 (2%)	1 (0%)	44	53
2	B	169/174 (97%)	161 (95%)	8 (5%)	0	100	100
2	D	169/174 (97%)	161 (95%)	8 (5%)	0	100	100
2	F	169/174 (97%)	161 (95%)	8 (5%)	0	100	100
All	All	1455/1491 (98%)	1408 (97%)	44 (3%)	3 (0%)	51	61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	ILE
1	C	62	ILE
1	E	62	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/283 (98%)	268 (97%)	9 (3%)	44	55
1	C	278/283 (98%)	269 (97%)	9 (3%)	44	55
1	E	282/283 (100%)	275 (98%)	7 (2%)	53	66
2	B	145/148 (98%)	138 (95%)	7 (5%)	30	37
2	D	145/148 (98%)	139 (96%)	6 (4%)	35	44
2	F	145/148 (98%)	140 (97%)	5 (3%)	42	53
All	All	1272/1293 (98%)	1229 (97%)	43 (3%)	42	53

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	186	SER
1	C	227	SER
2	F	38	LEU
1	C	189	GLN
1	C	208	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
2	B	53	ASN
1	E	22	ASN
1	E	33	GLN
1	E	225	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	A	401	1,3	14,14,15	0.42	0	15,19,21	1.03	1 (6%)
3	NAG	A	402	3,4	14,14,15	0.40	0	15,19,21	0.48	0
4	BMA	A	403	3,5	11,11,12	0.37	0	13,15,17	0.87	0
5	MAN	A	404	4	11,11,12	0.37	0	13,15,17	0.88	1 (7%)
3	NAG	A	405	1,3	14,14,15	0.45	0	15,19,21	0.92	0
3	NAG	A	406	3	14,14,15	0.32	0	15,19,21	1.32	1 (6%)
3	NAG	A	407	1	14,14,15	0.69	1 (7%)	15,19,21	2.92	7 (46%)
6	SIA	A	408	7	17,20,21	0.45	0	19,28,31	0.52	0
7	GAL	A	409	6	12,12,12	0.55	0	17,17,17	0.61	0
3	NAG	C	401	1,3	14,14,15	0.25	0	15,19,21	1.29	2 (13%)
3	NAG	C	402	3,4	14,14,15	0.30	0	15,19,21	1.05	2 (13%)
4	BMA	C	403	3,5	11,11,12	0.56	0	13,15,17	0.72	0
5	MAN	C	404	4	11,11,12	0.39	0	13,15,17	1.14	2 (15%)
5	MAN	C	405	4	11,11,12	0.44	0	13,15,17	0.95	1 (7%)
3	NAG	C	406	1,3	14,14,15	0.54	0	15,19,21	1.26	3 (20%)
3	NAG	C	407	3	14,14,15	0.56	0	15,19,21	1.96	1 (6%)
6	SIA	C	408	7	17,20,21	0.39	0	19,28,31	0.62	0
7	GAL	C	409	3,6	11,11,12	0.42	0	13,15,17	0.70	0
3	NAG	C	410	7	15,15,15	0.55	0	21,21,21	1.37	4 (19%)
3	NAG	E	501	1	14,14,15	0.35	0	15,19,21	1.09	1 (6%)
3	NAG	E	502	1,3	14,14,15	0.42	0	15,19,21	0.94	1 (6%)
3	NAG	E	503	3,4	14,14,15	0.33	0	15,19,21	0.92	1 (6%)
4	BMA	E	504	3,5	11,11,12	0.43	0	13,15,17	1.26	2 (15%)
5	MAN	E	505	4	11,11,12	0.45	0	13,15,17	1.26	2 (15%)
3	NAG	E	506	1,3	14,14,15	0.41	0	15,19,21	0.62	0
3	NAG	E	507	3	14,14,15	0.25	0	15,19,21	0.49	0
3	NAG	E	508	1	14,14,15	0.26	0	15,19,21	0.54	0
6	SIA	E	509	7	17,20,21	0.40	0	19,28,31	0.60	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GAL	E	510	3,6	11,11,12	0.68	0	13,15,17	1.23	2 (15%)
3	NAG	E	511	7	15,15,15	0.56	0	21,21,21	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	402	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	403	3,5	-	0/2/19/22	0/1/1/1
5	MAN	A	404	4	-	0/2/19/22	0/1/1/1
3	NAG	A	405	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	406	3	-	0/6/23/26	0/1/1/1
3	NAG	A	407	1	-	0/6/23/26	0/1/1/1
6	SIA	A	408	7	-	0/14/34/38	0/1/1/1
7	GAL	A	409	6	-	0/2/22/22	0/1/1/1
3	NAG	C	401	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	402	3,4	-	0/6/23/26	0/1/1/1
4	BMA	C	403	3,5	-	0/2/19/22	0/1/1/1
5	MAN	C	404	4	-	0/2/19/22	0/1/1/1
5	MAN	C	405	4	-	0/2/19/22	0/1/1/1
3	NAG	C	406	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	407	3	-	0/6/23/26	0/1/1/1
6	SIA	C	408	7	-	0/14/34/38	0/1/1/1
7	GAL	C	409	3,6	-	0/2/19/22	0/1/1/1
3	NAG	C	410	7	-	0/6/26/26	0/1/1/1
3	NAG	E	501	1	-	0/6/23/26	0/1/1/1
3	NAG	E	502	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	503	3,4	-	0/6/23/26	0/1/1/1
4	BMA	E	504	3,5	-	0/2/19/22	0/1/1/1
5	MAN	E	505	4	-	0/2/19/22	0/1/1/1
3	NAG	E	506	1,3	-	0/6/23/26	0/1/1/1
3	NAG	E	507	3	-	0/6/23/26	0/1/1/1
3	NAG	E	508	1	-	0/6/23/26	0/1/1/1
6	SIA	E	509	7	-	0/14/34/38	0/1/1/1
7	GAL	E	510	3,6	-	0/2/19/22	0/1/1/1
3	NAG	E	511	7	-	0/6/26/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	407	NAG	O5-C1	-2.08	1.40	1.43

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	407	NAG	C1-C2-N2	-6.01	100.22	110.49
3	A	407	NAG	C1-O5-C5	-4.13	106.47	112.17
3	C	401	NAG	O5-C1-C2	-3.05	107.23	111.47
7	E	510	GAL	O5-C1-C2	-2.88	106.28	110.79
3	E	502	NAG	O5-C1-C2	-2.51	107.99	111.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	407	NAG	1	0
3	C	402	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/323 (97%)	-0.13	3 (0%) 84 91	34, 47, 67, 99	0
1	C	317/323 (98%)	-0.02	3 (0%) 84 91	27, 40, 65, 115	0
1	E	321/323 (99%)	0.05	13 (4%) 39 51	31, 48, 79, 117	0
2	B	171/174 (98%)	0.27	3 (1%) 69 78	29, 56, 89, 100	0
2	D	171/174 (98%)	0.40	7 (4%) 38 50	28, 62, 97, 111	0
2	F	171/174 (98%)	0.62	20 (11%) 5 9	28, 62, 93, 107	0
All	All	1467/1491 (98%)	0.13	49 (3%) 47 59	27, 48, 87, 117	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	9	PRO	5.8
2	F	59	THR	4.3
2	F	31	GLY	4.1
2	D	171	PHE	4.1
2	F	161	VAL	4.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	A	407	14/15	0.78	0.28	10.57	71,80,88,89	0
3	NAG	E	506	14/15	0.93	0.14	1.62	45,50,57,67	0
3	NAG	C	406	14/15	0.94	0.12	1.24	44,49,60,70	0
6	SIA	A	408	20/21	0.94	0.14	0.99	53,61,66,66	0
3	NAG	A	405	14/15	0.95	0.14	0.72	48,56,60,68	0
6	SIA	E	509	20/21	0.93	0.16	0.31	66,72,74,76	0
6	SIA	C	408	20/21	0.94	0.11	-0.67	42,48,52,53	0
3	NAG	A	401	14/15	0.97	0.11	-1.39	41,46,52,59	0
3	NAG	C	401	14/15	0.95	0.10	-1.41	41,45,48,49	0
3	NAG	A	402	14/15	0.95	0.16	-1.62	53,57,68,74	0
3	NAG	C	402	14/15	0.94	0.10	-2.08	42,53,58,68	0
5	MAN	C	405	11/12	0.66	0.26	-	83,90,98,100	0
7	GAL	C	409	11/12	0.96	0.18	-	61,71,73,76	0
4	BMA	A	403	11/12	0.92	0.12	-	78,84,88,97	0
3	NAG	E	507	14/15	0.81	0.31	-	71,84,92,93	0
3	NAG	E	508	14/15	0.93	0.20	-	55,63,69,70	0
7	GAL	A	409	12/12	0.86	0.21	-	72,80,89,91	0
4	BMA	E	504	11/12	0.91	0.18	-	93,96,100,106	0
3	NAG	E	501	14/15	0.88	0.24	-	65,74,76,76	0
3	NAG	A	406	14/15	0.83	0.28	-	74,82,89,91	0
3	NAG	C	410	15/15	0.81	0.46	-	84,96,105,106	0
5	MAN	A	404	11/12	0.86	0.33	-	104,109,111,111	0
5	MAN	C	404	11/12	0.86	0.30	-	91,95,97,99	0
4	BMA	C	403	11/12	0.86	0.14	-	78,87,90,95	0
3	NAG	E	502	14/15	0.95	0.21	-	61,68,73,76	0
3	NAG	E	503	14/15	0.94	0.21	-	71,74,81,87	0
7	GAL	E	510	11/12	0.91	0.27	-	89,96,99,102	0
3	NAG	E	511	15/15	0.88	0.39	-	107,112,114,115	0
5	MAN	E	505	11/12	0.75	0.29	-	108,110,114,114	0
3	NAG	C	407	14/15	0.81	0.27	-	81,88,95,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.