



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 11:59 am GMT

PDB ID : 2VVM  
Title : THE STRUCTURE OF MAO-N-D5, A VARIANT OF MONOAMINE OXIDASE FROM ASPERGILLUS NIGER.  
Authors : Atkin, K.E.; Hart, S.; Turkenburg, J.P.; Brzozowski, A.M.; Grogan, G.J.  
Deposited on : 2008-06-10  
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1923 (1.86-1.86)
Clashscore	112137	2083 (1.86-1.86)
Ramachandran outliers	110173	2060 (1.86-1.86)
Sidechain outliers	110143	2060 (1.86-1.86)
RSRZ outliers	101464	1932 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div>.</div> </div> </div>
1	B	495	<div> <div>%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PRO	A	601	-	-	-	X
3	FAD	A	600	X	-	-	-
3	FAD	B	600	X	-	-	-
4	EDO	B	604	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MONOAMINE OXIDASE N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	481	Total	C	N	O	S	17	15	0
			3820	2413	677	707	23			
1	B	478	Total	C	N	O	S	50	25	0
			3795	2398	671	703	23			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	246	MET	ILE	ENGINEERED MUTATION	UNP P46882
A	336	SER	ASN	ENGINEERED MUTATION	UNP P46882
A	384	ASN	THR	ENGINEERED MUTATION	UNP P46882
A	385	SER	ASP	ENGINEERED MUTATION	UNP P46882
A	300	VAL	ALA	CONFLICT	UNP P46882
A	304	VAL	LEU	CONFLICT	UNP P46882
A	450	GLY	ARG	CONFLICT	UNP P46882
B	246	MET	ILE	ENGINEERED MUTATION	UNP P46882
B	336	SER	ASN	ENGINEERED MUTATION	UNP P46882
B	384	ASN	THR	ENGINEERED MUTATION	UNP P46882
B	385	SER	ASP	ENGINEERED MUTATION	UNP P46882
B	300	VAL	ALA	CONFLICT	UNP P46882
B	304	VAL	LEU	CONFLICT	UNP P46882
B	450	GLY	ARG	CONFLICT	UNP P46882

- Molecule 2 is PROLINE (three-letter code: PRO) (formula: C<sub>5</sub>H<sub>9</sub>NO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	5	1	2		
2	B	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	357	Total	O	0	0
			357	357		
5	B	348	Total	O	0	0
			348	348		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: MONOAMINE OXIDASE N



#### • Molecule 1: MONOAMINE OXIDASE N



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.42Å 107.42Å 235.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	97.59 – 1.85 29.79 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (97.59-1.85) 99.7 (29.79-1.85)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.45 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0065	Depositor
R, $R_{free}$	0.181 , 0.205 0.181 , 0.204	Depositor DCC
$R_{free}$ test set	5889 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.8	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8466	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/3936	0.57	2/5335 (0.0%)
1	B	0.56	3/3907 (0.1%)	0.68	5/5296 (0.1%)
All	All	0.52	3/7843 (0.0%)	0.63	7/10631 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	150[A]	ARG	CZ-NH2	-5.66	1.25	1.33
1	B	150[B]	ARG	CZ-NH2	-5.66	1.25	1.33
1	B	317[A]	GLN	CG-CD	5.05	1.62	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	196[A]	ARG	NE-CZ-NH1	16.57	128.59	120.30
1	B	196[A]	ARG	NE-CZ-NH2	-16.51	112.04	120.30
1	B	482[A]	LEU	CB-CG-CD2	8.40	125.28	111.00
1	A	435	PRO	CA-N-CD	-8.19	100.03	111.50
1	A	56	ARG	NE-CZ-NH2	-7.30	116.65	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3820	0	3620	23	0
1	B	3795	0	3554	22	0
2	A	8	0	7	1	0
2	B	8	0	7	1	0
3	A	53	0	28	3	0
3	B	53	0	28	4	0
4	A	12	0	18	2	0
4	B	12	0	18	5	0
5	A	357	0	0	3	0
5	B	348	0	0	2	0
All	All	8466	0	7280	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:ARG:NH2	1:A:105:GLU:OE1	2.01	0.93
1:B:146[A]:ASP:OD2	1:B:150[A]:ARG:CZ	2.19	0.90
2:B:601:PRO:N	4:B:602:EDO:HO2	1.86	0.73
1:A:8:GLN:NE2	1:A:414:ARG:HH21	1.87	0.72
1:B:8:GLN:NE2	1:B:414:ARG:HH21	1.90	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	482/495 (97%)	471 (98%)	11 (2%)	0	100	100
1	B	478/495 (97%)	467 (98%)	11 (2%)	0	100	100
All	All	960/990 (97%)	938 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	406/414 (98%)	403 (99%)	3 (1%)	87	83
1	B	403/414 (97%)	398 (99%)	5 (1%)	75	67
All	All	809/828 (98%)	801 (99%)	8 (1%)	80	74

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	22	LEU
1	B	461	LEU
1	B	286[A]	ARG
1	A	485	LEU
1	B	127	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	404	GLN
1	B	8	GLN
1	B	284	ASN
1	A	388	HIS
1	B	229	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FAD	A	600	-	51,58,58	2.22	21 (41%)	54,89,89	2.67	19 (35%)
2	PRO	A	601	-	5,8,8	0.55	0	6,10,10	1.10	0
4	EDO	A	602	-	3,3,3	0.38	0	2,2,2	0.58	0
4	EDO	A	603	-	3,3,3	0.40	0	2,2,2	0.58	0
4	EDO	A	604	-	3,3,3	0.38	0	2,2,2	0.09	0
3	FAD	B	600	-	51,58,58	2.07	20 (39%)	54,89,89	2.95	20 (37%)
2	PRO	B	601	-	5,8,8	0.53	0	6,10,10	1.06	1 (16%)
4	EDO	B	602	-	3,3,3	0.36	0	2,2,2	0.58	0
4	EDO	B	603	-	3,3,3	0.48	0	2,2,2	0.27	0
4	EDO	B	604	-	3,3,3	0.46	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	600	-	3/3/9/9	0/28/50/50	0/6/6/6
2	PRO	A	601	-	-	0/0/11/11	0/1/1/1
4	EDO	A	602	-	-	0/1/1/1	0/0/0/0
4	EDO	A	603	-	-	0/1/1/1	0/0/0/0
4	EDO	A	604	-	-	0/1/1/1	0/0/0/0
3	FAD	B	600	-	3/3/9/9	0/28/50/50	0/6/6/6
2	PRO	B	601	-	-	0/0/11/11	0/1/1/1
4	EDO	B	602	-	-	0/1/1/1	0/0/0/0
4	EDO	B	603	-	-	0/1/1/1	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	604	-	-	0/1/1/1	0/0/0/0

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600	FAD	P-O1P	-5.40	1.30	1.50
3	A	600	FAD	PA-O1A	-3.95	1.36	1.50
3	B	600	FAD	PA-O1A	-3.63	1.37	1.50
3	A	600	FAD	O4'-C4'	-3.62	1.35	1.43
3	A	600	FAD	PA-O2A	-3.52	1.37	1.55

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	FAD	C2B-C3B-C4B	-8.36	86.33	102.62
3	B	600	FAD	O4B-C4B-C3B	-7.36	90.54	105.17
3	B	600	FAD	N3A-C2A-N1A	-6.81	122.92	128.86
3	A	600	FAD	N3A-C2A-N1A	-6.81	122.93	128.86
3	A	600	FAD	C2B-C3B-C4B	-6.01	90.91	102.62

5 of 6 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	B	600	FAD	C4B
3	B	600	FAD	C2'
3	B	600	FAD	C3B
3	A	600	FAD	C4B
3	A	600	FAD	C2'

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	FAD	3	0
2	A	601	PRO	1	0
4	A	602	EDO	1	0
4	A	604	EDO	1	0
3	B	600	FAD	4	0
2	B	601	PRO	1	0
4	B	602	EDO	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	604	EDO	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	481/495 (97%)	-0.21	6 (1%) 79 79	16, 24, 36, 52	8 (1%)
1	B	478/495 (96%)	-0.22	6 (1%) 77 78	16, 24, 40, 49	19 (3%)
All	All	959/990 (96%)	-0.21	12 (1%) 77 78	16, 24, 38, 52	27 (2%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	487	THR	4.6
1	A	38	GLY	3.9
1	B	287[A]	ASP	3.6
1	A	1	MET	3.3
1	B	1	MET	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	EDO	B	604	4/4	0.76	0.17	11.29	39,40,40,40	0
2	PRO	A	601	8/8	0.79	0.24	6.18	37,37,37,37	0
2	PRO	B	601	8/8	0.87	0.19	1.43	33,34,34,34	0
4	EDO	A	604	4/4	0.95	0.13	0.92	28,30,31,32	0
4	EDO	B	603	4/4	0.90	0.12	0.88	39,39,40,41	0
3	FAD	B	600	53/53	0.96	0.13	0.18	18,21,28,29	0
3	FAD	A	600	53/53	0.97	0.11	-0.25	18,21,24,28	0
4	EDO	A	603	4/4	0.97	0.08	-0.45	23,23,24,25	0
4	EDO	B	602	4/4	0.89	0.25	-	31,31,32,32	0
4	EDO	A	602	4/4	0.93	0.24	-	30,30,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.