



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:47 pm GMT

PDB ID : 2VW9  
Title : SINGLE STRANDED DNA BINDING PROTEIN COMPLEX FROM HELI-COBACTER PYLORI  
Authors : Chan, K.-W.; Wang, C.-H.; Lee, Y.-J.; Sun, Y.-J.  
Deposited on : 2008-06-18  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	134	<div> <div>4%</div> <div>56%</div> <div>21%</div> <div>• •</div> <div>19%</div> </div>
1	B	134	<div> <div>10%</div> <div>51%</div> <div>20%</div> <div>8%</div> <div>21%</div> </div>
2	C	35	<div> <div>29%</div> <div>9%</div> <div>49%</div> <div>11%</div> <div>31%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2272 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SINGLE-STRANDED DNA BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	1
			836	522	150	159	5			
1	B	106	Total	C	N	O	S	0	0	1
			811	508	146	152	5			

- Molecule 2 is a DNA chain called POLY-DT.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	24	Total	C	N	O	P	0	0	0
			465	240	48	158	19			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	85	Total	O	0	0
			85	85		
3	B	47	Total	O	0	0
			47	47		
3	C	28	Total	O	0	0
			28	28		



- Molecule 1: SINGLE-STRANDED DNA BINDING PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.56Å 75.56Å 117.69Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.13 – 2.30 25.13 – 2.30	Depositor EDS
% Data completeness (in resolution range)	89.5 (25.13-2.30) 93.9 (25.13-2.30)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.74 (at 2.31Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.238 , 0.260 0.241 , 0.253	Depositor DCC
$R_{free}$ test set	812 reflections (5.11%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.8	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	2272	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/847	0.78	0/1138
1	B	0.56	0/822	0.79	0/1105
2	C	0.91	0/508	1.41	7/776 (0.9%)
All	All	0.66	0/2177	0.99	7/3019 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	20	DT	O4'-C1'-N1	8.16	113.71	108.00
2	C	25	DT	O4'-C1'-C2'	6.40	111.02	105.90
2	C	25	DT	O4'-C4'-C3'	5.59	109.35	106.00
2	C	32	DT	O4'-C1'-C2'	5.56	110.35	105.90
2	C	18	DT	O4'-C1'-N1	5.50	111.85	108.00
2	C	25	DT	N1-C1'-C2'	5.20	122.47	112.60
2	C	21	DT	O4'-C1'-C2'	5.16	110.03	105.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1035	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	836	0	832	20	0
1	B	811	0	811	34	0
2	C	465	0	298	44	0
3	A	85	0	0	2	0
3	B	47	0	0	1	0
3	C	28	0	0	0	0
All	All	2272	0	1941	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1091:LYS:H	1:A:1091:LYS:HE2	1.23	0.98
1:B:2087:GLN:H	1:B:2087:GLN:HE21	1.21	0.86
2:C:19:DT:O2	2:C:20:DT:H73	1.78	0.83
1:B:2087:GLN:N	1:B:2087:GLN:HE21	1.82	0.77
1:A:1074:LEU:HB3	1:A:1103:GLN:HB3	1.71	0.72
2:C:30:DT:H2''	2:C:31:DT:OP1	1.89	0.70
1:A:1075:ILE:HD11	1:A:1099:ALA:HB2	1.72	0.69
1:B:2084:TRP:CZ2	2:C:23:DT:H72	2.29	0.67
2:C:19:DT:O2	2:C:20:DT:C7	2.42	0.67
1:B:2075:ILE:HG22	1:B:2102:LEU:HD22	1.75	0.65
2:C:14:DT:H6	2:C:14:DT:H3'	1.60	0.65
1:A:1041:ASP:OD2	1:A:1041:ASP:N	2.27	0.65
2:C:7:DT:H4'	2:C:8:DT:OP1	1.96	0.64
1:B:2056:PHE:CE2	2:C:20:DT:H71	2.32	0.64
1:B:2084:TRP:CZ2	2:C:23:DT:C7	2.80	0.64
1:B:2035:ARG:HD3	1:B:2047:GLU:OE2	1.98	0.63
1:A:1075:ILE:HD11	1:A:1099:ALA:CB	2.29	0.62
1:B:2084:TRP:HZ2	2:C:23:DT:C7	2.12	0.62
2:C:14:DT:H4'	2:C:15:DT:OP1	2.00	0.62
1:B:2018:LYS:NZ	2:C:20:DT:O2	2.27	0.61
2:C:25:DT:C2'	2:C:26:DT:H71	2.29	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1:DT:H4'	2:C:2:DT:O5'	2.02	0.60
2:C:18:DT:H2''	2:C:19:DT:H5''	1.84	0.60
1:B:2056:PHE:CZ	2:C:20:DT:H71	2.37	0.59
1:A:1052:ASP:OD1	1:A:1094:ARG:NH1	2.36	0.59
1:B:2038:LYS:HA	1:B:2044:LEU:HA	1.85	0.59
1:A:1091:LYS:N	1:A:1091:LYS:HE2	2.07	0.58
1:A:1055:LEU:HD12	1:A:1099:ALA:HB3	1.83	0.58
1:B:2056:PHE:CE2	2:C:20:DT:C7	2.88	0.57
1:B:2084:TRP:CH2	2:C:23:DT:H72	2.40	0.56
1:A:1091:LYS:H	1:A:1091:LYS:CE	2.10	0.55
1:B:2017:LEU:HD11	1:B:2025:ALA:HB1	1.90	0.54
2:C:14:DT:C6	2:C:14:DT:H3'	2.43	0.54
2:C:25:DT:H2'	2:C:26:DT:H71	1.90	0.54
3:B:3017:HOH:O	2:C:33:DT:H73	2.07	0.53
2:C:18:DT:H1'	2:C:19:DT:H5''	1.90	0.53
2:C:23:DT:H2'	2:C:23:DT:O2	2.08	0.53
1:B:2040:GLN:O	1:B:2042:GLY:N	2.42	0.53
1:B:2104:PHE:HE1	2:C:14:DT:H2'	1.74	0.52
1:A:1040:GLN:HG2	1:A:1040:GLN:O	2.10	0.52
1:B:2043:THR:HG23	1:B:2044:LEU:N	2.25	0.52
2:C:18:DT:C2'	2:C:19:DT:H5''	2.39	0.51
1:B:2040:GLN:HG3	1:B:2043:THR:O	2.11	0.50
2:C:19:DT:C2	2:C:20:DT:C7	2.94	0.49
1:B:2056:PHE:HE2	2:C:20:DT:C7	2.24	0.49
1:A:1075:ILE:HD13	1:A:1076:GLU:N	2.27	0.49
1:B:2074:LEU:HD13	1:B:2075:ILE:N	2.28	0.48
3:A:2006:HOH:O	2:C:33:DT:H4'	2.13	0.48
1:B:2065:GLN:HG3	1:B:2066:TYR:CD1	2.48	0.48
2:C:24:DT:H2''	2:C:25:DT:OP1	2.13	0.48
1:A:1039:LYS:O	1:A:1041:ASP:N	2.47	0.48
2:C:7:DT:H2''	2:C:8:DT:H72	1.96	0.48
1:A:1010:ARG:HA	1:A:1071:SER:O	2.15	0.47
2:C:14:DT:C6	2:C:14:DT:C3'	2.97	0.47
1:B:2074:LEU:HD13	1:B:2074:LEU:C	2.35	0.47
2:C:20:DT:H2''	2:C:21:DT:OP1	2.13	0.46
1:A:1007:MET:HB3	1:A:1007:MET:HE3	1.75	0.46
1:B:2085:MET:SD	1:B:2091:LYS:HB3	2.56	0.46
2:C:19:DT:H2'	2:C:20:DT:C7	2.45	0.45
2:C:7:DT:H2''	2:C:8:DT:C7	2.46	0.45
1:B:2010:ARG:HA	1:B:2071:SER:O	2.17	0.45
2:C:18:DT:H2''	2:C:19:DT:C5'	2.45	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:24:DT:O4'	2:C:24:DT:O2	2.34	0.45
1:B:2091:LYS:HE2	1:B:2091:LYS:HB2	1.74	0.45
1:A:1053:ALA:HA	1:A:1097:ILE:O	2.17	0.44
1:B:2053:ALA:HA	1:B:2097:ILE:O	2.16	0.44
2:C:8:DT:C6	2:C:9:DT:H72	2.52	0.44
2:C:7:DT:C4'	2:C:8:DT:OP1	2.65	0.44
2:C:19:DT:C2	2:C:20:DT:H72	2.53	0.44
1:B:2074:LEU:CD1	1:B:2074:LEU:C	2.87	0.43
1:A:1040:GLN:NE2	3:A:2033:HOH:O	2.51	0.43
1:A:1011:LEU:HA	1:A:1030:GLY:O	2.18	0.43
1:B:2056:PHE:CE2	2:C:20:DT:O4	2.72	0.43
2:C:30:DT:O2	2:C:30:DT:O4'	2.35	0.43
1:B:2003:ASN:HD21	1:B:2079:LEU:H	1.67	0.43
2:C:1:DT:H4'	2:C:2:DT:O4'	2.20	0.42
2:C:4:DT:H1'	2:C:5:DT:H71	2.01	0.42
1:B:2035:ARG:CD	1:B:2047:GLU:OE2	2.66	0.42
1:B:2058:ARG:HG3	1:B:2058:ARG:O	2.19	0.42
1:B:2044:LEU:O	1:B:2044:LEU:HG	2.20	0.42
1:B:2087:GLN:H	1:B:2087:GLN:NE2	2.02	0.41
1:A:1007:MET:HE3	1:A:1033:THR:OG1	2.20	0.41
1:B:2070:GLY:HA2	2:C:26:DT:H2''	2.02	0.41
1:A:1016:GLU:HG2	1:A:1016:GLU:H	1.73	0.41
1:A:1008:VAL:O	1:A:1033:THR:HG23	2.21	0.40
2:C:20:DT:C2'	2:C:21:DT:OP1	2.70	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	107/134 (80%)	103 (96%)	2 (2%)	2 (2%)	<b>9</b> <b>8</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	104/134 (78%)	100 (96%)	3 (3%)	1 (1%)	18	20
All	All	211/268 (79%)	203 (96%)	5 (2%)	3 (1%)	13	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1040	GLN
1	B	2041	ASP
1	A	1042	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/113 (78%)	72 (82%)	16 (18%)	2	1
1	B	85/113 (75%)	66 (78%)	19 (22%)	1	1
All	All	173/226 (76%)	138 (80%)	35 (20%)	1	1

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1010	ARG
1	A	1015	VAL
1	A	1017	LEU
1	A	1018	LYS
1	A	1020	LEU
1	A	1031	LEU
1	A	1036	ARG
1	A	1040	GLN
1	A	1041	ASP
1	A	1043	THR
1	A	1044	LEU
1	A	1055	LEU
1	A	1075	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1088	THR
1	A	1090	LYS
1	A	1091	LYS
1	B	2003	ASN
1	B	2007	MET
1	B	2036	ARG
1	B	2040	GLN
1	B	2043	THR
1	B	2044	LEU
1	B	2046	GLU
1	B	2047	GLU
1	B	2055	LEU
1	B	2058	ARG
1	B	2074	LEU
1	B	2086	ASP
1	B	2087	GLN
1	B	2088	THR
1	B	2091	LYS
1	B	2102	LEU
1	B	2103	GLN
1	B	2104	PHE
1	B	2105	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1065	GLN
1	A	1103	GLN
1	B	2003	ASN
1	B	2040	GLN
1	B	2087	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	109/134 (81%)	0.38	6 (5%) 26 32	21, 36, 61, 68	0
1	B	106/134 (79%)	0.76	14 (13%) 4 5	23, 45, 67, 72	0
2	C	24/35 (68%)	1.66	10 (41%) 0 0	46, 80, 91, 96	0
All	All	239/303 (78%)	0.68	30 (12%) 4 6	21, 43, 80, 96	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1044	LEU	5.8
1	B	2044	LEU	5.4
1	B	2038	LYS	5.3
2	C	26	DT	4.9
1	A	1040	GLN	4.6
2	C	18	DT	4.5
1	B	2040	GLN	4.4
1	B	2085	MET	4.1
2	C	25	DT	3.8
2	C	24	DT	3.8
1	A	1041	ASP	3.7
1	B	2105	MET	3.7
1	B	2087	GLN	3.7
1	B	2088	THR	3.7
2	C	14	DT	3.5
2	C	30	DT	3.3
1	B	2039	LYS	3.2
2	C	20	DT	3.2
1	B	2091	LYS	3.2
1	A	1042	GLY	3.0
1	B	2037	PHE	2.8
1	B	2045	GLY	2.8
1	A	1043	THR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	C	16	DT	2.3
1	A	1053	ALA	2.3
2	C	19	DT	2.3
2	C	8	DT	2.2
1	B	2090	LYS	2.2
1	B	2043	THR	2.2
1	B	2089	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.