



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 01:52 pm GMT

PDB ID : 3W04
Title : Crystal structure of Oryza sativa DWARF14 (D14)
Authors : Kagiya, M.; Hirano, Y.; Mori, T.; Kim, S.Y.; Kyojuka, J.; Seto, Y.; Yamaguchi, S.; Hakoshima, T.
Deposited on : 2012-10-19
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1510 (1.48-1.44)
Clashscore	112137	1573 (1.48-1.44)
Ramachandran outliers	110173	1555 (1.48-1.44)
Sidechain outliers	110143	1555 (1.48-1.44)
RSRZ outliers	101464	1516 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 84%, grey 13%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 84% 12% .. </div> </div>
1	B	266	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 2%, yellow 18%, green 79%, grey 19%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 2% 79% 18% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MPD	B	402	-	-	-	X
3	EDO	A	403	-	-	-	X
3	EDO	A	404	-	-	X	-
3	EDO	B	406	-	X	X	X
4	EPE	B	401	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4714 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dwarf 88 esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	9	0
			2096	1336	372	378	10			
1	B	263	Total	C	N	O	S	0	11	0
			2107	1340	381	376	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	GLY	-	EXPRESSION TAG	UNP Q10QA5
A	54	PRO	-	EXPRESSION TAG	UNP Q10QA5
B	53	GLY	-	EXPRESSION TAG	UNP Q10QA5
B	54	PRO	-	EXPRESSION TAG	UNP Q10QA5

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	6	2		
2	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	232	Total	O	0	0
			232	232		
5	B	220	Total	O	0	0
			220	220		

- Molecule 1: Dwarf 88 esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.99Å 88.19Å 121.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.62 – 1.45 26.45 – 1.45	Depositor EDS
% Data completeness (in resolution range)	99.7 (25.62-1.45) 99.6 (26.45-1.45)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.160 , 0.194 0.161 , 0.195	Depositor DCC
R_{free} test set	4586 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	10.9	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4714	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, EPE, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	5/2156 (0.2%)	1.43	22/2933 (0.8%)
1	B	1.37	7/2160 (0.3%)	1.40	23/2939 (0.8%)
All	All	1.38	12/4316 (0.3%)	1.42	45/5872 (0.8%)

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	218	VAL	CB-CG1	-7.06	1.38	1.52
1	A	278	TYR	CE2-CZ	6.09	1.46	1.38
1	A	224	GLU	CB-CG	-5.88	1.41	1.52
1	A	140	ARG	CB-CG	-5.84	1.36	1.52
1	B	147	SER	CA-CB	5.59	1.61	1.52
1	B	246	LYS	CB-CG	-5.54	1.37	1.52
1	B	173	SER	CB-OG	5.44	1.49	1.42
1	A	124	ALA	CA-CB	5.42	1.63	1.52
1	B	270	SER	CA-CB	5.35	1.60	1.52
1	B	278	TYR	CG-CD1	5.33	1.46	1.39
1	A	164	PHE	CE1-CZ	5.13	1.47	1.37
1	B	190	GLU	CD-OE1	-5.07	1.20	1.25

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	ARG	NE-CZ-NH1	-11.56	114.52	120.30
1	A	70	ARG	NE-CZ-NH1	10.34	125.47	120.30
1	A	230	PHE	CB-CG-CD1	9.69	127.58	120.80
1	B	134	ASP	CB-CG-OD2	-9.61	109.65	118.30
1	A	159	ARG	NE-CZ-NH2	-9.49	115.55	120.30
1	B	127	ASP	CB-CG-OD2	-9.38	109.85	118.30
1	A	115	ASP	CB-CG-OD2	-8.94	110.26	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	120	ASP	CB-CG-OD2	-8.93	110.26	118.30
1	A	118	ARG	CG-CD-NE	-8.87	93.17	111.80
1	A	97	ARG	NE-CZ-NH1	8.60	124.60	120.30
1	B	230	PHE	CB-CG-CD1	-8.57	114.80	120.80
1	A	230	PHE	CB-CG-CD2	-8.48	114.87	120.80
1	A	134	ASP	CB-CG-OD2	-8.44	110.70	118.30
1	B	312	ARG	NE-CZ-NH2	-8.06	116.27	120.30
1	B	233	ARG	NE-CZ-NH2	-8.04	116.28	120.30
1	B	230	PHE	CB-CG-CD2	7.97	126.38	120.80
1	B	120	ASP	CB-CG-OD1	7.22	124.80	118.30
1	B	250	ARG	NE-CZ-NH1	-7.18	116.71	120.30
1	B	159	ARG	NE-CZ-NH1	7.15	123.88	120.30
1	B	118	ARG	NE-CZ-NH1	-7.11	116.74	120.30
1	A	162	ASP	CB-CG-OD2	7.03	124.62	118.30
1	B	248	ASP	CB-CG-OD2	-7.02	111.98	118.30
1	A	123	ASP	CB-CG-OD1	6.99	124.59	118.30
1	A	128	ASP	CB-CG-OD2	-6.88	112.11	118.30
1	A	248	ASP	CB-CG-OD1	6.88	124.49	118.30
1	A	248	ASP	CB-CG-OD2	-6.86	112.12	118.30
1	A	229	LEU	CA-CB-CG	6.45	130.13	115.30
1	B	182	TYR	CD1-CE1-CZ	-6.24	114.18	119.80
1	B	112	ASP	CB-CG-OD2	-6.21	112.72	118.30
1	B	137	ARG	NE-CZ-NH2	-6.15	117.23	120.30
1	B	117	ARG	NE-CZ-NH1	6.12	123.36	120.30
1	A	159	ARG	NE-CZ-NH1	6.07	123.33	120.30
1	B	123	ASP	CB-CG-OD2	-6.02	112.88	118.30
1	B	94	ARG	NE-CZ-NH2	-5.97	117.31	120.30
1	A	175	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	A	235	ASP	CB-CG-OD1	5.84	123.55	118.30
1	B	112	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	63	ARG	NE-CZ-NH1	5.59	123.10	120.30
1	B	195	PHE	CB-CG-CD2	-5.38	117.03	120.80
1	B	235	ASP	CB-CG-OD1	5.31	123.08	118.30
1	B	166	LYS	CD-CE-NZ	-5.28	99.56	111.70
1	A	195	PHE	CB-CG-CD2	-5.19	117.17	120.80
1	A	163	LEU	CB-CG-CD2	-5.14	102.27	111.00
1	A	117	ARG	CA-CB-CG	5.13	124.69	113.40
1	B	205	TRP	CZ3-CH2-CZ2	5.13	127.75	121.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2096	0	2109	17	0
1	B	2107	0	2120	42	0
2	A	8	0	14	0	0
2	B	8	0	14	0	0
3	A	12	0	18	7	0
3	B	16	0	24	8	0
4	B	15	0	17	1	0
5	A	232	0	0	8	0
5	B	220	0	0	5	0
All	All	4714	0	4316	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177[A]:LEU:HD21	1:B:191:ILE:CD1	1.59	1.31
1:B:63[B]:ARG:HE	1:B:82:GLN:NE2	1.45	1.15
1:B:177[A]:LEU:CD2	1:B:191:ILE:CD1	2.27	1.12
1:B:177[A]:LEU:CD2	1:B:191:ILE:HD12	1.81	1.09
1:B:177[A]:LEU:HD21	1:B:191:ILE:HD12	0.98	0.97
1:B:231:ASN:HB2	5:B:626:HOH:O	1.67	0.94
1:B:177[A]:LEU:CD2	1:B:191:ILE:HD11	2.04	0.85
1:B:121[B]:ASN:HD22	1:B:123:ASP:H	1.24	0.82
1:A:115:ASP:OD1	1:A:117:ARG:HG2	1.80	0.82
1:B:63[B]:ARG:NE	1:B:82:GLN:NE2	2.29	0.80
1:B:63[B]:ARG:HE	1:B:82:GLN:HE22	1.26	0.80
1:B:183:HIS:HD2	5:B:640:HOH:O	1.63	0.80
1:B:267[B]:ARG:NH2	3:B:406:EDO:H11	1.97	0.78
1:B:267[B]:ARG:NH2	3:B:406:EDO:O1	2.16	0.78
1:B:267[A]:ARG:HB3	1:B:295:GLU:HG3	1.70	0.73
1:B:267[B]:ARG:NH2	3:B:406:EDO:C1	2.52	0.72
1:B:231:ASN:CB	5:B:626:HOH:O	2.29	0.71
1:A:111:PRO:HA	5:A:698:HOH:O	1.92	0.68

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:177[A]:LEU:HD23	1:B:191:ILE:HD11	1.75	0.67
1:B:267[B]:ARG:HH22	3:B:406:EDO:H11	1.57	0.67
1:B:63[B]:ARG:NE	1:B:82:GLN:HE22	1.91	0.66
5:A:681:HOH:O	1:B:309:GLN:HG3	1.96	0.65
1:B:121[B]:ASN:ND2	1:B:123:ASP:H	1.97	0.61
1:A:224:GLU:OE2	5:A:629:HOH:O	2.16	0.61
1:A:293:GLN:NE2	5:A:690:HOH:O	2.26	0.60
1:A:305[B]:SER:HB2	3:A:404:EDO:H12	1.83	0.59
1:A:117:ARG:HH11	1:A:117:ARG:HG3	1.67	0.59
1:B:267[B]:ARG:HG3	1:B:267[B]:ARG:HH11	1.69	0.58
1:A:303:ALA:C	3:A:404:EDO:H11	2.27	0.56
1:A:117:ARG:HH11	1:A:117:ARG:CG	2.21	0.54
1:B:77:GLY:HA3	1:B:147:SER:HB3	1.89	0.54
1:B:177[A]:LEU:HD23	1:B:191:ILE:CD1	2.27	0.54
1:B:267[B]:ARG:CZ	3:B:406:EDO:O1	2.57	0.53
1:B:267[B]:ARG:HH22	3:B:406:EDO:C1	2.17	0.52
1:B:231:ASN:CG	5:B:626:HOH:O	2.46	0.50
1:B:272:PRO:HG3	3:B:406:EDO:H12	1.93	0.50
1:A:292:LEU:HD22	1:A:306:LEU:HG	1.93	0.49
1:A:77:GLY:HA3	1:A:147:SER:HB3	1.96	0.48
1:A:190[B]:GLU:OE2	5:A:685:HOH:O	2.20	0.47
5:A:681:HOH:O	1:B:309:GLN:CG	2.59	0.47
1:B:89:LEU:HB3	1:B:90:PRO:HD3	1.96	0.47
1:A:121:ASN:ND2	1:A:123:ASP:H	2.13	0.46
1:B:267[B]:ARG:HH11	1:B:267[B]:ARG:CG	2.28	0.46
1:B:216:ALA:CB	4:B:401:EPE:H82	2.45	0.46
1:A:70:ARG:HG2	5:A:677:HOH:O	2.16	0.46
1:B:269:VAL:O	3:B:406:EDO:H22	2.15	0.45
1:B:63[A]:ARG:HG2	1:B:65:VAL:HG13	1.99	0.45
1:A:233:ARG:HG2	1:B:90:PRO:HG2	1.99	0.44
1:B:191:ILE:HG22	1:B:195:PHE:CE2	2.55	0.42
1:B:246:LYS:HE3	1:B:246:LYS:HB3	1.54	0.42
1:B:175:ARG:NH2	1:B:177[B]:LEU:HD23	2.35	0.42
1:B:78:PHE:CE1	1:B:241[B]:CYS:SG	3.11	0.41
1:B:110:ASN:HB3	1:B:113:HIS:CD2	2.55	0.41
1:A:223[A]:GLN:HG3	5:A:708:HOH:O	2.20	0.41
1:B:167:LEU:O	1:B:260[B]:CYS:HA	2.21	0.41
1:A:305[B]:SER:HB2	3:A:404:EDO:C1	2.49	0.41
1:A:78:PHE:CZ	1:A:241[B]:CYS:SG	3.14	0.40
1:B:94:ARG:HD2	5:B:561:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	271/266 (102%)	266 (98%)	5 (2%)	0	100	100
1	B	272/266 (102%)	265 (97%)	7 (3%)	0	100	100
All	All	543/532 (102%)	531 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/219 (104%)	222 (98%)	5 (2%)	57	20
1	B	227/219 (104%)	225 (99%)	2 (1%)	82	56
All	All	454/438 (104%)	447 (98%)	7 (2%)	68	35

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LEU
1	A	94	ARG
1	A	117	ARG
1	A	121	ASN
1	A	229	LEU
1	B	246	LYS
1	B	264	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
1	A	193	GLN
1	B	58	GLN
1	B	61	ASN
1	B	82	GLN
1	B	183	HIS
1	B	264	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MPD	A	401	-	7,7,7	0.48	0	9,10,10	1.17	0
3	EDO	A	402	-	3,3,3	0.75	0	2,2,2	0.88	0
3	EDO	A	403	-	3,3,3	0.73	0	2,2,2	0.28	0
3	EDO	A	404	-	3,3,3	0.55	0	2,2,2	1.81	1 (50%)
4	EPE	B	401	-	15,15,15	1.15	2 (13%)	18,20,20	2.88	9 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MPD	B	402	-	7,7,7	0.57	0	9,10,10	0.88	0
3	EDO	B	403	-	3,3,3	0.57	0	2,2,2	0.68	0
3	EDO	B	404	-	3,3,3	0.53	0	2,2,2	1.03	0
3	EDO	B	405	-	3,3,3	1.11	0	2,2,2	1.11	0
3	EDO	B	406	-	3,3,3	1.56	1 (33%)	2,2,2	2.66	2 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	A	401	-	-	0/5/5/5	0/0/0/0
3	EDO	A	402	-	-	0/1/1/1	0/0/0/0
3	EDO	A	403	-	-	0/1/1/1	0/0/0/0
3	EDO	A	404	-	-	0/1/1/1	0/0/0/0
4	EPE	B	401	-	-	0/9/19/19	0/1/1/1
2	MPD	B	402	-	-	0/5/5/5	0/0/0/0
3	EDO	B	403	-	-	0/1/1/1	0/0/0/0
3	EDO	B	404	-	-	0/1/1/1	0/0/0/0
3	EDO	B	405	-	-	0/1/1/1	0/0/0/0
3	EDO	B	406	-	-	0/1/1/1	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	401	EPE	C6-N1	2.32	1.53	1.47
3	B	406	EDO	O2-C2	2.54	1.55	1.42
4	B	401	EPE	C10-S	2.58	1.81	1.77

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	EPE	C9-N1-C2	-3.67	101.84	111.26
4	B	401	EPE	O2S-S-C10	-3.54	103.75	106.79
4	B	401	EPE	O2S-S-O1S	-3.17	102.87	113.86
3	B	406	EDO	O2-C2-C1	-2.88	91.44	112.08
4	B	401	EPE	C8-C7-N4	-2.79	103.08	113.33
3	A	404	EDO	O1-C1-C2	-2.22	96.18	112.08
3	B	406	EDO	O1-C1-C2	2.42	129.41	112.08
4	B	401	EPE	C6-N1-C2	2.62	114.81	108.87
4	B	401	EPE	C9-N1-C6	2.65	118.04	111.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	EPE	O1S-S-C10	3.64	109.92	106.79
4	B	401	EPE	C5-N4-C3	5.42	121.14	108.87
4	B	401	EPE	O3S-S-C10	6.23	113.72	106.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	404	EDO	7	0
4	B	401	EPE	1	0
3	B	406	EDO	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/266 (99%)	-0.43	2 (0%) 86 86	6, 10, 21, 32	0
1	B	263/266 (98%)	-0.37	4 (1%) 74 74	5, 11, 21, 41	0
All	All	527/532 (99%)	-0.40	6 (1%) 80 81	5, 10, 21, 41	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	69	GLU	4.9
1	A	181	ASP	3.4
1	B	54	PRO	3.1
1	A	117	ARG	2.4
1	B	180	SER	2.2
1	B	181	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	EDO	B	406	4/4	0.67	0.21	5.84	17,25,30,34	0
4	EPE	B	401	15/15	0.93	0.16	2.90	21,24,30,32	0
3	EDO	A	403	4/4	0.91	0.11	2.12	14,19,20,22	0
2	MPD	B	402	8/8	0.92	0.10	2.11	15,19,22,23	0
3	EDO	A	404	4/4	0.91	0.11	1.25	19,20,26,40	0
3	EDO	B	405	4/4	0.98	0.08	0.66	13,16,18,23	0
2	MPD	A	401	8/8	0.95	0.07	-0.09	17,20,24,25	0
3	EDO	B	403	4/4	0.97	0.07	-0.23	10,12,13,13	0
3	EDO	A	402	4/4	0.96	0.07	-0.32	15,16,17,18	0
3	EDO	B	404	4/4	0.93	0.11	-	18,21,24,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.