



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 10:47 pm GMT

PDB ID : 3W16
Title : Structure of Aurora kinase A complexed to pyrazole-aminoquinoline inhibitor III
Authors : Oliveira, T.M.; Kairies, N.A.; Engh, R.A.
Deposited on : 2012-11-07
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	<div> <div>4%</div> <div>59%</div> <div>24%</div> <div>6%</div> <div>10%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aurora kinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	S	0	0	0
			2041	1318	354	365	4			

- Molecule 2 is 4-[(5-METHYL-1H-PYRAZOL-3-YL)AMINO]-2-PHENYLPHTHALAZIN-1(2H)-ONE (three-letter code: P9J) (formula: C₁₈H₁₅N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			24	18	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	8	Total	O	0	0
			8	8		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

4% 59% 24% 6% 10%

V324 G325 K326 P327 P328 E336 R340 R343 V344 E345 F346 T347 F348 D358 R362 N367 P368 S369 Q370 R375 R380 P381 W392 S388 LYS PRO SER ASN CYS GLN CYS ASN LYS GLU SER ALA SER LYS GLN SER E308 D307 P311 S314 L315 V316 G317 L318 C319 Y320

R126 E131 D132 K141 G142 K143 F144 G145 N146 R151 E152 K153 F157 L164 F165 K166 A167 Q168 L169 W170 K171 V174 E175 H176 Q177 L178 R179 R180 I184 Q185 R189 H190 P191 N192 I193 L194 R195 F200 K203 T204 R205 Y206 Y207 Y212 A213 P214 L225 S226

E230 T235 E239 L240 C247 K250 T253 H254 R255 D256 T257 K258 E269 L270 D274 R275 G276 W277 S278 W279 HIS ALA PRO SER SER ARG ARG THR THR LEU CYS G291 L293 G303 ALA ARG MET LYS HIS D307 E308 P311 S314 L315 V316 G317 L318 C319 Y320

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	83.11Å 83.11Å 169.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	23.99 – 2.80 23.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (23.99-2.80) 99.8 (23.33-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.41 (at 2.80Å)	Xtriage
Refinement program	phenix.refine	Depositor
R, R_{free}	0.231 , 0.322 0.222 , 0.301	Depositor DCC
R_{free} test set	419 reflections (4.63%)	DCC
Wilson B-factor (Å ²)	37.2	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2073	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P9J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	1/2090 (0.0%)	0.81	1/2823 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	336	GLU	CG-CD	5.95	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	318	LEU	CA-CB-CG	5.09	127.01	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2041	0	2046	54	0
2	A	24	0	15	3	0
3	A	8	0	0	1	1
All	All	2073	0	2061	57	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 14.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:NZ	1:A:291:GLY:HA2	1.55	1.20
1:A:258:LYS:HZ1	1:A:291:GLY:HA2	1.28	0.96
1:A:258:LYS:HZ2	1:A:291:GLY:HA2	1.29	0.95
1:A:195:ARG:HH11	1:A:195:ARG:HG3	1.34	0.90
1:A:144:PHE:HB3	1:A:169:LEU:HD21	1.59	0.83
1:A:190:HIS:HD2	1:A:192:ASN:H	1.23	0.83
1:A:307:ASP:O	1:A:308:GLU:HB3	1.84	0.75
1:A:254:HIS:HE1	1:A:274:ASP:OD2	1.70	0.74
1:A:195:ARG:HG3	1:A:195:ARG:NH1	2.03	0.73
1:A:258:LYS:HZ1	1:A:291:GLY:CA	2.01	0.73
1:A:375:ARG:HB3	1:A:375:ARG:HH11	1.54	0.72
1:A:170:GLU:OE2	1:A:175:GLU:HB3	1.89	0.71
1:A:144:PHE:HB2	1:A:164:LEU:HD23	1.74	0.69
1:A:235:THR:O	1:A:239:GLU:HG3	1.93	0.68
1:A:258:LYS:NZ	1:A:291:GLY:CA	2.47	0.65
1:A:254:HIS:CE1	1:A:274:ASP:OD2	2.49	0.64
1:A:190:HIS:HB3	1:A:193:ILE:HG13	1.82	0.62
1:A:195:ARG:HH11	1:A:195:ARG:CG	2.11	0.57
1:A:292:THR:O	1:A:292:THR:CG2	2.52	0.56
1:A:190:HIS:CD2	1:A:192:ASN:H	2.14	0.56
1:A:132:ASP:N	1:A:132:ASP:OD2	2.40	0.55
1:A:346:PHE:HE1	1:A:348:PHE:CZ	2.25	0.55
1:A:317:VAL:HG13	1:A:328:PRO:HD2	1.88	0.54
1:A:204:THR:OG1	1:A:205:ARG:NH1	2.41	0.53
1:A:367:ASN:HB3	1:A:370:GLN:HG3	1.88	0.53
1:A:180:ARG:O	1:A:184:ILE:HG13	2.09	0.53
1:A:185:GLN:NE2	1:A:275:PHE:CD1	2.78	0.52
1:A:131:GLU:O	1:A:153:LYS:HE3	2.10	0.52
1:A:320:TYR:O	1:A:324:VAL:HG13	2.10	0.52
1:A:225:LEU:O	1:A:226:SER:HB2	2.10	0.51
1:A:340:ARG:NH1	1:A:345:GLU:O	2.43	0.51
1:A:230:GLU:HG3	3:A:604:HOH:O	2.12	0.50
1:A:375:ARG:HB3	1:A:375:ARG:NH1	2.26	0.48
1:A:293:LEU:HA	1:A:293:LEU:HD23	1.69	0.48
1:A:190:HIS:CD2	1:A:191:PRO:HD2	2.48	0.47
2:A:501:P9J:H19	2:A:501:P9J:N10	2.30	0.47
1:A:358:ASP:O	1:A:362:ARG:HG3	2.15	0.47
1:A:256:ASP:OD1	1:A:291:GLY:HA3	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:GLU:O	1:A:269:GLU:HG2	2.15	0.46
1:A:367:ASN:HD22	1:A:368:PRO:HD2	1.81	0.46
1:A:311:ASP:O	1:A:314:SER:HB2	2.17	0.45
1:A:240:LEU:HG	1:A:270:LEU:HD11	1.99	0.45
1:A:380:HIS:CD2	1:A:382:TRP:H	2.34	0.45
2:A:501:P9J:O24	2:A:501:P9J:C12	2.63	0.45
1:A:190:HIS:HD2	1:A:192:ASN:N	2.03	0.45
1:A:292:THR:O	1:A:292:THR:HG23	2.17	0.44
1:A:380:HIS:HD2	1:A:382:TRP:H	1.66	0.44
1:A:200:PHE:CE2	1:A:207:TYR:CD1	3.07	0.43
1:A:212:TYR:CZ	1:A:214:PRO:HB3	2.54	0.43
1:A:326:LYS:HB2	1:A:327:PRO:HD2	2.00	0.43
2:A:501:P9J:N10	2:A:501:P9J:C19	2.82	0.42
1:A:166:LYS:HB2	1:A:204:THR:O	2.19	0.42
1:A:179:ARG:HB2	1:A:179:ARG:HH11	1.83	0.42
1:A:166:LYS:HG3	1:A:206:VAL:HG23	2.02	0.41
1:A:247:CYS:SG	1:A:277:TRP:CZ3	3.13	0.41
1:A:151:ARG:HA	1:A:157:PHE:O	2.21	0.40
1:A:190:HIS:CG	1:A:191:PRO:HD2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:606:HOH:O	3:A:607:HOH:O[12_565]	1.74	0.46

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	243/278 (87%)	223 (92%)	13 (5%)	7 (3%)	5 18

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	175	GLU
1	A	292	THR
1	A	293	LEU
1	A	255	ARG
1	A	166	LYS
1	A	308	GLU
1	A	381	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	218/246 (89%)	190 (87%)	28 (13%)	5 15

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	131	GLU
1	A	132	ASP
1	A	141	LYS
1	A	143	LYS
1	A	144	PHE
1	A	146	ASN
1	A	151	ARG
1	A	165	PHE
1	A	170	GLU
1	A	174	VAL
1	A	178	LEU
1	A	189	ARG
1	A	194	LEU
1	A	195	ARG
1	A	204	THR
1	A	250	LYS
1	A	253	ILE
1	A	255	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	292	THR
1	A	308	GLU
1	A	315	LEU
1	A	318	LEU
1	A	324	VAL
1	A	336	GLU
1	A	340	ARG
1	A	343	ARG
1	A	375	ARG
1	A	388	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	GLN
1	A	190	HIS
1	A	192	ASN
1	A	242	ASN
1	A	366	HIS
1	A	367	ASN
1	A	370	GLN
1	A	380	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	P9J	A	501	-	23,27,27	1.55	3 (13%)	25,38,38	1.31	4 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	P9J	A	501	-	-	0/6/8/8	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	P9J	C11-N2	-4.45	1.34	1.44
2	A	501	P9J	C9-C8	-2.95	1.41	1.44
2	A	501	P9J	C19-C20	-2.41	1.34	1.39

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	P9J	C8-C9-N17	-2.30	117.66	119.67
2	A	501	P9J	C15-C16-C11	2.37	121.67	118.66
2	A	501	P9J	C23-C20-N21	2.42	124.87	120.07
2	A	501	P9J	C12-C11-N2	2.84	123.60	119.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	P9J	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/278 (89%)	-0.01	10 (4%) 39 28	8, 25, 58, 66	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	GLY	3.7
1	A	203	ALA	3.1
1	A	303	GLY	2.7
1	A	291	GLY	2.6
1	A	279	VAL	2.6
1	A	126	ARG	2.4
1	A	177	GLN	2.2
1	A	171	LYS	2.2
1	A	167	ALA	2.1
1	A	143	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	P9J	A	501	24/24	0.97	0.13	-0.46	11,15,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.