



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 12, 2018 – 02:21 PM EST

PDB ID : 4W2R  
Title : Structure of Hs/AcPRC2 in complex with 5,8-dichloro-2-[(4-methoxy-6-methyl-2-oxo-1,2-dihydropyridin-3-yl)methyl]-7-[(R)-methoxy(oxetan-3-yl)methyl]-3,4-dihydroisoquinolin-1(2H)-one  
Authors : Gajiwala, K.S.; Brooun, A.; Liu, W.; Deng, Y.; Stewart, A.E.  
Deposited on : 2017-09-25  
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2917 (2.84-2.80)
Clashscore	112137	3382 (2.84-2.80)
Ramachandran outliers	110173	3324 (2.84-2.80)
Sidechain outliers	110143	3326 (2.84-2.80)
RSRZ outliers	101464	2948 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	643	<div> <div>0.1%</div> <div>39%</div> <div>29%</div> <div>0.1%</div> <div>28%</div> </div>
1	B	643	<div> <div>2%</div> <div>42%</div> <div>27%</div> <div>0.1%</div> <div>28%</div> </div>
2	E	362	<div> <div>0.1%</div> <div>60%</div> <div>36%</div> <div>0.1%</div> <div>0.1%</div> </div>
2	F	362	<div> <div>0.1%</div> <div>60%</div> <div>36%</div> <div>0.1%</div> <div>0.1%</div> </div>
3	S	191	<div> <div>0.1%</div> <div>31%</div> <div>31%</div> <div>5%</div> <div>33%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	T	191	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZN	B	705	-	-	-	X
4	ZN	B	706	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enhancer of zeste 2 polycomb repressive complex 2 subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	463	Total	C	N	O	S	0	0	0
			3769	2365	678	688	38			
1	B	465	Total	C	N	O	S	0	0	0
			3777	2368	679	692	38			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	LEU	-	linker	UNP G1KPH4
A	390	GLY	-	linker	UNP G1KPH4
A	391	GLY	-	linker	UNP G1KPH4
A	392	GLY	-	linker	UNP G1KPH4
A	393	GLY	-	linker	UNP G1KPH4
A	394	SER	-	linker	UNP G1KPH4
A	395	GLY	-	linker	UNP G1KPH4
A	396	GLY	-	linker	UNP G1KPH4
A	397	GLY	-	linker	UNP G1KPH4
A	398	GLY	-	linker	UNP G1KPH4
A	399	SER	-	linker	UNP G1KPH4
A	400	GLY	-	linker	UNP G1KPH4
A	401	GLY	-	linker	UNP G1KPH4
A	402	GLY	-	linker	UNP G1KPH4
A	403	GLY	-	linker	UNP G1KPH4
A	404	SER	-	linker	UNP G1KPH4
A	405	ALA	-	linker	UNP G1KPH4
A	406	ALA	-	linker	UNP G1KPH4
A	407	ALA	-	linker	UNP G1KPH4
B	389	LEU	-	linker	UNP G1KPH4
B	390	GLY	-	linker	UNP G1KPH4
B	391	GLY	-	linker	UNP G1KPH4
B	392	GLY	-	linker	UNP G1KPH4
B	393	GLY	-	linker	UNP G1KPH4
B	394	SER	-	linker	UNP G1KPH4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	395	GLY	-	linker	UNP G1KPH4
B	396	GLY	-	linker	UNP G1KPH4
B	397	GLY	-	linker	UNP G1KPH4
B	398	GLY	-	linker	UNP G1KPH4
B	399	SER	-	linker	UNP G1KPH4
B	400	GLY	-	linker	UNP G1KPH4
B	401	GLY	-	linker	UNP G1KPH4
B	402	GLY	-	linker	UNP G1KPH4
B	403	GLY	-	linker	UNP G1KPH4
B	404	SER	-	linker	UNP G1KPH4
B	405	ALA	-	linker	UNP G1KPH4
B	406	ALA	-	linker	UNP G1KPH4
B	407	ALA	-	linker	UNP G1KPH4

- Molecule 2 is a protein called Polycomb protein EED.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	358	Total	C	N	O	S	0	0	0
			2898	1834	509	534	21			
2	F	359	Total	C	N	O	S	0	0	0
			2906	1840	510	535	21			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	66	MET	-	expression tag	UNP O75530
F	66	MET	-	expression tag	UNP O75530

- Molecule 3 is a protein called Polycomb protein SUZ12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	S	128	Total	C	N	O	S	0	0	0
			1070	675	185	198	12			
3	T	128	Total	C	N	O	S	0	0	0
			1070	675	185	198	12			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	535	MET	-	expression tag	UNP Q15022
S	536	ASP	-	expression tag	UNP Q15022
S	537	TYR	-	expression tag	UNP Q15022

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
S	538	LYS	-	expression tag	UNP Q15022
S	539	ASP	-	expression tag	UNP Q15022
S	540	ASP	-	expression tag	UNP Q15022
S	541	ASP	-	expression tag	UNP Q15022
S	542	ASP	-	expression tag	UNP Q15022
S	543	LYS	-	expression tag	UNP Q15022
S	544	GLY	-	expression tag	UNP Q15022
S	583	ASP	SER	conflict	UNP Q15022
T	535	MET	-	expression tag	UNP Q15022
T	536	ASP	-	expression tag	UNP Q15022
T	537	TYR	-	expression tag	UNP Q15022
T	538	LYS	-	expression tag	UNP Q15022
T	539	ASP	-	expression tag	UNP Q15022
T	540	ASP	-	expression tag	UNP Q15022
T	541	ASP	-	expression tag	UNP Q15022
T	542	ASP	-	expression tag	UNP Q15022
T	543	LYS	-	expression tag	UNP Q15022
T	544	GLY	-	expression tag	UNP Q15022
T	583	ASP	SER	conflict	UNP Q15022

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	7	Total	Zn	0	0
			7	7		
4	A	7	Total	Zn	0	0
			7	7		

- Molecule 5 is 5,8-dichloro-2-[(4-methoxy-6-methyl-2-oxo-1,2-dihydropyridin-3-yl)methyl]-7-[(R)-methoxy(oxetan-3-yl)methyl]-3,4-dihydroisoquinolin-1(2H)-one (three-letter code: CJD) (formula: C<sub>22</sub>H<sub>24</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 31	C 22	Cl 2	N 2	O 5	0	0
5	B	1	Total 31	C 22	Cl 2	N 2	O 5	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Enhancer of zeste 2 polycomb repressive complex 2 subunit




- Molecule 1: Enhancer of zeste 2 polycomb repressive complex 2 subunit







• Molecule 2: Polycomb protein EED



• Molecule 2: Polycomb protein EED





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	70.72Å 115.48Å 151.54Å 90.00° 102.40° 90.00°	Depositor
Resolution (Å)	148.01 – 2.81 148.00 – 2.81	Depositor EDS
% Data completeness (in resolution range)	99.8 (148.01-2.81) 99.8 (148.00-2.81)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.82Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.218 , 0.266 0.214 , 0.261	Depositor DCC
$R_{free}$ test set	2938 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	74.6	Xtriage
Anisotropy	0.394	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	15566	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CJD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3850	0.52	0/5176
1	B	0.40	0/3856	0.53	0/5184
2	E	0.40	0/2972	0.57	0/4026
2	F	0.39	0/2980	0.58	1/4037 (0.0%)
3	S	0.39	0/1091	0.51	0/1464
3	T	0.40	0/1091	0.52	0/1464
All	All	0.40	0/15840	0.54	1/21351 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
3	T	0	1
All	All	0	4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	201	LEU	CA-CB-CG	5.50	127.95	115.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	250	PRO	Peptide
1	A	559	MET	Peptide
1	B	559	MET	Peptide
3	T	577	GLN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3769	0	3657	203	0
1	B	3777	0	3673	185	0
2	E	2898	0	2812	159	0
2	F	2906	0	2823	149	0
3	S	1070	0	1052	83	0
3	T	1070	0	1052	78	0
4	A	7	0	0	0	0
4	B	7	0	0	0	0
5	A	31	0	0	0	0
5	B	31	0	0	0	0
All	All	15566	0	15069	777	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 777 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:165:THR:HG22	2:E:167:GLN:HG2	1.32	1.11
2:F:340:GLY:HA3	2:F:382:PRO:HG3	1.40	1.03
2:F:165:THR:HG22	2:F:167:GLN:HG2	1.40	1.02
2:E:340:GLY:HA3	2:E:382:PRO:HG3	1.45	0.96
1:A:96:ILE:HG12	1:A:97:PRO:HD2	1.43	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	449/643 (70%)	396 (88%)	43 (10%)	10 (2%)	8	25
1	B	451/643 (70%)	408 (90%)	36 (8%)	7 (2%)	11	34
2	E	356/362 (98%)	335 (94%)	20 (6%)	1 (0%)	44	76
2	F	357/362 (99%)	335 (94%)	20 (6%)	2 (1%)	28	61
3	S	126/191 (66%)	109 (86%)	12 (10%)	5 (4%)	3	10
3	T	126/191 (66%)	108 (86%)	13 (10%)	5 (4%)	3	10
All	All	1865/2392 (78%)	1691 (91%)	144 (8%)	30 (2%)	11	34

5 of 30 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	S	605	ASP
3	S	607	ASN
3	T	578	GLU
3	T	605	ASP
1	A	10	LYS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	418/560 (75%)	388 (93%)	30 (7%)	17	42
1	B	420/560 (75%)	392 (93%)	28 (7%)	19	46
2	E	321/325 (99%)	305 (95%)	16 (5%)	28	60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	F	322/325 (99%)	310 (96%)	12 (4%)	39 72
3	S	122/175 (70%)	109 (89%)	13 (11%)	8 22
3	T	122/175 (70%)	112 (92%)	10 (8%)	13 36
All	All	1725/2120 (81%)	1616 (94%)	109 (6%)	21 50

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	422	HIS
2	E	254	LYS
3	T	579	MET
1	B	460	LYS
1	B	567	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 60 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	429	ASN
1	B	545	GLN
3	S	664	HIS
1	B	504	GLN
1	B	603	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 16 ligands modelled in this entry, 14 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	CJD	A	708	-	30,34,34	2.06	9 (30%)	31,50,50	1.84	3 (9%)
5	CJD	B	708	-	30,34,34	2.07	10 (33%)	31,50,50	1.94	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CJD	A	708	-	-	0/12/35/35	0/4/4/4
5	CJD	B	708	-	-	0/12/35/35	0/4/4/4

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	708	CJD	C3-C4	2.30	1.43	1.39
5	A	708	CJD	C16-N17	2.44	1.39	1.34
5	B	708	CJD	C5-C4	2.48	1.45	1.40
5	B	708	CJD	C5-C6	2.52	1.45	1.39
5	A	708	CJD	C5-C8	2.58	1.52	1.47

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	708	CJD	O21-C14-C15	-3.19	118.79	124.17
5	A	708	CJD	O21-C14-C15	-2.98	119.15	124.17
5	B	708	CJD	O23-C8-C5	-2.49	119.58	124.23
5	B	708	CJD	C2-C3-C4	-2.13	119.86	123.39
5	A	708	CJD	C11-C10-N9	2.28	112.97	109.91

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	463/643 (72%)	0.20	9 (1%) 67 58	43, 74, 104, 121	0
1	B	465/643 (72%)	0.19	15 (3%) 48 37	42, 74, 106, 121	0
2	E	358/362 (98%)	0.15	3 (0%) 86 81	42, 63, 87, 122	0
2	F	359/362 (99%)	0.10	5 (1%) 75 69	45, 64, 87, 122	0
3	S	128/191 (67%)	0.23	2 (1%) 72 65	51, 79, 114, 120	0
3	T	128/191 (67%)	0.29	3 (2%) 61 51	47, 76, 113, 124	0
All	All	1901/2392 (79%)	0.18	37 (1%) 67 58	42, 69, 104, 124	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	383	HIS	5.9
1	B	367	VAL	4.1
2	F	385	ALA	3.6
1	A	166	ILE	3.5
2	F	383	HIS	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	ZN	B	705	1/1	0.91	0.23	2.89	79,79,79,79	1
4	ZN	B	706	1/1	0.98	0.20	2.61	72,72,72,72	1
4	ZN	B	701	1/1	0.97	0.20	1.96	78,78,78,78	1
4	ZN	A	701	1/1	0.97	0.19	1.60	75,75,75,75	1
4	ZN	A	703	1/1	0.98	0.19	0.91	77,77,77,77	1
4	ZN	B	704	1/1	0.98	0.17	0.84	54,54,54,54	1
4	ZN	A	706	1/1	0.99	0.17	0.81	58,58,58,58	1
5	CJD	B	708	31/31	0.97	0.22	0.80	42,56,63,66	0
4	ZN	A	705	1/1	0.98	0.21	0.78	65,65,65,65	1
4	ZN	B	703	1/1	0.95	0.18	0.72	72,72,72,72	1
5	CJD	A	708	31/31	0.97	0.22	0.56	43,53,61,65	0
4	ZN	A	704	1/1	0.98	0.16	0.27	74,74,74,74	1
4	ZN	B	707	1/1	0.99	0.18	0.09	60,60,60,60	1
4	ZN	B	702	1/1	0.98	0.16	-0.36	71,71,71,71	1
4	ZN	A	707	1/1	0.99	0.17	-0.52	63,63,63,63	1
4	ZN	A	702	1/1	1.00	0.12	-1.46	65,65,65,65	1

## 6.5 Other polymers ⓘ

There are no such residues in this entry.