



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 11:21 AM EST

PDB ID : 5W2U  
Title : INFLUENZA VIRUS NEURAMINIDASE N9 IN COMPLEX WITH 7-DEOXYGENATED 2,3-DIFLUORO-N-ACETYLNEURAMINIC ACID  
Authors : Streltsov, V.A.; Mckimm-Breschkin, J.; Barrett, S.; Pilling, P.; Hader, S.; Watt, A.G.  
Deposited on : 2017-06-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	9SM	A	501	-	-	-	X
5	MAN	A	506	-	-	-	X

## 2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 3715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3094	1928	548	595	23	0	9	0

- Molecule 2 is (2 {R},3 {R},4 {R},5 {R},6 {S})-5-acetamido-6-[(2 {S})-2,3-bis(oxidanyl)propyl]-2,3-bis(fluoranyl)-4-oxidanyl-oxane-2-carboxylic acid (three-letter code: 9SM) (formula:  $C_{11}H_{17}F_2NO_7$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	F	N	O		
2	A	1	21	11	2	1	7	0	0

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is (2 {S},3 {R},4 {R},5 {R})-3-acetamido-2-[(2 {S})-2,3-bis(oxidanyl)propyl]-5-fluoranyl-4-oxidanyl-2,3,4,5-tetrahydropyran-1-ium-6-carboxylic acid (three-letter code: 9S4) (formula: C<sub>11</sub>H<sub>17</sub>FN<sub>7</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	F	N	O	0	1
			20	11	1	1	7		

- Molecule 8 is 5-(acetylamino)-2,6-anhydro-3,5,7-trideoxy-3-fluoro-D-threo-L-galacto-nononic acid (three-letter code: 9SJ) (formula:  $C_{11}H_{18}FNO_7$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	F	N	O	0	1
			20	11	1	1	7		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	423	Total 423	O 423	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neuraminidase





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	181.03Å 181.03Å 181.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.38 – 2.00 45.26 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.38-2.00) 99.6 (45.26-2.00)	Depositor EDS
$R_{merge}$	0.31	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.142 , 0.187 0.144 , 0.191	Depositor DCC
$R_{free}$ test set	1723 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 61.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3715	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9SJ, BMA, NAG, 9SM, CA, 9S4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.09	3/3226 (0.1%)	0.96	10/4391 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	372	SER	CB-OG	-6.34	1.34	1.42
1	A	132	TYR	CE1-CZ	5.37	1.45	1.38
1	A	316	GLN	CD-NE2	5.06	1.45	1.32

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	NE-CZ-NH1	8.13	124.37	120.30
1	A	447[A]	MET	CG-SD-CE	7.62	112.40	100.20
1	A	447[B]	MET	CG-SD-CE	7.62	112.40	100.20
1	A	173	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	A	125	ASP	CB-CG-OD1	6.69	124.32	118.30
1	A	210	ARG	NE-CZ-NH1	-6.04	117.28	120.30
1	A	435	ASP	CB-CG-OD1	5.67	123.40	118.30
1	A	210	ARG	NE-CZ-NH2	5.41	123.01	120.30
1	A	339	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	A	325	ASP	CB-CG-OD1	5.34	123.11	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3094	0	2925	27	1
2	A	21	0	0	0	0
3	A	70	0	62	1	0
4	A	11	0	8	0	0
5	A	55	0	47	0	0
6	A	1	0	0	0	0
7	A	20	0	0	0	0
8	A	20	0	0	0	0
9	A	423	0	0	4	7
All	All	3715	0	3042	27	7

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:LEU:H	1:A:234:HIS:HD2	1.14	0.95
1:A:168:THR:H	1:A:171:ASN:HD21	1.14	0.88
1:A:173:ARG:HD3	1:A:210:ARG:NH1	1.92	0.84
1:A:87:LEU:H	1:A:234:HIS:CD2	2.03	0.76
1:A:285[B]:ARG:HD2	9:A:602:HOH:O	1.91	0.70
1:A:372:SER:HB2	1:A:399:ASN:ND2	2.07	0.69
1:A:140[B]:ILE:HD11	9:A:681:HOH:O	1.94	0.67
1:A:98:HIS:HE1	1:A:420:ARG:HH11	1.45	0.63
1:A:168:THR:H	1:A:171:ASN:ND2	1.94	0.62
1:A:168:THR:OG1	1:A:171:ASN:ND2	2.33	0.61
1:A:173:ARG:CD	1:A:210:ARG:NH1	2.64	0.60
1:A:97:TRP:H	1:A:394:GLN:HE22	1.50	0.60
1:A:168:THR:N	1:A:171:ASN:HD21	1.95	0.59
1:A:98:HIS:CE1	1:A:420:ARG:HH11	2.22	0.57
1:A:98:HIS:HD2	1:A:99:ILE:O	1.87	0.57
1:A:285[B]:ARG:NH2	9:A:601:HOH:O	2.16	0.55
1:A:285[B]:ARG:NH1	9:A:602:HOH:O	2.18	0.54
1:A:173:ARG:HD3	1:A:210:ARG:CZ	2.39	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:372:SER:HB2	1:A:399:ASN:HD21	1.74	0.52
1:A:171:ASN:HD22	1:A:171:ASN:H	1.57	0.51
1:A:279:CYS:HB3	1:A:290:CYS:HB3	1.95	0.48
1:A:121:TYR:CG	1:A:229:SER:HA	2.49	0.47
1:A:229:SER:HB3	1:A:350:LYS:HE2	1.96	0.46
1:A:287:GLU:OE2	1:A:305[B]:ARG:HD3	2.20	0.42
1:A:189:THR:HG21	1:A:209:ASN:HB2	2.01	0.42
1:A:438:TRP:CD1	3:A:512:NAG:H82	2.56	0.41
1:A:208:TYR:CZ	1:A:260:GLU:HA	2.56	0.41

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:650:HOH:O	9:A:982:HOH:O[15_555]	1.89	0.31
9:A:632:HOH:O	9:A:995:HOH:O[15_555]	1.92	0.28
9:A:953:HOH:O	9:A:964:HOH:O[9_555]	1.92	0.28
9:A:723:HOH:O	9:A:982:HOH:O[15_555]	1.97	0.23
9:A:993:HOH:O	9:A:995:HOH:O[15_555]	1.97	0.23
9:A:640:HOH:O	9:A:959:HOH:O[15_555]	2.06	0.14
1:A:171:ASN:CB	9:A:837:HOH:O[16_555]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	394/388 (102%)	379 (96%)	15 (4%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	350/341 (103%)	340 (97%)	10 (3%)	48 47

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82[A]	ARG
1	A	82[B]	ARG
1	A	171	ASN
1	A	210	ARG
1	A	222	ASN
1	A	231	CYS
1	A	334	VAL
1	A	372	SER
1	A	415	GLU
1	A	457	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	HIS
1	A	171	ASN
1	A	222	ASN
1	A	234	HIS
1	A	345	ASN
1	A	346	ASN
1	A	392	GLN
1	A	394	GLN
1	A	399	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	9SM	A	501	-	15,21,21	0.97	2 (13%)	16,31,31	3.34	6 (37%)
3	NAG	A	502	1,3	14,14,15	0.79	0	15,19,21	2.35	6 (40%)
3	NAG	A	503	3,4	14,14,15	0.81	1 (7%)	15,19,21	1.88	4 (26%)
4	BMA	A	504	3,5	11,11,12	0.79	0	13,15,17	1.31	3 (23%)
5	MAN	A	505	5,4	11,11,12	1.07	1 (9%)	13,15,17	2.52	5 (38%)
5	MAN	A	506	5	11,11,12	0.84	0	13,15,17	1.31	2 (15%)
5	MAN	A	507	5	11,11,12	1.00	1 (9%)	13,15,17	1.18	1 (7%)
5	MAN	A	508	5,4	11,11,12	0.65	0	13,15,17	1.59	2 (15%)
5	MAN	A	509	5	11,11,12	1.00	0	13,15,17	1.99	4 (30%)
3	NAG	A	510	1,3	14,14,15	0.79	0	15,19,21	1.31	2 (13%)
3	NAG	A	511	3	14,14,15	1.35	2 (14%)	15,19,21	2.05	4 (26%)
3	NAG	A	512	1	14,14,15	0.67	0	15,19,21	2.04	5 (33%)
7	9S4	A	514[B]	-	14,20,20	0.74	0	14,28,28	1.33	3 (21%)
8	9SJ	A	515[A]	1	17,20,20	0.86	1 (5%)	18,28,28	2.58	7 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9SM	A	501	-	-	0/10/39/39	0/1/1/1
3	NAG	A	502	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	503	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	504	3,5	-	0/2/19/22	0/1/1/1
5	MAN	A	505	5,4	-	0/2/19/22	0/1/1/1
5	MAN	A	506	5	-	0/2/19/22	0/1/1/1
5	MAN	A	507	5	-	0/2/19/22	0/1/1/1
5	MAN	A	508	5,4	-	0/2/19/22	0/1/1/1
5	MAN	A	509	5	-	0/2/19/22	0/1/1/1
3	NAG	A	510	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	511	3	-	0/6/23/26	0/1/1/1
3	NAG	A	512	1	-	0/6/23/26	0/1/1/1
7	9S4	A	514[B]	-	-	0/10/34/34	0/0/1/1
8	9SJ	A	515[A]	1	-	0/10/34/34	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	A	515[A]	9SJ	F1-C3	-2.56	1.33	1.40
5	A	507	MAN	C2-C3	2.00	1.55	1.52
2	A	501	9SM	CAQ-CAP	2.03	1.56	1.52
2	A	501	9SM	OAF-CAP	2.08	1.49	1.43
3	A	511	NAG	C2-N2	2.18	1.50	1.46
5	A	505	MAN	O5-C5	2.35	1.48	1.43
3	A	503	NAG	C1-C2	2.47	1.55	1.52
3	A	511	NAG	C1-C2	2.98	1.56	1.52

All (54) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	515[A]	9SJ	F1-C3-C2	-8.40	96.74	108.00
5	A	505	MAN	O2-C2-C1	-5.26	98.48	109.18
3	A	502	NAG	C1-C2-N2	-5.02	101.92	110.49
3	A	512	NAG	C2-N2-C7	-4.88	115.83	122.94
2	A	501	9SM	CAU-CAQ-CAP	-4.24	104.99	114.63
5	A	505	MAN	C6-C5-C4	-3.99	103.66	113.00
5	A	505	MAN	O6-C6-C5	-3.84	98.40	111.34
2	A	501	9SM	OAM-CAU-CAQ	-3.75	101.15	105.95
3	A	502	NAG	C8-C7-N2	-3.53	109.73	116.11
8	A	515[A]	9SJ	C6-C7-C8	-3.41	106.89	114.63
8	A	515[A]	9SJ	C4-C5-N5	-3.37	104.16	110.61
3	A	503	NAG	O6-C6-C5	-3.35	100.06	111.34
5	A	509	MAN	C3-C4-C5	-3.24	104.51	110.22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	508	MAN	O4-C4-C3	-3.22	103.35	110.36
3	A	502	NAG	C3-C4-C5	-3.18	104.62	110.22
3	A	510	NAG	O6-C6-C5	-3.17	100.68	111.34
3	A	502	NAG	O4-C4-C3	-2.84	104.17	110.36
2	A	501	9SM	OAG-CAR-CAS	-2.76	103.87	109.15
2	A	501	9SM	OAG-CAR-CAT	-2.70	104.15	109.61
7	A	514[B]	9S4	C4-C5-N5	-2.64	105.54	110.61
3	A	502	NAG	C2-N2-C7	-2.55	119.22	122.94
5	A	506	MAN	O4-C4-C3	-2.55	104.81	110.36
4	A	504	BMA	O2-C2-C3	-2.54	105.18	110.17
3	A	512	NAG	O4-C4-C3	-2.52	104.88	110.36
4	A	504	BMA	O3-C3-C4	-2.44	105.05	110.36
7	A	514[B]	9S4	C6-C7-C8	-2.39	109.19	114.63
8	A	515[A]	9SJ	F1-C3-C4	-2.36	106.77	108.49
5	A	506	MAN	C1-C2-C3	-2.31	106.72	109.65
7	A	514[B]	9S4	O4-C4-C3	-2.29	104.82	109.15
4	A	504	BMA	C6-C5-C4	-2.22	107.81	113.00
8	A	515[A]	9SJ	C4-C5-C6	-2.21	106.31	110.44
5	A	509	MAN	O5-C1-C2	-2.18	107.37	110.79
8	A	515[A]	9SJ	O6-C2-C3	-2.16	104.77	109.24
5	A	507	MAN	O2-C2-C1	-2.08	104.94	109.18
3	A	511	NAG	O3-C3-C4	-2.03	105.94	110.36
5	A	505	MAN	O2-C2-C3	2.10	114.30	110.17
3	A	510	NAG	C8-C7-N2	2.24	120.16	116.11
5	A	509	MAN	O2-C2-C3	2.40	114.88	110.17
3	A	503	NAG	C8-C7-N2	2.41	120.46	116.11
8	A	515[A]	9SJ	C6-O6-C2	2.41	116.67	112.06
3	A	512	NAG	C4-C3-C2	2.43	114.58	111.02
3	A	512	NAG	C8-C7-N2	2.62	120.84	116.11
3	A	511	NAG	C1-O5-C5	2.68	115.86	112.17
3	A	503	NAG	C1-C2-N2	2.98	115.58	110.49
3	A	512	NAG	O5-C1-C2	3.04	115.70	111.47
3	A	502	NAG	C1-O5-C5	3.07	116.40	112.17
3	A	511	NAG	O4-C4-C3	3.24	117.41	110.36
3	A	503	NAG	C6-C5-C4	3.47	121.12	113.00
5	A	508	MAN	C1-O5-C5	3.80	117.40	112.17
5	A	505	MAN	C1-O5-C5	3.81	117.42	112.17
5	A	509	MAN	C1-O5-C5	4.05	117.75	112.17
2	A	501	9SM	OAF-CAP-CAQ	5.16	120.80	109.14
3	A	511	NAG	C2-N2-C7	5.34	130.73	122.94
2	A	501	9SM	FAI-CAS-CAR	9.39	115.37	108.49

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	512	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/388 (100%)	-0.47	2 (0%) 90 90	17, 22, 32, 54	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	82[A]	ARG	2.8
1	A	83	ASP	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	9SM	A	501	21/21	0.93	0.19	4.12	28,38,43,50	0
5	MAN	A	506	11/12	0.97	0.12	2.41	25,26,29,29	0
4	BMA	A	504	11/12	0.99	0.08	1.44	23,25,28,31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	NAG	A	502	14/15	0.96	0.11	0.83	19,25,39,40	0
7	9S4	A	514[B]	20/20	0.97	0.10	0.65	21,26,30,45	20
8	9SJ	A	515[A]	20/20	0.97	0.10	0.56	19,21,23,25	20
5	MAN	A	507	11/12	0.98	0.07	-1.01	23,25,28,28	0
6	CA	A	513	1/1	0.99	0.05	-1.69	25,25,25,25	0
5	MAN	A	508	11/12	0.96	0.19	-	31,38,45,46	0
3	NAG	A	503	14/15	0.98	0.10	-	20,26,32,43	0
5	MAN	A	505	11/12	0.98	0.10	-	21,25,32,35	0
3	NAG	A	511	14/15	0.78	0.43	-	54,64,69,74	0
5	MAN	A	509	11/12	0.77	0.40	-	58,66,72,77	0
3	NAG	A	510	14/15	0.94	0.25	-	32,35,42,52	0
3	NAG	A	512	14/15	0.93	0.33	-	45,52,63,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.