



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2017 – 02:10 AM EST

PDB ID : 4W5O  
Title : The Crystal Structure of Human Argonaute2 Bound to a Guide and Target RNA Containing Seed Pairing from 2-9  
Authors : Schirle, N.T.; MacRae, I.J.  
Deposited on : unknown  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)
RNA backbone	2435	1035 (2.60-1.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	859	<div> <div>11%</div> <div> <div></div> <div>85%</div> <div>8%</div> <div>•</div> <div>7%</div> </div> </div>
2	B	21	<div> <div>5%</div> <div> <div></div> <div>71%</div> <div>19%</div> <div>5%</div> <div>5%</div> </div> </div>
3	D	11	<div> <div></div> <div> <div></div> <div>91%</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	IPA	A	902	-	-	-	X
5	IPA	A	903	-	-	-	X

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein argonaute-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	801	Total	C	N	O	S	0	0	0
			6419	4089	1154	1136	40			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	387	ASP	SER	engineered mutation	UNP Q9UKV8

- Molecule 2 is a RNA chain called RNA (5'-R(P\*UP\*UP\*CP\*AP\*CP\*AP\*UP\*UP\*GP\*CP\*CP\*CP\*AP\*AP\*GP\*UP\*CP\*UP\*UP\*U)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	20	Total	C	N	O	P	0	0	0
			399	177	62	140	20			


- Molecule 3 is a RNA chain called RNA (5'-R(\*CP\*AP\*AP\*UP\*GP\*UP\*GP\*AP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	10	Total	C	N	O	P	0	0	0
			196	88	37	62	9			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		
4	D	1	Total	Mg	0	0
			1	1		

- Molecule 5 is ISOPROPYL ALCOHOL (three-letter code: IPA) (formula:  $C_3H_8O$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	3	1		
5	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 6 is PHENOL (three-letter code: IPH) (formula:  $C_6H_6O$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	6	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	6	1		
6	A	1	Total	C	O	0	0
			7	6	1		
6	A	1	Total	C	O	0	0
			7	6	1		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	354	Total	O	0	0
			354	354		
8	B	45	Total	O	0	0
			45	45		
8	D	34	Total	O	0	0
			34	34		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Protein argonaute-2



#### • Molecule 2: RNA (5'-R(P\*UP\*UP\*CP\*AP\*CP\*AP\*UP\*UP\*GP\*CP\*CP\*CP\*AP\*AP\*GP\*UP\*CP\*UP\*UP\*U)-3')



#### • Molecule 3: RNA (5'-R(\*CP\*AP\*AP\*UP\*GP\*UP\*GP\*AP\*AP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.81Å 116.99Å 69.77Å 90.00° 92.45° 90.00°	Depositor
Resolution (Å)	34.47 – 1.80 34.47 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.6 (34.47-1.80) 97.6 (34.47-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.21 (at 1.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.167 , 0.197 0.167 , 0.195	Depositor DCC
$R_{free}$ test set	4009 reflections (4.99%)	DCC
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtriage
Anisotropy	0.442	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7491	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG, IPA, IPH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/6570	0.55	0/8889
2	B	0.72	1/441 (0.2%)	1.05	3/679 (0.4%)
3	D	0.47	0/219	0.95	0/340
All	All	0.40	1/7230 (0.0%)	0.61	3/9908 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	U	OP3-P	-10.20	1.49	1.61

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	20	U	C2-N1-C1'	6.11	125.04	117.70
2	B	20	U	C6-N1-C1'	-5.33	113.74	121.20
2	B	20	U	N1-C1'-C2'	5.05	120.57	114.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6419	0	6487	55	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	399	0	203	2	0
3	D	196	0	98	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	D	1	0	0	0	0
5	A	8	0	16	2	0
6	A	28	0	24	1	0
7	B	5	0	0	0	0
8	A	354	0	0	1	0
8	B	45	0	0	0	0
8	D	34	0	0	0	0
All	All	7491	0	6828	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:ALA:H	5:A:902:IPA:H12	1.37	0.90
1:A:243:ILE:HD11	1:A:325:LEU:HD11	1.65	0.77
1:A:484:PRO:HB2	1:A:486:GLN:HE21	1.49	0.77
1:A:231:ILE:HG23	1:A:243:ILE:HD12	1.69	0.73
1:A:112:LYS:HG2	1:A:133:LYS:HG2	1.73	0.70
1:A:696:LYS:HB3	1:A:697:ASP:HB2	1.74	0.69
1:A:74:ILE:HG23	1:A:117:VAL:HG21	1.73	0.68
1:A:169:LEU:HD21	1:A:173:ARG:HH21	1.61	0.64
1:A:55:TYR:HD2	1:A:134:TRP:HZ3	1.48	0.61
1:A:110:ARG:HG2	1:A:134:TRP:CD1	2.37	0.60
1:A:296:LEU:HD11	1:A:306:CYS:HB2	1.84	0.59
1:A:395:ARG:NE	1:A:395:ARG:HA	2.18	0.59
1:A:119:LEU:HD12	1:A:120:PRO:HD2	1.85	0.58
1:A:696:LYS:HB3	1:A:697:ASP:CA	2.35	0.56
1:A:756:ILE:HD11	1:A:795:ARG:NH2	2.21	0.56
1:A:484:PRO:HB2	1:A:486:GLN:NE2	2.20	0.55
1:A:695:GLU:HG2	1:A:696:LYS:HB2	1.88	0.55
1:A:186:GLU:CD	1:A:186:GLU:H	2.11	0.54
1:A:114:GLU:OE2	1:A:129:LYS:HB3	2.08	0.53
1:A:296:LEU:H	1:A:296:LEU:HD12	1.74	0.53
1:A:696:LYS:HB3	1:A:697:ASP:CB	2.38	0.53
1:A:55:TYR:HD2	1:A:134:TRP:CZ3	2.27	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:PRO:HB2	1:A:52:ILE:HD12	1.93	0.50
1:A:279:TYR:HB3	1:A:330:VAL:HB	1.94	0.50
1:A:77:HIS:HA	1:A:80:GLN:HG2	1.93	0.50
1:A:696:LYS:HB3	1:A:697:ASP:C	2.32	0.49
1:A:45:PHE:CZ	1:A:383:MET:HG3	2.47	0.49
1:A:151:LEU:HG	1:A:152:PRO:HD2	1.93	0.49
1:A:77:HIS:CD2	1:A:119:LEU:HD13	2.47	0.49
1:A:199:TRP:HZ2	5:A:902:IPA:H32	1.77	0.48
1:A:277:ARG:HG2	1:A:279:TYR:CE2	2.47	0.48
2:B:3:C:H2'	2:B:4:A:C8	2.49	0.48
1:A:569:VAL:HG21	1:A:791:VAL:HB	1.96	0.48
1:A:106:LEU:O	1:A:108:ILE:HG13	2.14	0.48
1:A:86:ILE:HG22	1:A:87:PHE:CG	2.50	0.47
1:A:55:TYR:CD2	1:A:134:TRP:HZ3	2.32	0.47
1:A:61:ILE:O	1:A:64:GLU:HB3	2.15	0.47
1:A:336:HIS:ND1	2:B:21:U:O2	2.48	0.46
1:A:673:GLU:HG3	8:A:1202:HOH:O	2.13	0.46
1:A:110:ARG:HG2	1:A:134:TRP:HD1	1.79	0.46
1:A:333:GLU:HG3	1:A:334:GLN:HG3	1.98	0.46
1:A:245:GLU:HA	1:A:246:GLN:HA	1.76	0.45
1:A:475:ARG:HG2	1:A:485:ILE:HB	1.98	0.45
1:A:370:ARG:O	1:A:758:GLY:HA2	2.16	0.45
1:A:112:LYS:HG2	1:A:133:LYS:CG	2.45	0.45
1:A:119:LEU:HD12	1:A:120:PRO:CD	2.46	0.45
1:A:108:ILE:HD12	1:A:109:GLY:H	1.83	0.43
1:A:322:TYR:N	1:A:342:GLU:OE2	2.40	0.43
1:A:277:ARG:HD3	1:A:279:TYR:OH	2.20	0.42
1:A:696:LYS:CB	1:A:697:ASP:HB2	2.47	0.42
1:A:84:THR:O	1:A:85:GLN:HG2	2.18	0.42
1:A:695:GLU:HG2	1:A:696:LYS:CB	2.50	0.41
1:A:117:VAL:HG22	1:A:128:PHE:HB2	2.02	0.41
1:A:425:LYS:HB3	1:A:425:LYS:HE2	1.79	0.40
1:A:691:CYS:HB2	6:A:904:IPH:H3	2.02	0.40
1:A:69:ARG:NH1	1:A:73:GLU:HG2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	789/859 (92%)	766 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	708/752 (94%)	699 (99%)	9 (1%)	73	66

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	108	ILE
1	A	111	ASP
1	A	134	TRP
1	A	296	LEU
1	A	370	ARG
1	A	387	ASP
1	A	425	LYS
1	A	804	TYR
1	A	815	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	HIS
1	A	228	GLN
1	A	459	GLN
1	A	486	GLN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	17/21 (80%)	1 (5%)	0
3	D	8/11 (72%)	0	0
All	All	25/32 (78%)	1 (4%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	21	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 3 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	IPA	A	902	-	3,3,3	0.55	0	3,3,3	0.31	0
5	IPA	A	903	-	3,3,3	0.57	0	3,3,3	0.27	0
6	IPH	A	904	-	7,7,7	0.52	0	8,8,8	0.25	0
6	IPH	A	905	-	7,7,7	0.38	0	8,8,8	0.33	0
6	IPH	A	906	-	7,7,7	0.59	0	8,8,8	0.23	0
6	IPH	A	907	-	7,7,7	0.46	0	8,8,8	0.38	0
7	PO4	B	102	-	4,4,4	0.72	0	6,6,6	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	IPA	A	902	-	-	0/0/0/0	0/0/0/0
5	IPA	A	903	-	-	0/0/0/0	0/0/0/0
6	IPH	A	904	-	-	0/0/0/0	0/1/1/1
6	IPH	A	905	-	-	0/0/0/0	0/1/1/1
6	IPH	A	906	-	-	0/0/0/0	0/1/1/1
6	IPH	A	907	-	-	0/0/0/0	0/1/1/1
7	PO4	B	102	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	902	IPA	2	0
6	A	904	IPH	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	801/859 (93%)	0.49	98 (12%) 5 4	13, 31, 76, 96	0
2	B	20/21 (95%)	-0.02	1 (5%) 30 25	15, 32, 91, 118	0
3	D	10/11 (90%)	-0.52	0 100 100	24, 26, 45, 69	0
All	All	831/891 (93%)	0.47	99 (11%) 5 4	13, 31, 76, 118	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	GLN	7.4
1	A	110	ARG	7.4
1	A	120	PRO	6.3
1	A	296	LEU	6.2
1	A	423	ARG	6.1
1	A	134	TRP	5.7
1	A	149	GLY	5.6
1	A	245	GLU	5.6
1	A	119	LEU	5.6
1	A	84	THR	5.2
1	A	65	LYS	5.2
1	A	152	PRO	5.0
1	A	23	PHE	5.0
1	A	108	ILE	4.9
1	A	151	LEU	4.7
1	A	64	GLU	4.6
1	A	85	GLN	4.6
1	A	311	TYR	4.4
1	A	248	LYS	4.3
1	A	133	LYS	4.3
1	A	117	VAL	4.2
1	A	87	PHE	4.1
1	A	241	LYS	4.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	277	ARG	4.1
1	A	88	GLY	3.9
1	A	150	ARG	3.9
1	A	52	ILE	3.8
1	A	109	GLY	3.8
1	A	316	HIS	3.8
1	A	86	ILE	3.8
1	A	424	ASN	3.7
1	A	336	HIS	3.7
1	A	295	PRO	3.7
1	A	156	PHE	3.6
1	A	315	ARG	3.6
1	A	51	LYS	3.6
1	A	279	TYR	3.5
1	A	154	VAL	3.5
1	A	127	ILE	3.5
1	A	318	LEU	3.5
1	A	80	GLN	3.5
1	A	247	GLN	3.3
1	A	114	GLU	3.3
1	A	313	LYS	3.3
1	A	63	PRO	3.3
1	A	142	ALA	3.3
1	A	211	TRP	3.2
1	A	132	ILE	3.2
1	A	355	LYS	3.2
1	A	81	HIS	3.1
1	A	115	LEU	3.1
1	A	243	ILE	3.0
1	A	312	PHE	3.0
1	A	293	THR	3.0
1	A	113	VAL	3.0
1	A	244	GLU	2.9
1	A	707	VAL	2.9
1	A	269	ILE	2.8
2	B	20	U	2.7
1	A	322	TYR	2.7
1	A	294	PHE	2.5
1	A	140	LEU	2.5
1	A	55	TYR	2.5
1	A	307	THR	2.5
1	A	332	GLN	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	249	PRO	2.5
1	A	61	ILE	2.4
1	A	291	HIS	2.4
1	A	60	ASP	2.4
1	A	319	VAL	2.4
1	A	790	TYR	2.3
1	A	338	TYR	2.3
1	A	317	LYS	2.3
1	A	791	VAL	2.3
1	A	324	HIS	2.3
1	A	118	THR	2.3
1	A	104	MET	2.2
1	A	138	VAL	2.2
1	A	803	ALA	2.2
1	A	62	LYS	2.2
1	A	111	ASP	2.2
1	A	314	ASP	2.2
1	A	22	ALA	2.2
1	A	331	GLY	2.2
1	A	808	LEU	2.1
1	A	604	GLY	2.1
1	A	116	GLU	2.1
1	A	188	CYS	2.1
1	A	422	GLY	2.1
1	A	53	ASP	2.1
1	A	460	ARG	2.1
1	A	367	ALA	2.1
1	A	147	LEU	2.1
1	A	323	PRO	2.1
1	A	805	TYR	2.1
1	A	797	VAL	2.1
1	A	486	GLN	2.1
1	A	112	LYS	2.0
1	A	605	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	IPA	A	902	4/4	0.72	0.24	5.88	24,37,38,49	0
5	IPA	A	903	4/4	0.79	0.21	4.09	37,42,43,47	0
6	IPH	A	907	7/7	0.88	0.17	1.88	33,38,38,39	0
6	IPH	A	904	7/7	0.83	0.13	1.62	43,46,49,52	0
6	IPH	A	906	7/7	0.93	0.12	1.10	21,24,27,38	0
6	IPH	A	905	7/7	0.98	0.09	0.12	24,26,28,35	0
4	MG	A	901	1/1	0.99	0.06	-2.32	17,17,17,17	0
7	PO4	B	102	5/5	0.78	0.42	-	90,91,92,93	0
4	MG	D	101	1/1	0.98	0.06	-	23,23,23,23	0
4	MG	B	101	1/1	0.99	0.13	-	33,33,33,33	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.