



wwPDB/EMDataBank EM Map/Model Validation Summary Report ⓘ

Sep 12, 2017 – 08:31 AM EDT

PDB ID : 5W5E
EMDB ID: : EMD-6270
Title : Re-refinement of the pyocin tube structure
Authors : Wang, F.; Zheng, W.; Taylor, N.M.; Guerrero-Ferreira, R.C.; Leiman, P.G.;
Egelman, E.H.
Deposited on : unknown
Resolution : 3.50 Å(reported)

This is a wwPDB/EMDataBank EM Map/Model Validation Summary Report
for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.
























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	125131	1336
Ramachandran outliers	121729	1120
Sidechain outliers	121581	1026

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	168	90% 10% .
1	B	168	90% 9% .
1	C	168	90% 10% .
1	D	168	90% 10% .
1	E	168	90% 9% .
1	F	168	90% 9% .
1	G	168	89% 10% .
1	H	168	89% 10% .
1	I	168	90% 9% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	168	 92% 8% .
1	K	168	 91% 8% .
1	L	168	 90% 10% .
1	M	168	 91% 8% .
1	N	168	 91% 8% .
1	O	168	 90% 10% .
1	P	168	 90% 10% .
1	Q	168	 90% 9% .
1	R	168	 91% 8% .
1	S	168	 92% 7% .
1	T	168	 93% 7% .
1	U	168	 92% 7% .
1	V	168	 93% 7% .
1	W	168	 90% 9% .
1	X	168	 91% 8% .
1	Y	168	 92% 8% .
1	Z	168	 92% 7% .
1	a	168	 99% .
1	b	168	 99% .
1	c	168	 99% .
1	d	168	 99% .

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 37890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FIIR2 protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	B	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	C	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	D	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	E	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	F	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	G	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	H	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	I	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	J	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	K	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	L	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	M	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	N	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	O	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	P	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	Q	167	Total 1263	C 800	N 215	O 243	S 5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
1	R	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	S	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	T	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	U	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	V	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	W	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	X	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	Y	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	Z	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	a	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	b	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	c	167	Total 1263	C 800	N 215	O 243	S 5	0	0
1	d	167	Total 1263	C 800	N 215	O 243	S 5	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FIIR2 protein




- Molecule 1: FIIR2 protein




- Molecule 1: FIIR2 protein




- Molecule 1: FIIR2 protein



- Molecule 1: FIIR2 protein



- Molecule 1: FIIR2 protein





- Molecule 1: FIIR2 protein

Chain G: 89% 10%




- Molecule 1: FIIR2 protein

Chain H: 89% 10%



- Molecule 1: FIIR2 protein

Chain I: 90% 9%



- Molecule 1: FIIR2 protein

Chain J: 92% 8%




- Molecule 1: FIIR2 protein

Chain K: 91% 8%



- Molecule 1: FIIR2 protein

Chain L: 90% 10%



- Molecule 1: FIIR2 protein

Chain M: 91% 8%




- Molecule 1: FIIR2 protein

Chain N:  91% 8%




- Molecule 1: FIIR2 protein

Chain O:  90% 10%




- Molecule 1: FIIR2 protein

Chain P:  90% 10%




- Molecule 1: FIIR2 protein

Chain Q:  90% 9%



- Molecule 1: FIIR2 protein

Chain R:  91% 8%



- Molecule 1: FIIR2 protein

Chain S:  92% 7%



- Molecule 1: FIIR2 protein

Chain T:  93% 7%



- Molecule 1: FIIR2 protein

Chain U:  92% 7%



- Molecule 1: FIIR2 protein

Chain V:  93% 7%



- Molecule 1: FIIR2 protein

Chain W:  90% 9%



- Molecule 1: FIIR2 protein

Chain X:  91% 8%




- Molecule 1: FIIR2 protein

Chain Y:  92% 8%



- Molecule 1: FIIR2 protein

Chain Z:  92% 7%



- Molecule 1: FIIR2 protein

Chain a:  99%



- Molecule 1: FIIR2 protein

Chain b:  99%



- Molecule 1: FIIR2 protein

Chain c:  99% .



- Molecule 1: FIIR2 protein

Chain d:  99% .



4 Experimental information

Property	Value	Source
Reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=18.3°, rise=38.4 Å, axial sym=C6	Depositor
Number of segments used	50000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	Not provided	Depositor
Minimum defocus (nm)	Not provided	Depositor
Maximum defocus (nm)	Not provided	Depositor
Magnification	Not provided	Depositor
Image detector	KODAK SO-163 FILM	Depositor

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >2	RMSZ	# Z >2
1	A	0.50	0/1285	0.69	0/1734
1	B	0.50	0/1285	0.69	0/1734
1	C	0.50	0/1285	0.69	0/1734
1	D	0.50	0/1285	0.69	0/1734
1	E	0.50	0/1285	0.69	0/1734
1	F	0.50	0/1285	0.69	0/1734
1	G	0.50	0/1285	0.69	0/1734
1	H	0.50	0/1285	0.69	0/1734
1	I	0.50	0/1285	0.69	0/1734
1	J	0.50	0/1285	0.69	0/1734
1	K	0.50	0/1285	0.69	0/1734
1	L	0.50	0/1285	0.69	0/1734
1	M	0.50	0/1285	0.69	0/1734
1	N	0.50	0/1285	0.69	0/1734
1	O	0.50	0/1285	0.69	0/1734
1	P	0.50	0/1285	0.69	0/1734
1	Q	0.50	0/1285	0.69	0/1734
1	R	0.50	0/1285	0.69	0/1734
1	S	0.50	0/1285	0.69	0/1734
1	T	0.50	0/1285	0.69	0/1734
1	U	0.50	0/1285	0.69	0/1734
1	V	0.50	0/1285	0.69	0/1734
1	W	0.50	0/1285	0.69	0/1734
1	X	0.50	0/1285	0.69	0/1734
1	Y	0.50	0/1285	0.69	0/1734
1	Z	0.50	0/1285	0.69	0/1734
1	a	0.50	0/1285	0.69	0/1734
1	b	0.50	0/1285	0.69	0/1734
1	c	0.50	0/1285	0.69	0/1734
1	d	0.50	0/1285	0.69	0/1734
All	All	0.50	0/38550	0.69	0/52020

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1263	0	1255	17	0
1	B	1263	0	1255	14	0
1	C	1263	0	1255	15	0
1	D	1263	0	1255	16	0
1	E	1263	0	1255	12	0
1	F	1263	0	1255	13	0
1	G	1263	0	1255	14	0
1	H	1263	0	1255	13	0
1	I	1263	0	1255	11	0
1	J	1263	0	1255	11	0
1	K	1263	0	1255	11	0
1	L	1263	0	1255	14	0
1	M	1263	0	1255	12	0
1	N	1263	0	1255	12	0
1	O	1263	0	1255	12	0
1	P	1263	0	1255	13	0
1	Q	1263	0	1255	12	0
1	R	1263	0	1255	11	0
1	S	1263	0	1255	9	0
1	T	1263	0	1255	9	0
1	U	1263	0	1255	9	0
1	V	1263	0	1255	8	0
1	W	1263	0	1255	12	0
1	X	1263	0	1255	11	0
1	Y	1263	0	1255	10	0
1	Z	1263	0	1255	9	0
1	a	1263	0	1255	0	0
1	b	1263	0	1255	0	0
1	c	1263	0	1255	0	0
1	d	1263	0	1255	0	0
All	All	37890	0	37650	201	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 201 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:96:SER:OG	1:I:165:ASP:OD2	2.03	0.77
1:D:96:SER:OG	1:R:165:ASP:OD2	92.37	0.77
1:B:96:SER:OG	1:P:165:ASP:OD2	92.37	0.77
1:J:96:SER:OG	1:V:165:ASP:OD2	2.03	0.77
1:N:165:ASP:OD2	1:Z:96:SER:OG	2.03	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	B	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	C	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	D	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	E	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	F	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	G	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	H	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	I	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	J	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	K	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	L	165/168 (98%)	155 (94%)	10 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	M	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	N	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	O	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	P	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	Q	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	R	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	S	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	T	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	U	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	V	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	W	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	X	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	Y	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	Z	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	a	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	b	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	c	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
1	d	165/168 (98%)	155 (94%)	10 (6%)	0	100	100
All	All	4950/5040 (98%)	4650 (94%)	300 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/131 (99%)	130 (100%)	0	100	100
1	B	130/131 (99%)	130 (100%)	0	100	100
1	C	130/131 (99%)	130 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	130/131 (99%)	130 (100%)	0	100	100
1	E	130/131 (99%)	130 (100%)	0	100	100
1	F	130/131 (99%)	130 (100%)	0	100	100
1	G	130/131 (99%)	130 (100%)	0	100	100
1	H	130/131 (99%)	130 (100%)	0	100	100
1	I	130/131 (99%)	130 (100%)	0	100	100
1	J	130/131 (99%)	130 (100%)	0	100	100
1	K	130/131 (99%)	130 (100%)	0	100	100
1	L	130/131 (99%)	130 (100%)	0	100	100
1	M	130/131 (99%)	130 (100%)	0	100	100
1	N	130/131 (99%)	130 (100%)	0	100	100
1	O	130/131 (99%)	130 (100%)	0	100	100
1	P	130/131 (99%)	130 (100%)	0	100	100
1	Q	130/131 (99%)	130 (100%)	0	100	100
1	R	130/131 (99%)	130 (100%)	0	100	100
1	S	130/131 (99%)	130 (100%)	0	100	100
1	T	130/131 (99%)	130 (100%)	0	100	100
1	U	130/131 (99%)	130 (100%)	0	100	100
1	V	130/131 (99%)	130 (100%)	0	100	100
1	W	130/131 (99%)	130 (100%)	0	100	100
1	X	130/131 (99%)	130 (100%)	0	100	100
1	Y	130/131 (99%)	130 (100%)	0	100	100
1	Z	130/131 (99%)	130 (100%)	0	100	100
1	a	130/131 (99%)	130 (100%)	0	100	100
1	b	130/131 (99%)	130 (100%)	0	100	100
1	c	130/131 (99%)	130 (100%)	0	100	100
1	d	130/131 (99%)	130 (100%)	0	100	100
All	All	3900/3930 (99%)	3900 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.