



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 3, 2017 – 07:41 PM EDT

PDB ID : 2W6C
Title : ACHE IN COMPLEX WITH A BIS-(-)-NOR-MEPTAZINOL DERIVATIVE
Authors : Paz, A.; Xie, Q.; Greenblatt, H.M.; Fu, W.; Tang, Y.; Silman, I.; Qiu, Z.;
Sussman, J.L.
Deposited on : unknown
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	586	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; right: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="position: absolute; bottom: 0; left: 0; width: 68%;"></div> <div style="position: absolute; bottom: 0; left: 68%; width: 17%;"></div> <div style="position: absolute; bottom: 0; left: 85%; width: 10%;"></div> </div> <div>68% 17% . . 10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:


Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	X	602	X	-	-	-

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	528	Total	C	N	O	S	0	0	0
			4171	2689	700	760	22			

- Molecule 2 is 3-[(3R)-3-ethyl-1-{9-[(3S)-3-ethyl-3-(3-hydroxyphenyl)azepan-1-yl]nonyl}azepan-3-yl]phenol (three-letter code: BM4) (formula: $\text{C}_{37}\text{H}_{58}\text{N}_2\text{O}_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	X	1	Total	C	N	O	0	0
			26	23	2	1		

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	X	1	Total	C	N	O	0	0
			14	8	1	5		
3	X	1	Total	C	N	O	0	0
			14	8	1	5		
3	X	1	Total	C	N	O	0	0
			14	8	1	5		
3	X	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	X	115	Total	O	0	0
			115	115		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACETYLCHOLINESTERASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	112.17Å 112.17Å 137.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.68 – 2.69 39.68 – 2.69	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.68-2.69) 99.6 (39.68-2.69)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.44 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.4.0067	Depositor
R, R_{free}	0.185 , 0.235 0.189 , 0.237	Depositor DCC
R_{free} test set	1452 reflections (5.44%)	DCC
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4368	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BM4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.96	3/4292 (0.1%)	0.98	17/5829 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	260	GLU	CB-CG	6.79	1.65	1.52
1	X	268	GLU	CB-CG	5.05	1.61	1.52
1	X	260	GLU	CG-CD	5.01	1.59	1.51

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	494	LEU	CA-CB-CG	7.09	131.61	115.30
1	X	243	ARG	NE-CZ-NH2	-7.02	116.79	120.30
1	X	531	LEU	CA-CB-CG	6.94	131.27	115.30
1	X	143	LEU	CA-CB-CG	-6.62	100.08	115.30
1	X	430	LEU	CA-CB-CG	6.54	130.35	115.30
1	X	295	VAL	CB-CA-C	-6.34	99.35	111.40
1	X	518	VAL	CB-CA-C	-6.24	99.55	111.40
1	X	244	ARG	NE-CZ-NH2	-6.21	117.19	120.30
1	X	244	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	X	430	LEU	CB-CG-CD2	5.60	120.53	111.00
1	X	358	LEU	CB-CG-CD1	5.58	120.49	111.00
1	X	243	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	X	467	ARG	NE-CZ-NH1	-5.38	117.61	120.30
1	X	175	MET	CG-SD-CE	-5.26	91.78	100.20
1	X	267	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	X	468	ARG	NE-CZ-NH1	-5.16	117.72	120.30
1	X	287	ILE	CB-CA-C	5.02	121.65	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	4171	0	4007	105	0
2	X	26	0	38	1	0
3	X	56	0	51	3	0
4	X	115	0	0	17	0
All	All	4368	0	4096	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:526:GLN:HG3	4:X:774:HOH:O	1.39	1.19
1:X:497:THR:HB	4:X:702:HOH:O	0.98	1.15
1:X:74:GLN:HE21	1:X:74:GLN:HA	1.34	0.91
1:X:287:ILE:HD11	1:X:335:GLY:HA3	1.55	0.89
1:X:22:VAL:HG13	1:X:133:LYS:HG3	1.56	0.87
1:X:383:ASN:C	1:X:383:ASN:HD22	1.79	0.86
1:X:16:MET:SD	4:X:811:HOH:O	2.37	0.81
1:X:439:ILE:HG22	1:X:440:HIS:O	1.83	0.79
1:X:504:ASP:HB2	4:X:761:HOH:O	1.84	0.76
1:X:74:GLN:HE21	1:X:74:GLN:CA	2.01	0.72
1:X:23:LEU:HD11	1:X:452:LEU:HD12	1.71	0.72
1:X:484:GLU:OE2	1:X:484:GLU:HA	1.91	0.69
1:X:49:GLU:HG2	4:X:813:HOH:O	1.92	0.69
1:X:450:LEU:N	1:X:450:LEU:CD2	2.57	0.68
1:X:5:GLU:OE2	1:X:104:PRO:HA	1.94	0.67
1:X:310:ASN:OD1	1:X:410:LYS:NZ	2.21	0.67
1:X:456:LEU:O	1:X:457:ASN:HB2	1.96	0.66
1:X:421:TYR:HB2	1:X:505:LEU:HD22	1.78	0.64
1:X:450:LEU:H	1:X:450:LEU:HD23	1.62	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:74:GLN:NE2	1:X:74:GLN:HA	2.11	0.63
1:X:497:THR:CB	4:X:702:HOH:O	1.79	0.62
1:X:484:GLU:OE2	1:X:485:PRO:HD2	2.01	0.61
1:X:321:LEU:HD23	1:X:321:LEU:N	2.18	0.58
1:X:287:ILE:HD11	1:X:335:GLY:CA	2.33	0.56
1:X:383:ASN:ND2	1:X:383:ASN:C	2.54	0.56
1:X:16:MET:CG	4:X:811:HOH:O	2.54	0.56
1:X:531:LEU:C	1:X:531:LEU:HD12	2.26	0.55
1:X:453:VAL:CG2	1:X:456:LEU:HG	2.36	0.55
1:X:330:PHE:HE2	1:X:439:ILE:HG21	1.72	0.55
1:X:325:LYS:HB2	1:X:422:PHE:CZ	2.43	0.54
1:X:43:MET:O	1:X:46:ARG:HB2	2.08	0.54
1:X:190:ASP:OD1	1:X:192:LYS:HG2	2.07	0.54
3:X:605:NAG:H62	4:X:806:HOH:O	2.07	0.54
1:X:37:GLU:OE2	1:X:52:LYS:HG3	2.09	0.53
1:X:19:ARG:NH2	1:X:26:HIS:HB2	2.24	0.53
1:X:299:GLU:HB2	4:X:803:HOH:O	2.08	0.53
1:X:528:LEU:HB3	1:X:529:PRO:HD3	1.90	0.53
1:X:207:GLY:HA3	1:X:229:PRO:HD3	1.91	0.53
1:X:74:GLN:NE2	1:X:74:GLN:CA	2.70	0.52
1:X:16:MET:HG3	4:X:811:HOH:O	2.09	0.52
1:X:452:LEU:HD13	1:X:467:ARG:NH2	2.25	0.52
1:X:451:PRO:HA	1:X:458:TYR:CD1	2.45	0.51
3:X:602:NAG:H3	3:X:602:NAG:H83	1.93	0.51
1:X:450:LEU:H	1:X:450:LEU:CD2	2.22	0.50
1:X:405:MET:HA	1:X:408:VAL:HG12	1.92	0.50
1:X:468:ARG:HH21	1:X:510:MET:HE3	1.76	0.50
1:X:419:TYR:CZ	1:X:494:LEU:HD13	2.46	0.50
1:X:484:GLU:OE2	1:X:485:PRO:CD	2.58	0.50
1:X:497:THR:C	4:X:702:HOH:O	2.49	0.49
1:X:450:LEU:HD22	1:X:450:LEU:N	2.25	0.49
1:X:405:MET:HA	1:X:408:VAL:CG1	2.43	0.49
1:X:68:GLN:HE21	1:X:68:GLN:HA	1.77	0.48
1:X:269:LYS:HB2	1:X:274:LEU:HD11	1.95	0.48
1:X:320:LEU:C	1:X:320:LEU:HD23	2.34	0.48
1:X:162:GLN:HG3	4:X:815:HOH:O	2.14	0.48
1:X:25:SER:OG	1:X:26:HIS:N	2.45	0.47
1:X:515:ARG:O	1:X:518:VAL:HG22	2.13	0.47
1:X:197:PHE:CB	1:X:223:ILE:HB	2.44	0.47
1:X:260:GLU:H	1:X:260:GLU:CD	2.18	0.47
1:X:121:TYR:CD1	1:X:122:SER:HB3	2.50	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:321:LEU:CD2	1:X:321:LEU:N	2.78	0.47
1:X:468:ARG:HD3	4:X:701:HOH:O	2.14	0.47
1:X:35:PHE:O	1:X:175:MET:CE	2.64	0.46
1:X:282:LEU:HD23	1:X:282:LEU:HA	1.77	0.46
1:X:206:VAL:CG1	1:X:222:ALA:HB1	2.46	0.46
1:X:404:LEU:O	1:X:408:VAL:HG12	2.16	0.45
1:X:383:ASN:HD22	1:X:384:GLY:N	2.15	0.45
1:X:357:LYS:HD2	4:X:797:HOH:O	2.16	0.45
1:X:502:PHE:CZ	1:X:513:HIS:HB2	2.51	0.45
1:X:82:GLU:HG3	4:X:798:HOH:O	2.18	0.44
1:X:452:LEU:N	1:X:452:LEU:HD23	2.32	0.44
1:X:379:MET:O	1:X:380:ASP:HB2	2.18	0.44
1:X:50:PRO:HA	1:X:175:MET:HE3	1.99	0.44
1:X:35:PHE:O	1:X:175:MET:HE3	2.16	0.43
1:X:468:ARG:HH21	1:X:510:MET:CE	2.31	0.43
1:X:15:VAL:CG1	1:X:30:PHE:HD2	2.32	0.43
1:X:404:LEU:HD23	1:X:404:LEU:C	2.38	0.43
1:X:197:PHE:HB3	1:X:223:ILE:HB	2.01	0.43
1:X:303:THR:HB	4:X:734:HOH:O	2.17	0.42
1:X:325:LYS:NZ	1:X:326:ASP:OD1	2.52	0.42
1:X:454:LYS:HA	1:X:454:LYS:HD3	1.73	0.42
1:X:204:ALA:O	1:X:208:MET:HG3	2.19	0.42
1:X:113:VAL:HG22	1:X:144:VAL:HB	2.01	0.42
1:X:456:LEU:O	1:X:457:ASN:CB	2.67	0.42
1:X:78:PHE:O	1:X:82:GLU:HB2	2.19	0.42
1:X:223:ILE:HA	1:X:320:LEU:O	2.19	0.42
1:X:21:PRO:HA	1:X:25:SER:O	2.19	0.42
1:X:302:PRO:HD2	1:X:308:MET:SD	2.59	0.42
1:X:27:ILE:HG21	1:X:27:ILE:HD13	1.79	0.42
1:X:366:LEU:HD23	1:X:531:LEU:CD1	2.50	0.42
2:X:601:BM4:HAF	2:X:601:BM4:HABA	1.84	0.42
1:X:240:GLU:OE1	1:X:243:ARG:HD3	2.20	0.41
1:X:115:ILE:HG23	1:X:146:LEU:HD11	2.01	0.41
1:X:39:PRO:HG3	1:X:95:LEU:HD11	2.03	0.41
1:X:252:LEU:HB3	1:X:269:LYS:NZ	2.35	0.41
1:X:426:ARG:CZ	1:X:430:LEU:HD12	2.50	0.41
1:X:330:PHE:CE2	1:X:439:ILE:HG21	2.54	0.41
1:X:139:GLU:O	1:X:140:GLU:CB	2.68	0.41
1:X:531:LEU:C	1:X:531:LEU:CD1	2.89	0.41
1:X:240:GLU:O	1:X:244:ARG:HG3	2.20	0.41
1:X:383:ASN:ND2	1:X:386:LYS:H	2.18	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:299:GLU:HB3	1:X:300:PHE:H	1.78	0.41
1:X:450:LEU:N	1:X:450:LEU:HD23	2.23	0.41
1:X:274:LEU:HD12	1:X:274:LEU:N	2.36	0.40
3:X:602:NAG:H3	3:X:602:NAG:C8	2.51	0.40
1:X:221:ARG:HD3	1:X:480:GLY:HA2	2.04	0.40
1:X:46:ARG:CD	4:X:731:HOH:O	2.68	0.40
1:X:132:GLY:HA3	1:X:143:LEU:HD22	2.03	0.40
1:X:515:ARG:HB3	1:X:518:VAL:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	524/586 (89%)	492 (94%)	30 (6%)	2 (0%)	38 66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	380	ASP
1	X	457	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	X	446/513 (87%)	413 (93%)	33 (7%)	16	37

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	5	GLU
1	X	28	SER
1	X	49	GLU
1	X	68	GLN
1	X	74	GLN
1	X	89	GLU
1	X	91	SER
1	X	129	VAL
1	X	140	GLU
1	X	143	LEU
1	X	156	LEU
1	X	194	VAL
1	X	243	ARG
1	X	287	ILE
1	X	288	PHE
1	X	295	VAL
1	X	325	LYS
1	X	354	SER
1	X	356	VAL
1	X	358	LEU
1	X	383	ASN
1	X	391	LEU
1	X	430	LEU
1	X	450	LEU
1	X	452	LEU
1	X	453	VAL
1	X	473	TRP
1	X	484	GLU
1	X	497	THR
1	X	499	GLU
1	X	505	LEU
1	X	518	VAL
1	X	531	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	68	GLN
1	X	74	GLN
1	X	383	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BM4	X	601	-	25,27,44	0.73	0	23,34,59	1.35	3 (13%)
3	NAG	X	602	1	14,14,15	0.98	1 (7%)	15,19,21	1.86	4 (26%)
3	NAG	X	603	1,3	14,14,15	0.76	1 (7%)	15,19,21	1.68	4 (26%)
3	NAG	X	604	3	14,14,15	0.62	0	15,19,21	1.97	5 (33%)
3	NAG	X	605	1	14,14,15	0.75	0	15,19,21	3.62	8 (53%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BM4	X	601	-	-	0/19/32/56	0/2/2/4
3	NAG	X	602	1	1/1/5/7	1/6/23/26	0/1/1/1
3	NAG	X	603	1,3	-	0/6/23/26	0/1/1/1
3	NAG	X	604	3	-	0/6/23/26	0/1/1/1
3	NAG	X	605	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	603	NAG	O5-C1	-2.32	1.39	1.43
3	X	602	NAG	C1-C2	2.42	1.55	1.52

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	605	NAG	C2-N2-C7	-8.35	110.75	122.94
3	X	605	NAG	C3-C4-C5	-7.88	96.33	110.22
2	X	601	BM4	CAN-CAM-CAC	-4.67	110.81	115.89
3	X	604	NAG	O5-C1-C2	-3.51	106.59	111.47
3	X	603	NAG	O5-C1-C2	-3.50	106.61	111.47
3	X	604	NAG	C2-N2-C7	-3.29	118.14	122.94
3	X	603	NAG	C3-C4-C5	-3.08	104.78	110.22
2	X	601	BM4	CAF-CAG-NAA	-3.04	109.68	115.42
3	X	605	NAG	O7-C7-N2	-2.85	116.43	121.92
3	X	602	NAG	O7-C7-C8	-2.69	117.16	122.06
3	X	603	NAG	C2-N2-C7	-2.63	119.11	122.94
3	X	605	NAG	C4-C3-C2	-2.45	107.42	111.02
2	X	601	BM4	CAI-CAH-NAA	-2.25	108.25	113.90
3	X	604	NAG	O6-C6-C5	-2.16	104.07	111.34
3	X	605	NAG	C8-C7-N2	2.18	120.05	116.11
3	X	603	NAG	O4-C4-C5	2.35	115.20	109.28
3	X	604	NAG	O4-C4-C5	2.64	115.93	109.28
3	X	605	NAG	O3-C3-C2	2.67	115.11	109.39
3	X	602	NAG	C1-C2-N2	2.75	115.19	110.49
3	X	605	NAG	O4-C4-C5	2.88	116.54	109.28
3	X	602	NAG	C2-N2-C7	2.95	127.25	122.94
3	X	604	NAG	C4-C3-C2	3.63	116.34	111.02
3	X	602	NAG	C4-C3-C2	3.97	116.84	111.02
3	X	605	NAG	C1-O5-C5	5.16	119.27	112.17

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	X	602	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	X	602	NAG	O7-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	X	601	BM4	1	0
3	X	602	NAG	2	0
3	X	605	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	528/586 (90%)	-0.53	5 (0%) 84 85	19, 32, 49, 62	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	55	SER	3.4
1	X	535	THR	2.3
1	X	56	GLY	2.2
1	X	509	PRO	2.1
1	X	264	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	BM4	X	601	26/41	0.97	0.22	1.14	17,26,41,41	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	X	603	14/15	0.96	0.12	-	22,33,38,41	14
3	NAG	X	605	14/15	0.91	0.38	-	55,57,60,60	14
3	NAG	X	602	14/15	0.85	0.16	-	52,55,57,57	14
3	NAG	X	604	14/15	0.86	0.33	-	41,44,49,50	14

6.5 Other polymers [i](#)

There are no such residues in this entry.