



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 17, 2017 – 08:01 AM EDT

PDB ID : 4W6P
Title : Crystal Structure of Full-Length Split GFP Mutant D102C Disulfide Dimer,
P 21 21 21 Space Group
Authors : Leibly, D.J.; Waldo, G.S.; Yeates, T.O.
Deposited on : unknown
Resolution : 3.08 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1116 (3.10-3.06)
Clashscore	112137	1220 (3.10-3.06)
Ramachandran outliers	110173	1176 (3.10-3.06)
Sidechain outliers	110143	1176 (3.10-3.06)
RSRZ outliers	101464	1123 (3.10-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	<div> <div></div> <div>76% 17% 7%</div> </div>
1	B	234	<div> <div></div> <div>70% 21% 8%</div> </div>
1	C	234	<div> <div>4%</div> <div>63% 21% 15%</div> </div>
1	D	234	<div> <div></div> <div>64% 21% 13%</div> </div>
1	E	234	<div> <div></div> <div>67% 22% 10%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	234	
1	G	234	
1	H	234	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	CRO	E	66	-	-	X	-
1	CRO	F	66	-	-	X	-

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 12960 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called fluorescent protein D102C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	218	Total	C	N	O	S	0	0	0
			1745	1112	292	338	3			
1	B	215	Total	C	N	O	S	0	0	0
			1735	1104	295	332	4			
1	C	199	Total	C	N	O	S	0	0	0
			1599	1024	264	308	3			
1	D	203	Total	C	N	O	S	0	0	0
			1621	1040	272	306	3			
1	E	210	Total	C	N	O	S	0	0	0
			1696	1085	284	323	4			
1	F	215	Total	C	N	O	S	0	0	0
			1718	1097	287	331	3			
1	G	198	Total	C	N	O	S	0	0	0
			1584	1014	264	303	3			
1	H	160	Total	C	N	O	S	0	0	0
			1262	808	208	244	2			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: fluorescent protein D102C



- Molecule 1: fluorescent protein D102C




- Molecule 1: fluorescent protein D102C



- Molecule 1: fluorescent protein D102C



WORLDWIDE
 **PDB**
PROTEIN DATA BANK



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.05Å 117.86Å 209.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	79.58 – 3.08 79.58 – 3.09	Depositor EDS
% Data completeness (in resolution range)	98.0 (79.58-3.08) 98.0 (79.58-3.09)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.95 (at 3.07Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.232 , 0.279 0.232 , 0.279	Depositor DCC
R_{free} test set	3913 reflections (10.00%)	DCC
Wilson B-factor (Å ²)	70.7	Xtrriage
Anisotropy	0.526	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	12960	wwPDB-VP
Average B, all atoms (Å ²)	80.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	1/1758 (0.1%)	0.64	2/2375 (0.1%)
1	B	0.34	0/1747	0.59	1/2356 (0.0%)
1	C	0.38	1/1606 (0.1%)	0.60	2/2167 (0.1%)
1	D	0.37	1/1629 (0.1%)	0.77	4/2196 (0.2%)
1	E	0.37	1/1709 (0.1%)	0.60	2/2307 (0.1%)
1	F	0.35	1/1731 (0.1%)	0.63	1/2339 (0.0%)
1	G	0.42	2/1614 (0.1%)	0.74	4/2177 (0.2%)
1	H	0.32	0/1281	0.60	0/1727
All	All	0.37	7/13075 (0.1%)	0.65	16/17644 (0.1%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	68	VAL	N-CA	-7.52	1.31	1.46
1	G	68	VAL	N-CA	-6.95	1.32	1.46
1	A	68	VAL	N-CA	-6.64	1.33	1.46
1	G	102	CYS	CB-SG	-6.63	1.71	1.82
1	E	68	VAL	N-CA	-6.08	1.34	1.46

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	64	LEU	CA-C-O	-18.78	80.65	120.10
1	G	68	VAL	N-CA-CB	-13.54	81.72	111.50
1	A	68	VAL	N-CA-CB	-10.92	87.47	111.50
1	G	68	VAL	N-CA-C	-9.26	86.00	111.00
1	G	102	CYS	CA-CB-SG	8.75	129.75	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1687	29	0
1	B	1735	0	1687	35	0
1	C	1599	0	1552	38	0
1	D	1621	0	1589	49	0
1	E	1696	0	1650	52	0
1	F	1718	0	1667	49	0
1	G	1584	0	1555	41	0
1	H	1262	0	1234	39	0
All	All	12960	0	12621	323	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 323 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:121:ASN:ND2	1:E:123:ILE:HD11	1.62	1.15
1:G:46:PHE:CE2	1:G:64:LEU:HD22	1.89	1.08
1:G:98:ILE:HB	1:G:106:TYR:HB2	1.38	1.02
1:E:66:CRO:HG11	1:E:222:GLU:OE2	1.59	1.00
1:D:60:LEU:O	1:D:63:THR:O	1.81	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/234 (89%)	203 (97%)	6 (3%)	0	100	100
1	B	206/234 (88%)	201 (98%)	5 (2%)	0	100	100
1	C	186/234 (80%)	178 (96%)	8 (4%)	0	100	100
1	D	190/234 (81%)	185 (97%)	5 (3%)	0	100	100
1	E	201/234 (86%)	194 (96%)	7 (4%)	0	100	100
1	F	206/234 (88%)	200 (97%)	6 (3%)	0	100	100
1	G	188/234 (80%)	179 (95%)	8 (4%)	1 (0%)	32	70
1	H	146/234 (62%)	139 (95%)	6 (4%)	1 (1%)	25	63
All	All	1532/1872 (82%)	1479 (96%)	51 (3%)	2 (0%)	55	87

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	98	ILE
1	H	199	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/203 (92%)	186 (100%)	1 (0%)	91	96
1	B	187/203 (92%)	187 (100%)	0	100	100
1	C	172/203 (85%)	171 (99%)	1 (1%)	89	95
1	D	173/203 (85%)	173 (100%)	0	100	100
1	E	182/203 (90%)	181 (100%)	1 (0%)	91	96
1	F	184/203 (91%)	184 (100%)	0	100	100
1	G	173/203 (85%)	172 (99%)	1 (1%)	89	95
1	H	139/203 (68%)	137 (99%)	2 (1%)	71	89
All	All	1397/1624 (86%)	1391 (100%)	6 (0%)	93	96

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	68	VAL
1	H	198	ASP
1	G	68	VAL
1	C	155	ASP
1	H	62	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	185	ASN
1	C	94	GLN
1	C	199	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CRO	A	66	1	23,23,24	3.62	7 (30%)	27,32,34	5.10	8 (29%)
1	CRO	B	66	1	23,23,24	3.60	8 (34%)	27,32,34	4.82	12 (44%)
1	CRO	C	66	1	23,23,24	3.56	7 (30%)	27,32,34	4.87	7 (25%)
1	CRO	D	66	1	23,23,24	3.47	4 (17%)	27,32,34	4.00	7 (25%)
1	CRO	E	66	1	23,23,24	3.53	5 (21%)	27,32,34	4.45	7 (25%)
1	CRO	F	66	1	23,23,24	3.62	5 (21%)	27,32,34	4.51	8 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	A	66	1	-	0/12/31/32	0/2/2/2
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2
1	CRO	C	66	1	-	0/12/31/32	0/2/2/2
1	CRO	D	66	1	-	0/12/31/32	0/2/2/2
1	CRO	E	66	1	-	0/12/31/32	0/2/2/2
1	CRO	F	66	1	-	0/12/31/32	0/2/2/2

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66	CRO	CA2-C2	-5.91	1.42	1.48
1	C	66	CRO	CA2-C2	-5.19	1.43	1.48
1	F	66	CRO	CA1-N1	-5.10	1.31	1.47
1	B	66	CRO	CA2-C2	-4.83	1.43	1.48
1	C	66	CRO	C2-N3	-4.50	1.28	1.39

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	CRO	O2-C2-CA2	-15.96	122.36	130.97
1	A	66	CRO	O2-C2-CA2	-15.19	122.78	130.97
1	C	66	CRO	O2-C2-CA2	-11.12	124.97	130.97
1	C	66	CRO	O3-C3-CA3	-10.85	90.53	126.38
1	E	66	CRO	O3-C3-CA3	-10.66	91.15	126.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 49 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	66	CRO	6	0
1	B	66	CRO	7	0
1	C	66	CRO	7	0
1	D	66	CRO	6	0
1	E	66	CRO	12	0
1	F	66	CRO	11	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/234 (92%)	0.22	1 (0%) 90 80	38, 61, 98, 142	0
1	B	214/234 (91%)	0.33	3 (1%) 75 56	38, 66, 127, 180	0
1	C	198/234 (84%)	0.43	10 (5%) 29 13	31, 68, 118, 152	0
1	D	202/234 (86%)	0.33	3 (1%) 74 54	35, 69, 113, 147	0
1	E	209/234 (89%)	0.16	0 100 100	47, 80, 122, 157	0
1	F	214/234 (91%)	0.31	2 (0%) 84 68	51, 82, 129, 186	0
1	G	198/234 (84%)	0.17	0 100 100	49, 82, 118, 136	0
1	H	160/234 (68%)	0.64	11 (6%) 18 7	60, 107, 152, 206	0
All	All	1612/1872 (86%)	0.31	30 (1%) 67 46	31, 76, 128, 206	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	118	THR	4.2
1	H	26	LYS	3.8
1	C	86	SER	3.8
1	C	37	ALA	3.4
1	D	161	ILE	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CRO	D	66	22/23	0.89	0.42	-	65,67,73,78	0
1	CRO	F	66	22/23	0.83	0.37	-	68,72,75,76	0
1	CRO	B	66	22/23	0.94	0.31	-	62,67,71,72	0
1	CRO	E	66	22/23	0.93	0.24	-	75,80,84,85	0
1	CRO	C	66	22/23	0.85	0.34	-	72,76,83,88	0
1	CRO	A	66	22/23	0.89	0.27	-	53,55,58,60	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.