



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 01:22 pm GMT

PDB ID : 3W9G  
Title : Crystal structure of the ankilin repeat domain of chicken TRPV4  
Authors : Itoh, Y.; Hamada-nakahara, S.; Suetsugu, S.  
Deposited on : 2013-04-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	<div> <div>3%</div> <div>89%</div> <div>9%</div> <div>•</div> </div>
1	B	260	<div> <div>5%</div> <div>93%</div> <div>5%</div> <div>•</div> </div>
1	C	260	<div> <div>3%</div> <div>88%</div> <div>8%</div> <div>•</div> </div>
1	D	260	<div> <div>27%</div> <div>80%</div> <div>17%</div> <div>•</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vanilloid receptor-related osmotically activated channel protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	6	0
			2065	1307	382	367	9			
1	B	256	Total	C	N	O	S	0	5	0
			2058	1302	379	368	9			
1	C	252	Total	C	N	O	S	0	3	0
			2009	1273	366	361	9			
1	D	252	Total	C	N	O	S	0	0	0
			2001	1267	366	360	8			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	MET	-	EXPRESSION TAG	UNP Q9DFS3
A	383	ALA	-	EXPRESSION TAG	UNP Q9DFS3
A	384	ALA	-	EXPRESSION TAG	UNP Q9DFS3
A	385	ALA	-	EXPRESSION TAG	UNP Q9DFS3
A	386	HIS	-	EXPRESSION TAG	UNP Q9DFS3
A	387	HIS	-	EXPRESSION TAG	UNP Q9DFS3
A	388	HIS	-	EXPRESSION TAG	UNP Q9DFS3
A	389	HIS	-	EXPRESSION TAG	UNP Q9DFS3
A	390	HIS	-	EXPRESSION TAG	UNP Q9DFS3
A	391	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	132	MET	-	EXPRESSION TAG	UNP Q9DFS3
B	383	ALA	-	EXPRESSION TAG	UNP Q9DFS3
B	384	ALA	-	EXPRESSION TAG	UNP Q9DFS3
B	385	ALA	-	EXPRESSION TAG	UNP Q9DFS3
B	386	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	387	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	388	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	389	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	390	HIS	-	EXPRESSION TAG	UNP Q9DFS3
B	391	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	132	MET	-	EXPRESSION TAG	UNP Q9DFS3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	383	ALA	-	EXPRESSION TAG	UNP Q9DFS3
C	384	ALA	-	EXPRESSION TAG	UNP Q9DFS3
C	385	ALA	-	EXPRESSION TAG	UNP Q9DFS3
C	386	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	387	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	388	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	389	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	390	HIS	-	EXPRESSION TAG	UNP Q9DFS3
C	391	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	132	MET	-	EXPRESSION TAG	UNP Q9DFS3
D	383	ALA	-	EXPRESSION TAG	UNP Q9DFS3
D	384	ALA	-	EXPRESSION TAG	UNP Q9DFS3
D	385	ALA	-	EXPRESSION TAG	UNP Q9DFS3
D	386	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	387	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	388	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	389	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	390	HIS	-	EXPRESSION TAG	UNP Q9DFS3
D	391	HIS	-	EXPRESSION TAG	UNP Q9DFS3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	205	Total O 205 205	0	0
2	B	232	Total O 232 232	0	0
2	C	132	Total O 132 132	0	0
2	D	39	Total O 39 39	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vanilloid receptor-related osmotically activated channel protein




- Molecule 1: Vanilloid receptor-related osmotically activated channel protein



- Molecule 1: Vanilloid receptor-related osmotically activated channel protein



- Molecule 1: Vanilloid receptor-related osmotically activated channel protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.95Å 48.27Å 134.03Å 90.00° 101.79° 90.00°	Depositor
Resolution (Å)	37.57 – 2.00 37.57 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.57-2.00) 99.7 (37.57-2.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.51 (at 2.00Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.179 , 0.211 0.179 , 0.212	Depositor DCC
$R_{free}$ test set	1816 reflections (2.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtrriage
Anisotropy	0.444	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 57.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8741	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.88 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.0735e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/2138	0.58	1/2882 (0.0%)
1	B	0.47	0/2125	0.59	0/2866
1	C	0.35	0/2065	0.52	0/2786
1	D	0.31	0/2041	0.53	0/2754
All	All	0.40	0/8369	0.56	1/11288 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	ARG	NE-CZ-NH2	-5.01	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	2095	11	0
1	B	2058	0	2086	8	0
1	C	2009	0	2023	14	0
1	D	2001	0	2016	23	0
2	A	205	0	0	1	0
2	B	232	0	0	1	0
2	C	132	0	0	0	0
2	D	39	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8741	0	8220	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:LYS:NZ	1:A:241:GLU:OE2	2.26	0.69
1:D:164:LYS:NZ	1:D:170:GLU:OE2	2.25	0.67
1:B:237:LYS:NZ	1:B:241:GLU:OE2	2.30	0.64
1:A:300:LEU:HD11	1:A:338:LYS:HG2	1.81	0.61
1:C:237:LYS:NZ	1:C:241:GLU:OE2	2.34	0.60
1:D:275:LEU:HD21	1:D:300:LEU:HD23	1.84	0.59
1:D:213:ILE:HG23	1:D:243:LEU:HD22	1.86	0.58
1:B:292:GLU:OE1	1:B:338:LYS:NZ	2.36	0.58
1:D:140:LEU:HA	1:D:143:ILE:HD12	1.87	0.57
1:B:300:LEU:HD11	1:B:338:LYS:HG2	1.86	0.56
1:D:136:ASN:HA	1:D:171:PHE:HE1	1.71	0.56
1:C:258:PHE:HD2	1:C:258:PHE:H	1.53	0.55
1:D:165:ARG:NH1	1:D:206:THR:O	2.41	0.54
1:D:165:ARG:NH2	1:D:206:THR:O	2.43	0.52
1:A:292:GLU:OE1	1:A:338:LYS:NZ	2.44	0.51
1:B:152:LEU:HD11	1:B:195:THR:HB	1.93	0.51
1:C:300:LEU:HD11	1:C:338:LYS:HG2	1.92	0.51
1:D:165:ARG:NH1	1:D:208:ASN:HB2	2.27	0.50
1:A:245:GLU:HG2	1:A:295[A]:HIS:NE2	2.27	0.50
1:D:214:ASN:OD1	1:D:249:ASP:N	2.35	0.49
1:C:144:VAL:HG13	1:C:184:ALA:HB2	1.93	0.49
1:D:293:ASN:OD1	1:D:296:LYS:N	2.27	0.49
1:D:172:ARG:NH2	1:D:177:GLY:O	2.43	0.47
2:A:489:HOH:O	1:C:387:HIS:HD2	1.98	0.47
1:C:245:GLU:HG2	1:C:295:HIS:NE2	2.29	0.46
1:D:229:HIS:CE1	1:D:254:ALA:HB2	2.50	0.46
1:B:164:LYS:NZ	2:B:533:HOH:O	2.49	0.46
1:C:234:ARG:HA	1:C:234:ARG:HD3	1.74	0.46
1:C:258:PHE:CD2	1:C:258:PHE:N	2.84	0.46
1:C:364:ALA:HB2	1:C:372:PHE:CE1	2.52	0.45
1:A:145:SER:HB3	1:A:183:LYS:HE3	1.99	0.44
1:D:264:GLU:HG2	1:D:266:GLY:H	1.82	0.44
1:B:181:LEU:HB3	1:B:182:PRO:HD3	2.00	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:173:GLU:HA	1:D:174:PRO:HD3	1.85	0.43
1:D:196:ILE:HB	1:D:197:PRO:HD3	2.01	0.43
1:A:152:LEU:HD11	1:A:195:THR:HB	2.00	0.42
1:C:258:PHE:HD2	1:C:258:PHE:N	2.14	0.42
1:A:204:GLU:HA	1:A:209:MET:HB2	2.01	0.42
1:D:181:LEU:HB3	1:D:182:PRO:HD3	2.00	0.42
1:D:252:ALA:O	1:D:274:PRO:HD3	2.20	0.42
1:A:150:ASP:O	1:A:153:GLU:HG3	2.19	0.42
1:B:378[B]:ARG:NH1	1:B:379:GLU:OE2	2.47	0.42
1:D:137:ARG:HB3	1:D:138:PRO:HD3	2.00	0.42
1:D:148:SER:HA	1:D:149:PRO:HD2	1.85	0.42
1:D:134:VAL:HA	1:D:158:PHE:CE1	2.55	0.42
1:D:219:ASP:O	1:D:257:ARG:HD3	2.20	0.41
1:D:195:THR:O	1:D:199:LEU:HG	2.20	0.41
1:D:193:ASN:O	1:D:196:ILE:HG12	2.20	0.41
1:C:181:LEU:HB3	1:C:182:PRO:HD3	2.02	0.41
1:A:217:PHE:CZ	1:A:230:ILE:HD11	2.56	0.41
1:A:375:ILE:O	1:A:379:GLU:HG3	2.21	0.41
1:A:181:LEU:HB3	1:A:182:PRO:HD3	2.03	0.41
1:C:137:ARG:HB2	1:C:170:GLU:O	2.21	0.41
1:C:272:GLU:OE1	1:C:304:ASP:HB2	2.20	0.40
1:B:296:LYS:HB2	1:B:296:LYS:HE3	1.92	0.40
1:C:193:ASN:O	1:C:196:ILE:HG13	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/260 (100%)	254 (98%)	6 (2%)	0	100	100
1	B	259/260 (100%)	253 (98%)	6 (2%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	251/260 (96%)	245 (98%)	6 (2%)	0	100	100
1	D	248/260 (95%)	241 (97%)	7 (3%)	0	100	100
All	All	1018/1040 (98%)	993 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/221 (101%)	220 (99%)	3 (1%)	73	78
1	B	222/221 (100%)	219 (99%)	3 (1%)	71	76
1	C	216/221 (98%)	214 (99%)	2 (1%)	82	87
1	D	213/221 (96%)	211 (99%)	2 (1%)	82	87
All	All	874/884 (99%)	864 (99%)	10 (1%)	82	82

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	327	PHE
1	A	378[A]	ARG
1	A	378[B]	ARG
1	B	327	PHE
1	B	378[A]	ARG
1	B	378[B]	ARG
1	C	258	PHE
1	C	327	PHE
1	D	220	VAL
1	D	327	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	256/260 (98%)	-0.10	9 (3%) 44 45	16, 27, 68, 105	0
1	B	256/260 (98%)	-0.06	13 (5%) 29 29	15, 28, 70, 142	0
1	C	252/260 (96%)	-0.13	9 (3%) 43 44	29, 43, 65, 119	0
1	D	252/260 (96%)	1.23	69 (27%) 1 1	37, 84, 125, 186	0
All	All	1016/1040 (97%)	0.23	100 (9%) 8 8	15, 41, 108, 186	0

All (100) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	258	PHE	10.4
1	D	134	VAL	10.4
1	D	159	LEU	10.1
1	D	294	GLY	9.0
1	D	157	SER	8.6
1	C	266	GLY	8.5
1	D	171	PHE	8.3
1	B	261	PRO	7.5
1	D	139	ILE	6.8
1	C	265	GLY	6.7
1	D	135	PHE	6.7
1	D	265	GLY	6.3
1	A	261	PRO	6.1
1	D	266	GLY	6.1
1	B	260	GLN	6.1
1	D	170	GLU	6.0
1	D	258	PHE	6.0
1	D	163	LYS	5.9
1	D	207	GLY	5.8
1	A	262	LYS	5.7
1	D	138	PRO	5.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	160	LEU	5.5
1	D	292	GLU	4.9
1	D	155	LEU	4.8
1	D	238	HIS	4.8
1	D	210	ARG	4.6
1	B	384	ALA	4.4
1	D	136	ASN	4.4
1	D	166	LEU	4.3
1	D	340	ALA	4.3
1	B	262	LYS	4.3
1	D	389	HIS	4.2
1	D	156	LEU	4.2
1	C	134	VAL	4.1
1	D	190	ALA	4.1
1	C	259	PHE	4.0
1	D	154	GLY	3.8
1	D	152	LEU	3.5
1	D	341	LYS	3.5
1	B	265	GLY	3.5
1	D	297	GLN	3.3
1	C	267	TYR	3.3
1	D	257	ARG	3.2
1	D	168	ASP	3.1
1	D	205	LYS	3.1
1	D	203	ALA	3.1
1	A	260	GLN	3.0
1	A	385	ALA	3.0
1	D	226	THR	3.0
1	B	386	HIS	3.0
1	D	247	GLY	2.9
1	D	267	TYR	2.9
1	D	148	SER	2.9
1	D	200	LEU	2.9
1	D	158	PHE	2.9
1	D	165	ARG	2.9
1	D	192	ARG	2.8
1	D	175	SER	2.8
1	D	174	PRO	2.7
1	D	191	GLY	2.7
1	D	256	GLY	2.7
1	D	209	MET	2.7
1	B	385	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	133	LYS	2.6
1	D	164	LYS	2.6
1	A	386	HIS	2.5
1	B	387	HIS	2.5
1	D	198	ILE	2.5
1	A	265	GLY	2.5
1	D	189	SER	2.5
1	A	387	HIS	2.5
1	D	289	TYR	2.4
1	D	295	HIS	2.4
1	D	161	THR	2.4
1	D	173	GLU	2.4
1	D	201	ASP	2.4
1	B	258	PHE	2.3
1	D	264	GLU	2.3
1	D	162	HIS	2.3
1	B	380	ILE	2.3
1	C	153	GLU	2.3
1	D	150	ASP	2.2
1	D	343	PHE	2.2
1	D	285	HIS	2.2
1	D	147	GLY	2.2
1	A	132	MET	2.2
1	B	383	ALA	2.2
1	D	153	GLU	2.2
1	D	141	PHE	2.2
1	C	264	GLU	2.1
1	C	388	HIS	2.1
1	D	167	THR	2.1
1	D	385	ALA	2.1
1	D	354	ASN	2.1
1	B	259	PHE	2.1
1	D	213	ILE	2.1
1	D	337	ILE	2.0
1	A	382	ASP	2.0
1	D	142	ASP	2.0
1	B	322	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.