



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 12:39 am GMT

PDB ID : 4W9K  
Title : pVHL:EloB:EloC in complex with (2S,4R)-1-((S)-2-((S)-2-acetamido-3-phenylpropanamido)-3,3-dimethylbutanoyl)-4-hydroxy-N-(4-(4-methylthiazol-5-yl)benzyl)pyrrolidine-2-carboxamide (ligand 14)  
Authors : Gadd, M.S.; Galdeano, C.; van Molle, I.; Ciulli, A.  
Deposited on : 2014-08-27  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	<div> <div>%</div> <div>95%</div> <div>• •</div> </div>
1	D	104	<div> <div>%</div> <div>95%</div> <div>• •</div> </div>
1	G	104	<div> <div>5%</div> <div>95%</div> <div>• •</div> </div>
1	J	104	<div> <div>5%</div> <div>97%</div> <div>• •</div> </div>
2	B	97	<div> <div>2%</div> <div>87%</div> <div>• 11%</div> </div>
2	E	97	<div> <div>90%</div> <div>• 9%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	97	<div><div></div><div>6%</div><div>84%</div><div>7%</div><div>9%</div></div>
2	K	97	<div><div></div><div>3%</div><div>77%</div><div>12%</div><div>9%</div></div>
3	C	162	<div><div></div><div>3%</div><div>81%</div><div>5%</div><div>13%</div></div>
3	F	162	<div><div></div><div>3%</div><div>87%</div><div></div><div>11%</div></div>
3	I	162	<div><div></div><div>10%</div><div>84%</div><div></div><div>12%</div></div>
3	L	162	<div><div></div><div>4%</div><div>82%</div><div>6%</div><div>12%</div></div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 11022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription elongation factor B polypeptide 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	103	Total	As	C	N	O	S	0	0	0
			809	2	514	136	152	5			
1	D	102	Total	As	C	N	O	S	0	0	0
			791	2	505	132	147	5			
1	G	103	Total	As	C	N	O	S	0	0	0
			812	2	515	136	154	5			
1	J	103	Total	As	C	N	O	S	0	0	0
			812	2	515	136	154	5			

- Molecule 2 is a protein called Transcription elongation factor B polypeptide 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	86	Total	C	N	O	S	0	0	0
			682	440	109	127	6			
2	E	88	Total	C	N	O	S	0	0	0
			689	444	111	128	6			
2	H	88	Total	C	N	O	S	0	0	0
			684	440	111	127	6			
2	K	88	Total	C	N	O	S	0	0	0
			691	448	111	125	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	initiating methionine	UNP Q15369
E	16	MET	-	initiating methionine	UNP Q15369
H	16	MET	-	initiating methionine	UNP Q15369
K	16	MET	-	initiating methionine	UNP Q15369


- Molecule 3 is a protein called Von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	141	Total 1125	As 1	C 717	N 200	O 205	S 2	0	1	0
3	F	144	Total 1128	As 1	C 722	N 201	O 202	S 2	0	0	0
3	I	142	Total 1133	As 1	C 724	N 202	O 204	S 2	0	0	0
3	L	143	Total 1143	As 1	C 730	N 206	O 204	S 2	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	52	GLY	-	expression tag	UNP P40337
C	53	SER	-	expression tag	UNP P40337
F	52	GLY	-	expression tag	UNP P40337
F	53	SER	-	expression tag	UNP P40337
I	52	GLY	-	expression tag	UNP P40337
I	53	SER	-	expression tag	UNP P40337
L	52	GLY	-	expression tag	UNP P40337
L	53	SER	-	expression tag	UNP P40337

- Molecule 4 is N-acetyl-L-phenylalanyl-3-methyl-L-valyl-(4R)-4-hydroxy-N-[4-(4-methyl-1,3-thiazol-5-yl)benzyl]-L-prolinamide (three-letter code: 3JO) (formula: C<sub>33</sub>H<sub>41</sub>N<sub>5</sub>O<sub>5</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	C	1	Total	C	N	O	S	0	0
			44	33	5	5	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	F	1	Total 44	C 33	N 5	O 5	S 1	0	0
4	I	1	Total 44	C 33	N 5	O 5	S 1	0	0
4	L	1	Total 44	C 33	N 5	O 5	S 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	61	Total 61	O 61	0	0
5	B	24	Total 24	O 24	0	0
5	C	31	Total 31	O 31	0	0
5	D	15	Total 15	O 15	0	0
5	E	17	Total 17	O 17	0	0
5	F	27	Total 27	O 27	0	0
5	G	25	Total 25	O 25	0	0
5	H	18	Total 18	O 18	0	0
5	I	27	Total 27	O 27	0	0
5	J	44	Total 44	O 44	0	0
5	K	19	Total 19	O 19	0	0
5	L	39	Total 39	O 39	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription elongation factor B polypeptide 2




- Molecule 1: Transcription elongation factor B polypeptide 2





- Molecule 1: Transcription elongation factor B polypeptide 2




- Molecule 1: Transcription elongation factor B polypeptide 2




- Molecule 2: Transcription elongation factor B polypeptide 1





- Molecule 2: Transcription elongation factor B polypeptide 1

Chain E:  90% 9%





- Molecule 2: Transcription elongation factor B polypeptide 1

Chain H:  6% 84% 7% 9%





- Molecule 2: Transcription elongation factor B polypeptide 1

Chain K:  3% 77% 12% 9%





- Molecule 3: Von Hippel-Lindau disease tumor suppressor

Chain C:  3% 81% 5% 13%





- Molecule 3: Von Hippel-Lindau disease tumor suppressor

Chain F:  3% 87% 11%




- Molecule 3: Von Hippel-Lindau disease tumor suppressor

Chain I:  10% 84% 12%



- Molecule 3: Von Hippel-Lindau disease tumor suppressor

Chain L:  4% 82% 6% 12%





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.70Å 93.70Å 361.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.70 – 2.10 48.86 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.2 (93.70-2.10) 96.2 (48.86-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.215 , 0.260 0.220 , 0.260	Depositor DCC
$R_{free}$ test set	4637 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtriage
Anisotropy	0.068	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 43.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11022	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.14 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9411e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, 3JO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/805	0.67	0/1086
1	D	0.37	0/786	0.58	0/1059
1	G	0.39	0/808	0.62	0/1090
1	J	0.43	0/808	0.66	0/1090
2	B	0.43	0/696	0.58	0/940
2	E	0.36	0/703	0.53	0/949
2	H	0.35	0/697	0.52	0/940
2	K	0.41	0/705	0.58	0/951
3	C	0.39	0/1147	0.64	0/1568
3	F	0.35	0/1148	0.61	0/1573
3	I	0.38	0/1152	0.61	0/1574
3	L	0.46	0/1163	0.69	0/1590
All	All	0.40	0/10618	0.62	0/14410

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	809	0	800	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	791	0	777	2	0
1	G	812	0	802	2	0
1	J	812	0	802	1	0
2	B	682	0	674	2	0
2	E	689	0	674	1	0
2	H	684	0	674	3	0
2	K	691	0	685	9	0
3	C	1125	0	1094	7	0
3	F	1128	0	1085	1	0
3	I	1133	0	1105	5	0
3	L	1143	0	1116	6	0
4	C	44	0	41	2	0
4	F	44	0	41	2	0
4	I	44	0	41	2	0
4	L	44	0	41	2	0
5	A	61	0	0	0	0
5	B	24	0	0	1	0
5	C	31	0	0	0	0
5	D	15	0	0	0	0
5	E	17	0	0	0	0
5	F	27	0	0	0	0
5	G	25	0	0	0	0
5	H	18	0	0	1	0
5	I	27	0	0	1	0
5	J	44	0	0	0	0
5	K	19	0	0	1	0
5	L	39	0	0	0	0
All	All	11022	0	10452	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:37:ARG:NH1	1:G:41:GLU:OE1	2.30	0.65
3:L:120:ARG:NH2	3:L:197:ASP:OD2	2.36	0.57
1:D:81:ALA:O	1:D:83:ASP:N	2.37	0.56
3:C:77:CAS:CE2	2:K:45:MET:HE1	2.36	0.56
3:C:170:VAL:HG21	3:C:178:LEU:HD11	1.90	0.54
4:L:301:3JO:CAR	4:L:301:3JO:H1	2.38	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:301:3JO:H1	4:I:301:3JO:CAR	2.38	0.54
1:A:11:LYS:HG3	1:A:91:GLU:HG3	1.91	0.52
4:C:301:3JO:CAR	4:C:301:3JO:H1	2.40	0.52
5:H:215:HOH:O	3:I:184:LEU:HD21	2.10	0.52
1:G:80:ARG:HA	1:G:85:PHE:HA	1.94	0.49
3:I:73:GLN:HG2	3:I:108:ARG:NH2	2.28	0.49
3:I:120:ARG:NH1	5:I:422:HOH:O	2.40	0.49
4:F:301:3JO:H1	4:F:301:3JO:H7	1.96	0.48
3:L:170:VAL:CG1	3:L:174:ASN:HB2	2.45	0.46
3:I:170:VAL:HG21	3:I:178:LEU:HD11	1.97	0.46
3:C:105:THR:HG22	2:K:64:GLU:HG2	1.96	0.46
2:B:101:LEU:HD11	3:C:178:LEU:HD22	1.98	0.46
4:C:301:3JO:H1	4:C:301:3JO:H7	1.99	0.45
3:F:90:ASN:ND2	3:F:94:GLU:HB2	2.32	0.45
4:L:301:3JO:H7	4:L:301:3JO:H1	1.98	0.45
2:K:101:LEU:HD11	3:L:178:LEU:HD22	1.99	0.45
3:L:182:ARG:HA	3:L:185:TYR:CD2	2.52	0.45
4:F:301:3JO:CAR	4:F:301:3JO:H1	2.48	0.44
2:H:69:VAL:HG21	2:H:102:GLU:HB3	1.99	0.44
1:J:17:ASP:OD1	2:K:32:LYS:NZ	2.49	0.44
4:I:301:3JO:H1	4:I:301:3JO:H7	1.99	0.44
2:K:98:GLU:N	2:K:98:GLU:OE1	2.49	0.43
3:L:142:VAL:HG13	3:L:143:ASP:N	2.35	0.42
2:H:19:VAL:HG12	2:H:58:ASN:HB3	2.02	0.42
3:C:154:PRO:HG2	3:C:156:TYR:CE1	2.55	0.42
2:B:58:ASN:N	5:B:214:HOH:O	2.53	0.41
1:D:94:SER:O	2:E:68:HIS:HB3	2.20	0.41
3:I:184:LEU:HA	3:I:184:LEU:HD23	1.92	0.41
2:K:38:THR:HG23	2:K:80:LYS:CD	2.50	0.41
2:H:72:LYS:HG3	2:H:99:ILE:CD1	2.51	0.41
2:K:68:HIS:O	5:K:205:HOH:O	2.21	0.41
3:L:170:VAL:HG13	3:L:174:ASN:HB2	2.03	0.41
2:K:69:VAL:HG21	2:K:102:GLU:HB3	2.03	0.40
3:C:73:GLN:OE1	3:C:108:ARG:NH1	2.54	0.40
3:C:148:PHE:CZ	2:K:45:MET:HG2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/104 (95%)	95 (96%)	3 (3%)	1 (1%)	18	12
1	D	96/104 (92%)	92 (96%)	4 (4%)	0	100	100
1	G	99/104 (95%)	96 (97%)	3 (3%)	0	100	100
1	J	99/104 (95%)	95 (96%)	4 (4%)	0	100	100
2	B	82/97 (84%)	82 (100%)	0	0	100	100
2	E	84/97 (87%)	82 (98%)	2 (2%)	0	100	100
2	H	84/97 (87%)	81 (96%)	3 (4%)	0	100	100
2	K	84/97 (87%)	82 (98%)	2 (2%)	0	100	100
3	C	139/162 (86%)	136 (98%)	3 (2%)	0	100	100
3	F	141/162 (87%)	137 (97%)	4 (3%)	0	100	100
3	I	137/162 (85%)	131 (96%)	6 (4%)	0	100	100
3	L	140/162 (86%)	131 (94%)	8 (6%)	1 (1%)	25	20
All	All	1284/1452 (88%)	1240 (97%)	42 (3%)	2 (0%)	51	52

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	82	ASP
3	L	183	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/90 (96%)	85 (99%)	1 (1%)	75	81
1	D	82/90 (91%)	82 (100%)	0	100	100
1	G	87/90 (97%)	87 (100%)	0	100	100
1	J	87/90 (97%)	86 (99%)	1 (1%)	78	83
2	B	76/86 (88%)	76 (100%)	0	100	100
2	E	75/86 (87%)	75 (100%)	0	100	100
2	H	74/86 (86%)	73 (99%)	1 (1%)	71	78
2	K	75/86 (87%)	71 (95%)	4 (5%)	26	24
3	C	124/147 (84%)	123 (99%)	1 (1%)	85	89
3	F	121/147 (82%)	120 (99%)	1 (1%)	85	89
3	I	125/147 (85%)	125 (100%)	0	100	100
3	L	125/147 (85%)	125 (100%)	0	100	100
All	All	1137/1292 (88%)	1128 (99%)	9 (1%)	85	89

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	HIS
3	C	148	PHE
3	F	148	PHE
2	H	17	MET
1	J	37	ARG
2	K	17	MET
2	K	61	ASN
2	K	64	GLU
2	K	85	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	35	HIS
3	I	73	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CAS	A	60	1	6,8,9	0.87	0	2,9,11	1.57	1 (50%)
1	CAS	A	89	1	6,8,9	1.05	0	2,9,11	1.19	0
3	CAS	C	77	3	6,8,9	0.59	0	2,9,11	1.02	0
1	CAS	D	60	1	6,8,9	0.62	0	2,9,11	1.51	1 (50%)
1	CAS	D	89	1	6,8,9	1.02	0	2,9,11	1.20	0
3	CAS	F	77	3	6,8,9	0.67	0	2,9,11	1.15	0
1	CAS	G	60	1	6,8,9	0.80	0	2,9,11	1.54	0
1	CAS	G	89	1	6,8,9	0.98	0	2,9,11	1.24	0
3	CAS	I	77	3	6,8,9	0.59	0	2,9,11	1.26	0
1	CAS	J	60	1	6,8,9	0.78	0	2,9,11	1.56	0
1	CAS	J	89	1	6,8,9	1.18	0	2,9,11	1.35	0
3	CAS	L	77	3	6,8,9	0.66	0	2,9,11	1.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	60	1	-	0/0/7/9	0/0/0/0
1	CAS	A	89	1	-	0/0/7/9	0/0/0/0
3	CAS	C	77	3	-	0/0/7/9	0/0/0/0
1	CAS	D	60	1	-	0/0/7/9	0/0/0/0
1	CAS	D	89	1	-	0/0/7/9	0/0/0/0
3	CAS	F	77	3	-	0/0/7/9	0/0/0/0
1	CAS	G	60	1	-	0/0/7/9	0/0/0/0
1	CAS	G	89	1	-	0/0/7/9	0/0/0/0
3	CAS	I	77	3	-	0/0/7/9	0/0/0/0
1	CAS	J	60	1	-	0/0/7/9	0/0/0/0
1	CAS	J	89	1	-	0/0/7/9	0/0/0/0
3	CAS	L	77	3	-	0/0/7/9	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	60	CAS	O-C-CA	-2.14	119.10	125.02
1	D	60	CAS	O-C-CA	-2.04	119.39	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	77	CAS	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	3JO	C	301	-	44,47,47	2.60	7 (15%)	56,67,67	1.16	4 (7%)
4	3JO	F	301	-	44,47,47	2.34	6 (13%)	56,67,67	1.13	4 (7%)
4	3JO	I	301	-	44,47,47	2.24	6 (13%)	56,67,67	1.16	4 (7%)
4	3JO	L	301	-	44,47,47	2.37	6 (13%)	56,67,67	1.21	4 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means



no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	3JO	C	301	-	-	0/43/55/55	0/4/4/4
4	3JO	F	301	-	-	0/43/55/55	0/4/4/4
4	3JO	I	301	-	-	0/43/55/55	0/4/4/4
4	3JO	L	301	-	-	0/43/55/55	0/4/4/4

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	301	3JO	CBK-CBL	-13.39	1.35	1.48
4	L	301	3JO	CBK-CBL	-12.01	1.36	1.48
4	F	301	3JO	CBK-CBL	-11.35	1.37	1.48
4	I	301	3JO	CBK-CBL	-11.16	1.37	1.48
4	L	301	3JO	CBO-CBE	-5.81	1.40	1.52
4	C	301	3JO	CBO-CBE	-5.58	1.40	1.52
4	F	301	3JO	CBO-CBE	-5.48	1.41	1.52
4	F	301	3JO	CAU-CBH	-5.39	1.39	1.51
4	I	301	3JO	CBO-CBE	-5.19	1.41	1.52
4	C	301	3JO	CAU-CBH	-5.09	1.39	1.51
4	I	301	3JO	CAU-CBH	-4.28	1.41	1.51
4	I	301	3JO	CB-CG	-4.10	1.41	1.51
4	F	301	3JO	CB-CG	-3.86	1.42	1.51
4	L	301	3JO	CB-CG	-3.84	1.42	1.51
4	C	301	3JO	CB-CG	-3.78	1.42	1.51
4	L	301	3JO	CAU-CBH	-3.64	1.43	1.51
4	C	301	3JO	CAT-SBC	-2.36	1.61	1.68
4	F	301	3JO	CAX-NBQ	2.17	1.50	1.47
4	C	301	3JO	CAB-CBJ	2.17	1.54	1.50
4	I	301	3JO	CBO-NBQ	2.28	1.51	1.47
4	L	301	3JO	CAX-NBQ	2.37	1.51	1.47
4	I	301	3JO	CAB-CBJ	2.37	1.54	1.50
4	F	301	3JO	CBO-NBQ	2.52	1.52	1.47
4	C	301	3JO	CAX-NBQ	2.55	1.51	1.47
4	L	301	3JO	CAB-CBJ	2.92	1.55	1.50

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	L	301	3JO	CBR-CBP-NBB	-4.06	107.03	111.85
4	C	301	3JO	CBR-CBP-NBB	-3.69	107.46	111.85
4	F	301	3JO	CG-CB-CA	-3.31	104.15	113.41
4	L	301	3JO	CG-CB-CA	-3.03	104.94	113.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	I	301	3JO	CBR-CBP-NBB	-2.96	108.34	111.85
4	C	301	3JO	CAX-NBQ-CBO	-2.96	107.21	111.69
4	C	301	3JO	CG-CB-CA	-2.75	105.70	113.41
4	F	301	3JO	CBR-CBP-NBB	-2.40	109.00	111.85
4	I	301	3JO	CG-CB-CA	-2.39	106.71	113.41
4	F	301	3JO	CAX-NBQ-CBO	-2.09	108.53	111.69
4	C	301	3JO	CA-C-NBB	-2.07	112.11	116.78
4	F	301	3JO	CBM-CAX-NBQ	2.05	105.34	103.08
4	L	301	3JO	CAU-NAZ-CBE	2.18	125.34	122.39
4	I	301	3JO	CAU-CBH-CAQ	2.29	125.77	120.89
4	L	301	3JO	CAU-CBH-CAQ	2.39	126.00	120.89
4	I	301	3JO	CAU-NAZ-CBE	2.43	125.69	122.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	301	3JO	2	0
4	F	301	3JO	2	0
4	I	301	3JO	2	0
4	L	301	3JO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	101/104 (97%)	-0.05	1 (0%) 82 85	24, 35, 72, 87	0
1	D	100/104 (96%)	0.27	1 (1%) 82 85	34, 58, 98, 117	0
1	G	101/104 (97%)	0.33	5 (4%) 30 36	36, 54, 86, 101	0
1	J	101/104 (97%)	0.16	5 (4%) 30 36	24, 36, 75, 128	0
2	B	86/97 (88%)	0.08	2 (2%) 61 66	25, 37, 67, 89	0
2	E	88/97 (90%)	0.20	0 100 100	34, 51, 82, 89	0
2	H	88/97 (90%)	0.60	6 (6%) 18 23	39, 57, 88, 102	0
2	K	88/97 (90%)	0.28	3 (3%) 46 53	28, 45, 75, 95	0
3	C	140/162 (86%)	0.29	5 (3%) 43 50	33, 50, 86, 117	0
3	F	143/162 (88%)	0.29	5 (3%) 44 51	36, 55, 89, 113	0
3	I	141/162 (87%)	0.55	17 (12%) 5 6	34, 52, 114, 138	0
3	L	142/162 (87%)	0.27	6 (4%) 37 44	26, 40, 79, 96	0
All	All	1319/1452 (90%)	0.28	56 (4%) 37 44	24, 48, 88, 138	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	I	185	TYR	6.8
1	J	102	VAL	6.1
3	I	178	LEU	6.0
2	H	46	LEU	5.8
3	C	140	LEU	5.5
3	I	188	LEU	4.9
3	I	171	LYS	4.6
1	J	103	MET	4.5
3	I	170	VAL	4.3
1	J	101	ASP	4.2
3	I	184	LEU	4.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
3	C	143	ASP	4.1
3	L	181	VAL	4.0
3	I	189	GLU	4.0
3	I	179	ASP	3.9
3	I	181	VAL	3.8
2	K	57	THR	3.8
3	F	185	TYR	3.4
3	C	62	VAL	3.2
1	J	100	PRO	3.2
2	H	57	THR	3.1
3	I	175	TYR	3.0
3	L	142	VAL	2.9
1	G	102	VAL	2.8
1	G	81	ALA	2.8
2	K	101	LEU	2.6
2	K	46	LEU	2.6
3	C	176	ARG	2.6
1	G	24	VAL	2.6
3	F	140	LEU	2.6
3	I	180	ILE	2.5
3	I	174	ASN	2.5
2	H	33	ARG	2.5
1	G	84	THR	2.4
3	I	169	LEU	2.4
3	I	176	ARG	2.4
2	H	19	VAL	2.4
3	L	176	ARG	2.4
3	F	148	PHE	2.4
3	I	186	GLU	2.3
1	G	1	MET	2.3
2	H	45	MET	2.3
3	I	172	PRO	2.3
1	J	99	LEU	2.2
3	C	147	ILE	2.2
2	H	101	LEU	2.2
3	F	201	LEU	2.2
2	B	87	SER	2.2
1	D	81	ALA	2.2
3	L	186	GLU	2.1
3	L	144	GLY	2.1
2	B	17	MET	2.1
3	F	147	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	102	VAL	2.0
3	L	170	VAL	2.0
3	I	62	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CAS	D	60	9/10	0.97	0.11	-	37,40,61,64	0
1	CAS	A	60	9/10	0.95	0.11	-	28,31,55,57	0
1	CAS	D	89	9/10	0.92	0.15	-	68,77,125,128	0
1	CAS	A	89	9/10	0.95	0.15	-	41,48,84,86	0
1	CAS	G	89	9/10	0.92	0.13	-	45,54,104,104	0
3	CAS	C	77	9/10	0.99	0.10	-	39,42,64,64	0
1	CAS	J	89	9/10	0.97	0.13	-	32,37,67,70	0
1	CAS	G	60	9/10	0.97	0.13	-	52,54,71,72	0
3	CAS	F	77	9/10	0.98	0.10	-	43,50,72,73	0
3	CAS	L	77	9/10	0.99	0.10	-	27,30,47,51	0
1	CAS	J	60	9/10	0.96	0.13	-	32,36,67,69	0
3	CAS	I	77	9/10	0.99	0.12	-	35,38,55,57	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	3JO	F	301	44/44	0.92	0.18	0.31	44,56,85,92	0
4	3JO	I	301	44/44	0.96	0.14	0.15	29,45,71,76	0
4	3JO	C	301	44/44	0.96	0.15	0.10	33,42,65,81	0
4	3JO	L	301	44/44	0.97	0.14	-0.10	24,33,64,70	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.