



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2017 – 04:16 PM EDT

PDB ID : 4WAM  
Title : H. influenzae beta-carbonic anhydrase variant W39V/G41A/P48S/A49P  
Authors : Rowlett, R.S.; Hoffman, K.M.  
Deposited on : unknown  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	<div> <div>3%</div> <div>60%</div> <div>15%</div> <div>22%</div> </div>
1	B	229	<div> <div>6%</div> <div>62%</div> <div>15%</div> <div>21%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	302	-	-	-	X
3	PO4	A	303	-	-	X	-
3	PO4	A	304	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-carbonic anhydrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	0	0
			1410	900	252	251	7			
1	B	180	Total	C	N	O	S	0	1	0
			1433	912	255	259	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	VAL	TRP	engineered mutation	UNP P45148
A	41	ALA	GLY	engineered mutation	UNP P45148
A	48	SER	PRO	engineered mutation	UNP P45148
A	49	PRO	ALA	engineered mutation	UNP P45148
B	39	VAL	TRP	engineered mutation	UNP P45148
B	41	ALA	GLY	engineered mutation	UNP P45148
B	48	SER	PRO	engineered mutation	UNP P45148
B	49	PRO	ALA	engineered mutation	UNP P45148

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	82	Total	O	0	0
			82	82		
4	B	62	Total	O	0	0
			62	62		



- Molecule 1: beta-carbonic anhydrase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.16Å 130.26Å 143.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	96.56 – 2.20 12.01 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.3 (96.56-2.20) 96.0 (12.01-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.41 (at 2.21Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.232 , 0.287 0.238 , 0.287	Depositor DCC
$R_{free}$ test set	1164 reflections (5.48%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.491	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 49.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	3009	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.76	0/1437	0.90	2/1943 (0.1%)
1	B	0.77	0/1465	0.95	3/1985 (0.2%)
All	All	0.77	0/2902	0.92	5/3928 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	182	ASP	CB-CG-OD1	6.24	123.92	118.30
1	B	109	ALA	C-N-CA	-6.14	106.35	121.70
1	A	125	ASP	CB-CG-OD1	6.11	123.80	118.30
1	A	211	ARG	NE-CZ-NH2	-5.87	117.36	120.30
1	B	46	ARG	NE-CZ-NH1	5.18	122.89	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	50	GLU	Peptide

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	B	34	THR	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1410	0	1421	27	0
1	B	1433	0	1432	20	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	20	0	0	8	0
4	A	82	0	0	1	1
4	B	62	0	0	2	0
All	All	3009	0	2853	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:ARG:NH1	3:A:303:PO4:O1	2.07	0.86
1:B:181:TYR:HE2	1:B:183:VAL:HG22	1.44	0.82
1:B:181:TYR:CE2	1:B:183:VAL:HG22	2.20	0.75
3:A:304:PO4:O3	1:B:110:ASP:OD2	2.07	0.72
1:B:183:VAL:HG23	4:B:454:HOH:O	1.93	0.68
1:A:110:ASP:OD2	3:A:305:PO4:O1	2.13	0.66
1:A:62:VAL:HG11	3:A:303:PO4:O3	1.98	0.63
1:A:127:TRP:HE3	1:A:146:LEU:HD22	1.69	0.58
1:A:51:LYS:HD3	1:A:52:LEU:HB3	1.85	0.58
1:A:48:SER:CB	1:A:50:GLU:HG3	2.35	0.56
1:A:188:LEU:O	1:A:189:VAL:C	2.45	0.55
1:B:183:VAL:HG21	4:B:462:HOH:O	2.06	0.54
1:B:34:THR:HG22	1:B:35:PRO:HD3	1.90	0.54
1:A:49:PRO:HD3	3:A:303:PO4:O1	2.07	0.53
1:A:48:SER:HB2	1:A:50:GLU:HG3	1.89	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:VAL:O	1:A:67:ALA:HB3	2.09	0.53
1:A:211:ARG:NH2	4:A:402:HOH:O	2.41	0.53
1:B:69:GLN:HE22	1:B:122:HIS:HB2	1.74	0.52
1:B:185:ASP:HB3	1:B:187:PHE:CD2	2.46	0.50
1:B:47:VAL:HG23	1:B:51:LYS:HD2	1.92	0.50
1:B:137:LEU:HD21	1:B:215:LEU:HD22	1.93	0.48
1:A:64:ARG:HD2	3:A:303:PO4:O1	2.13	0.48
1:A:34:THR:HB	1:A:35:PRO:CD	2.43	0.48
1:A:47:VAL:HG21	1:A:181:TYR:CE2	2.50	0.47
1:A:150:ASN:O	1:A:154:GLN:HG2	2.14	0.47
1:B:191:GLN:O	1:B:193:VAL:N	2.46	0.47
1:B:181:TYR:OH	1:B:183:VAL:HG13	2.15	0.47
1:A:137:LEU:HD12	1:A:219:ASN:HB3	1.97	0.47
1:B:150:ASN:O	1:B:154:GLN:HG2	2.16	0.46
1:A:181:TYR:HA	1:A:188:LEU:HD23	1.98	0.46
1:B:113:LEU:HB2	1:B:117:ASN:OD1	2.15	0.45
1:A:51:LYS:C	1:A:51:LYS:HD3	2.36	0.45
1:A:48:SER:HB3	1:A:50:GLU:HG3	1.99	0.45
1:A:39:VAL:HG11	1:A:49:PRO:HB3	1.98	0.45
1:B:202:GLU:OE1	1:B:206:ARG:NH1	2.50	0.44
1:A:156:TYR:O	1:A:160:ARG:HD3	2.18	0.44
1:B:166:SER:O	1:B:170:ARG:HG2	2.18	0.43
1:A:49:PRO:CD	3:A:303:PO4:O1	2.67	0.43
1:B:191:GLN:C	1:B:193:VAL:H	2.22	0.43
1:A:64:ARG:HH11	3:A:303:PO4:P	2.37	0.42
1:B:98:HIS:O	1:B:147:THR:HG21	2.20	0.42
1:B:93:ILE:HG21	1:B:158:LEU:HD21	2.02	0.42
1:A:34:THR:HB	1:A:35:PRO:HD3	2.02	0.41
1:A:93:ILE:HG21	1:A:158:LEU:HD21	2.01	0.41
1:B:191:GLN:O	1:B:193:VAL:HG23	2.21	0.41
1:A:155:VAL:HG11	1:A:193:VAL:CG1	2.52	0.40
1:A:217:GLU:O	1:A:220:ILE:HG22	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:464:HOH:O	4:A:464:HOH:O[7_455]	1.80	0.40

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	170/229 (74%)	165 (97%)	4 (2%)	1 (1%)	28	29
1	B	177/229 (77%)	172 (97%)	4 (2%)	1 (1%)	28	29
All	All	347/458 (76%)	337 (97%)	8 (2%)	2 (1%)	28	29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	50	GLU
1	B	192	GLY

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/202 (76%)	141 (92%)	13 (8%)	13	12
1	B	157/202 (78%)	146 (93%)	11 (7%)	18	19
All	All	311/404 (77%)	287 (92%)	24 (8%)	15	15

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	ARG
1	A	48	SER
1	A	50	GLU
1	A	51	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	52	LEU
1	A	75	PHE
1	A	78	LEU
1	A	133	LEU
1	A	194	MET
1	A	196	THR
1	A	212	LEU
1	A	217	GLU
1	A	221	LEU
1	B	34	THR
1	B	45	SER
1	B	46	ARG
1	B	64	ARG
1	B	183	VAL
1	B	184	ASN
1	B	188	LEU
1	B	189	VAL
1	B	194	MET
1	B	199	GLU
1	B	206	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	219	ASN
1	B	69	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	PO4	A	302	-	4,4,4	2.69	1 (25%)	6,6,6	1.21	0
3	PO4	A	303	-	4,4,4	2.39	3 (75%)	6,6,6	0.90	0
3	PO4	A	304	-	4,4,4	0.79	0	6,6,6	0.64	0
3	PO4	A	305	-	4,4,4	0.32	0	6,6,6	1.42	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	A	302	-	-	0/0/0/0	0/0/0/0
3	PO4	A	303	-	-	0/0/0/0	0/0/0/0
3	PO4	A	304	-	-	0/0/0/0	0/0/0/0
3	PO4	A	305	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	PO4	P-O3	-2.82	1.44	1.54
3	A	303	PO4	P-O4	-2.53	1.45	1.54
3	A	303	PO4	P-O2	-2.53	1.45	1.54
3	A	302	PO4	P-O1	4.73	1.60	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	305	PO4	O3-P-O2	2.78	118.12	107.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	PO4	6	0
3	A	304	PO4	1	0
3	A	305	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	178/229 (77%)	0.05	8 (4%) 34 32	9, 19, 58, 72	0
1	B	180/229 (78%)	0.15	13 (7%) 16 15	7, 22, 65, 74	0
All	All	358/458 (78%)	0.10	21 (5%) 23 22	7, 21, 62, 74	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	184	ASN	5.9
1	A	57	PRO	5.0
1	A	34	THR	4.8
1	B	217	GLU	4.5
1	B	192	GLY	4.3
1	A	187	PHE	4.2
1	B	34	THR	4.1
1	A	221	LEU	3.8
1	B	183	VAL	3.7
1	A	218	GLU	3.1
1	A	52	LEU	2.7
1	A	189	VAL	2.7
1	B	191	GLN	2.5
1	B	181	TYR	2.3
1	B	52	LEU	2.2
1	B	216	ASP	2.2
1	B	47	VAL	2.1
1	B	139	PRO	2.1
1	B	132	HIS	2.0
1	A	48	SER	2.0
1	B	187	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	PO4	A	302	5/5	0.90	0.17	5.98	16,17,17,26	0
3	PO4	A	304	5/5	0.97	0.17	2.94	46,47,51,53	0
3	PO4	A	305	5/5	0.79	0.26	2.00	55,56,66,66	0
3	PO4	A	303	5/5	0.95	0.29	1.90	10,12,13,14	5
2	ZN	A	301	1/1	0.99	0.05	-1.70	12,12,12,12	0
2	ZN	B	301	1/1	0.99	0.06	-1.91	15,15,15,15	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.