



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 12:09 pm GMT

PDB ID : 4WE4  
Title : The crystal structure of hemagglutinin from 1968 H3N2 influenza virus  
Authors : Yang, H.; Carney, P.J.; Chang, J.C.; Guo, Z.; Villanueva, J.M.; Stevens, J.  
Deposited on : 2014-09-09  
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 84%, grey 14%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>%</span> <span>84%</span> <span>14%</span> </div> </div>
2	B	172	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 1%, green 79%, grey 21%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>3%</span> <span>79%</span> <span>21%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	401	-	-	-	X
4	NAG	A	407	X	-	-	X
7	PEG	A	417	-	-	X	X
7	PEG	A	418	-	-	X	X
7	PEG	B	203	-	-	X	-
7	PEG	B	204	-	-	X	X
7	PEG	B	205	-	-	-	X
7	PEG	B	206	-	-	X	-
8	NAG	B	201	-	-	-	X
9	PE5	B	208	-	-	X	-

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 4484 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin HA1 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	321	Total	C	N	O	S	0	0	0
			2480	1552	436	479	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	ILE	VAL	conflict	UNP Q91MA7
A	329	GLN	ARG	conflict	UNP Q91MA7

- Molecule 2 is a protein called Hemagglutinin HA2 chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	172	Total	C	N	O	S	0	0	0
			1399	868	247	278	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	132	GLU	ASP	conflict	UNP Q91MA7
B	164	ASN	ASP	conflict	UNP Q91MA7

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	6	Total	C	N	O	0	0
			75	42	3	30		

- Molecule 5 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 6 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		
7	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 9 is 3,6,9,12,15,18,21,24-OCTAOXAHEXACOSAN-1-OL (three-letter code: PE5) (formula: C<sub>18</sub>H<sub>38</sub>O<sub>9</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			25	16	9		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	223	Total	O	0	0
			223	223		
10	B	77	Total	O	0	0
			77	77		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Hemagglutinin HA1 chain



#### • Molecule 2: Hemagglutinin HA2 chain





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	155.51Å 155.51Å 155.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.50 – 2.35 49.18 – 2.35	Depositor EDS
% Data completeness (in resolution range)	98.1 (30.50-2.35) 98.1 (49.18-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.94 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, $R_{free}$	0.193 , 0.236 0.192 , 0.235	Depositor DCC
$R_{free}$ test set	1306 reflections (5.10%)	DCC
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 41.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.023 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4484	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PE5, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.25	0/2536	0.45	0/3455
2	B	0.26	0/1423	0.41	0/1912
All	All	0.26	0/3959	0.44	0/5367

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	A	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	407	NAG	C1

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2480	0	2428	44	0
2	B	1399	0	1319	55	0
3	A	28	0	26	1	0
4	A	75	0	64	0	0
5	A	61	0	52	1	0
6	A	39	0	34	0	0
7	A	14	0	18	13	0
7	B	35	0	45	22	0
8	B	28	0	25	1	0
9	B	25	0	33	26	0
10	A	223	0	0	12	0
10	B	77	0	0	5	0
All	All	4484	0	4044	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:ARG:NH1	10:A:501:HOH:O	1.84	1.08
2:B:82:LYS:HA	7:B:204:PEG:H21	1.43	0.99
1:A:211:GLN:OE1	10:A:681:HOH:O	1.88	0.92
2:B:67:GLU:HB2	9:B:208:PE5:H121	1.52	0.91
2:B:66:ILE:HA	9:B:208:PE5:H101	1.54	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	319/321 (99%)	308 (97%)	10 (3%)	1 (0%)	44 53

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	170/172 (99%)	161 (95%)	9 (5%)	0	100	100
All	All	489/493 (99%)	469 (96%)	19 (4%)	1 (0%)	51	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	62	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/283 (100%)	277 (98%)	6 (2%)	59	72
2	B	147/147 (100%)	145 (99%)	2 (1%)	71	82
All	All	430/430 (100%)	422 (98%)	8 (2%)	62	75

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	MET
2	B	161	VAL
1	A	309	VAL
1	A	167	THR
1	A	216	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	211	GLN
2	B	65	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

16 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	NAG	A	402	1,4	14,14,15	0.71	0	15,19,21	1.17	2 (13%)
4	NAG	A	403	4	14,14,15	0.55	0	15,19,21	0.50	0
4	BMA	A	404	4	11,11,12	0.89	0	13,15,17	1.00	0
4	MAN	A	405	4	11,11,12	1.25	1 (9%)	13,15,17	1.05	1 (7%)
4	MAN	A	406	4	11,11,12	1.52	4 (36%)	13,15,17	1.89	2 (15%)
4	NAG	A	407	4	14,14,15	0.85	1 (7%)	15,19,21	0.82	1 (6%)
5	NAG	A	408	1,5	14,14,15	0.39	0	15,19,21	0.43	0
5	NAG	A	409	5	14,14,15	0.54	0	15,19,21	0.62	0
5	BMA	A	410	5	11,11,12	0.90	0	13,15,17	1.05	0
5	MAN	A	411	5	11,11,12	0.76	0	13,15,17	1.50	3 (23%)
5	MAN	A	412	5	11,11,12	0.90	0	13,15,17	1.47	2 (15%)
6	NAG	A	413	1,6	14,14,15	0.26	0	15,19,21	0.39	0
6	NAG	A	414	6	14,14,15	0.48	0	15,19,21	0.68	0
6	BMA	A	415	6	11,11,12	0.71	0	13,15,17	1.38	1 (7%)
8	NAG	B	201	8,2	14,14,15	0.67	1 (7%)	15,19,21	0.72	0
8	NAG	B	202	8	14,14,15	1.17	2 (14%)	15,19,21	0.78	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	402	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	403	4	-	0/6/23/26	0/1/1/1
4	BMA	A	404	4	-	0/2/19/22	0/1/1/1
4	MAN	A	405	4	-	0/2/19/22	0/1/1/1
4	MAN	A	406	4	-	0/2/19/22	1/1/1/1
4	NAG	A	407	4	1/1/5/7	0/6/23/26	0/1/1/1
5	NAG	A	408	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	409	5	-	0/6/23/26	0/1/1/1
5	BMA	A	410	5	-	0/2/19/22	0/1/1/1
5	MAN	A	411	5	-	0/2/19/22	0/1/1/1
5	MAN	A	412	5	-	0/2/19/22	0/1/1/1
6	NAG	A	413	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	414	6	-	0/6/23/26	0/1/1/1
6	BMA	A	415	6	-	0/2/19/22	1/1/1/1
8	NAG	B	201	8,2	-	0/6/23/26	0/1/1/1
8	NAG	B	202	8	-	0/6/23/26	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	MAN	O5-C1	-3.22	1.38	1.43
8	B	202	NAG	O5-C1	-2.55	1.39	1.43
8	B	201	NAG	O5-C1	-2.37	1.39	1.43
4	A	406	MAN	O2-C2	2.18	1.48	1.43
4	A	406	MAN	C4-C3	2.19	1.57	1.52

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	406	MAN	C1-C2-C3	-4.03	104.54	109.65
5	A	412	MAN	O2-C2-C3	-2.21	105.84	110.17
4	A	405	MAN	O2-C2-C3	-2.09	106.07	110.17
5	A	411	MAN	O2-C2-C3	-2.07	106.11	110.17
4	A	402	NAG	C3-C4-C5	2.24	114.17	110.22

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	407	NAG	C1

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	415	BMA	C1-C2-C3-C4-C5-O5
4	A	406	MAN	C1-C2-C3-C4-C5-O5

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	409	NAG	1	0
8	B	201	NAG	1	0

## 5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	401	1	14,14,15	0.23	0	15,19,21	0.52	0
3	NAG	A	416	1	14,14,15	0.19	0	15,19,21	0.64	0
7	PEG	A	417	-	6,6,6	0.62	0	5,5,5	0.29	0
7	PEG	A	418	-	6,6,6	0.57	0	5,5,5	0.30	0
7	PEG	B	203	-	6,6,6	0.60	0	5,5,5	0.46	0
7	PEG	B	204	-	6,6,6	0.65	0	5,5,5	0.31	0
7	PEG	B	205	-	6,6,6	0.56	0	5,5,5	0.37	0
7	PEG	B	206	-	6,6,6	0.63	0	5,5,5	0.33	0
7	PEG	B	207	-	6,6,6	0.57	0	5,5,5	0.25	0
9	PE5	B	208	-	24,24,26	0.65	0	23,23,25	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1	-	0/6/23/26	0/1/1/1
3	NAG	A	416	1	-	0/6/23/26	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PEG	A	417	-	-	0/4/4/4	0/0/0/0
7	PEG	A	418	-	-	0/4/4/4	0/0/0/0
7	PEG	B	203	-	-	0/4/4/4	0/0/0/0
7	PEG	B	204	-	-	0/4/4/4	0/0/0/0
7	PEG	B	205	-	-	0/4/4/4	0/0/0/0
7	PEG	B	206	-	-	0/4/4/4	0/0/0/0
7	PEG	B	207	-	-	0/4/4/4	0/0/0/0
9	PE5	B	208	-	-	0/22/22/24	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 62 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	NAG	1	0
7	A	417	PEG	10	0
7	A	418	PEG	4	0
7	B	203	PEG	4	0
7	B	204	PEG	9	0
7	B	205	PEG	1	0
7	B	206	PEG	8	0
9	B	208	PE5	26	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	321/321 (100%)	-0.13	3 (0%) 84 91	14, 28, 49, 71	0
2	B	172/172 (100%)	0.10	6 (3%) 44 57	11, 35, 56, 88	0
All	All	493/493 (100%)	-0.05	9 (1%) 69 78	11, 31, 53, 88	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	4	GLY	3.4
2	B	29	SER	3.1
2	B	58	LYS	2.6
2	B	171	PHE	2.5
2	B	77	ILE	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
8	NAG	B	201	14/15	0.65	0.32	6.00	55,92,110,114	0
4	NAG	A	407	14/15	0.61	0.23	4.94	66,78,89,95	0
5	NAG	A	409	14/15	0.90	0.22	1.20	32,44,59,65	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	A	408	14/15	0.93	0.13	-0.39	26,38,52,57	0
6	NAG	A	413	14/15	0.92	0.12	-0.58	28,40,49,71	0
4	NAG	A	402	14/15	0.89	0.15	-	38,49,56,59	0
4	NAG	A	403	14/15	0.78	0.22	-	56,64,78,80	0
8	NAG	B	202	14/15	0.45	0.33	-	91,110,122,124	0
4	BMA	A	404	11/12	0.86	0.22	-	84,90,100,101	0
6	NAG	A	414	14/15	0.81	0.24	-	56,75,100,105	0
5	BMA	A	410	11/12	0.77	0.21	-	67,83,92,95	0
4	MAN	A	406	11/12	0.83	0.19	-	82,89,96,100	0
6	BMA	A	415	11/12	0.76	0.22	-	99,102,111,112	0
5	MAN	A	412	11/12	0.51	0.20	-	86,95,109,111	0
4	MAN	A	405	11/12	0.65	0.24	-	84,96,108,109	0
5	MAN	A	411	11/12	0.56	0.30	-	79,100,103,104	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	PEG	A	417	7/7	0.86	0.28	3.88	19,26,38,41	0
7	PEG	A	418	7/7	0.89	0.28	3.84	19,25,37,40	0
3	NAG	A	401	14/15	0.73	0.29	3.54	43,67,75,76	0
7	PEG	B	205	7/7	0.81	0.23	2.59	24,33,42,50	0
7	PEG	B	204	7/7	0.89	0.25	2.27	18,24,37,40	0
7	PEG	B	206	7/7	0.94	0.24	1.98	21,29,39,39	0
9	PE5	B	208	25/27	0.88	0.24	1.67	8,31,48,56	0
7	PEG	B	203	7/7	0.86	0.22	1.19	29,37,41,46	0
3	NAG	A	416	14/15	0.83	0.14	0.36	41,60,71,75	0
7	PEG	B	207	7/7	0.91	0.13	-0.38	46,51,61,68	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.