



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 01:38 am GMT

PDB ID : 2WIT
Title : CRYSTAL STRUCTURE OF THE SODIUM-COUPLED GLYCINE BETTAINE SYMPORTER BETP FROM CORYNEBACTERIUM GLUTAMICUM WITH BOUND SUBSTRATE
Authors : Ressler, S.; Terwisscha Van Scheltinga, A.C.; Vonnrhein, C.; Ott, V.; Ziegler, C.
Deposited on : 2009-05-17
Resolution : 3.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1156 (3.42-3.30)
Clashscore	112137	1231 (3.42-3.30)
Ramachandran outliers	110173	1212 (3.42-3.30)
Sidechain outliers	110143	1211 (3.42-3.30)
RSRZ outliers	101464	1165 (3.42-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	566	<div> <div>11%</div> <div> <div></div> <div>66%</div> <div>27%</div> <div>• 6%</div> </div> </div>
1	B	566	<div> <div>9%</div> <div> <div></div> <div>61%</div> <div>26%</div> <div>• 12%</div> </div> </div>
1	C	566	<div> <div>5%</div> <div> <div></div> <div>62%</div> <div>27%</div> <div>• 10%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 11737 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GLYCINE BETAINES TRANSPORTER BETP.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	531	Total	C	N	O	S	Se	0	0	0
			4057	2659	676	706	1	15			
1	B	500	Total	C	N	O	S	Se	0	0	0
			3783	2497	606	664	1	15			
1	C	510	Total	C	N	O	S	Se	0	0	0
			3873	2551	629	677	1	15			


There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
A	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
A	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
B	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	44	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	45	ALA	GLU	ENGINEERED MUTATION	UNP P54582
C	46	ALA	GLU	ENGINEERED MUTATION	UNP P54582


- Molecule 2 is TRIMETHYL GLYCINE (three-letter code: BET) (formula: C₅H₁₂NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	5	1	2		
2	B	1	Total	C	N	O	0	0
			8	5	1	2		
2	C	1	Total	C	N	O	0	0
			8	5	1	2		



• Molecule 1: GLYCINE BETAINES TRANSPORTER BETP



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	118.09Å 129.42Å 182.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.47 – 3.35 39.37 – 3.35	Depositor EDS
% Data completeness (in resolution range)	90.4 (39.47-3.35) 90.6 (39.37-3.35)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.32Å)	Xtriage
Refinement program	BUSTER-TNT 2.7.0	Depositor
R, R_{free}	0.257 , 0.265 0.299 , 0.315	Depositor DCC
R_{free} test set	1883 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	46.7	Xtriage
Anisotropy	0.286	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.22$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.74	EDS
Total number of atoms	11737	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BET

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/4142	0.34	0/5609
1	B	0.24	0/3865	0.33	0/5245
1	C	0.25	0/3956	0.34	0/5365
All	All	0.24	0/11963	0.34	0/16219

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4057	0	4104	119	0
1	B	3783	0	3815	110	0
1	C	3873	0	3909	110	0
2	A	8	0	11	4	0
2	B	8	0	11	3	0
2	C	8	0	11	0	0
All	All	11737	0	11861	332	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 332 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:558:ARG:HG2	1:A:558:ARG:HH11	1.36	0.91
1:A:565:ARG:HH11	1:A:565:ARG:HG3	1.42	0.83
1:B:548:VAL:HG13	1:B:549:ILE:H	1.43	0.83
1:B:270:ILE:HG23	1:B:271:ILE:HG23	1.64	0.79
1:B:334:PRO:HG3	1:C:105:LEU:HB2	1.64	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/566 (93%)	486 (92%)	39 (7%)	2 (0%)	38	75
1	B	496/566 (88%)	454 (92%)	39 (8%)	3 (1%)	28	66
1	C	506/566 (89%)	466 (92%)	38 (8%)	2 (0%)	38	75
All	All	1529/1698 (90%)	1406 (92%)	116 (8%)	7 (0%)	32	70

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	225	ILE
1	B	225	ILE
1	B	547	ASP
1	C	225	ILE
1	A	431	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	415/424 (98%)	410 (99%)	5 (1%)	75	88
1	B	389/424 (92%)	385 (99%)	4 (1%)	80	90
1	C	397/424 (94%)	392 (99%)	5 (1%)	73	88
All	All	1201/1272 (94%)	1187 (99%)	14 (1%)	75	88

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	271	ILE
1	B	371	TRP
1	C	371	TRP
1	B	145	MSE
1	C	271	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	484	GLN
1	B	519	GLN
1	C	484	GLN
1	B	309	ASN
1	B	481	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BET	A	1001	-	4,7,7	1.08	0	7,10,10	2.29	1 (14%)
2	BET	B	1001	-	4,7,7	1.08	0	7,10,10	2.14	1 (14%)
2	BET	C	1001	-	4,7,7	1.04	0	7,10,10	2.13	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BET	A	1001	-	-	0/3/5/5	0/0/0/0
2	BET	B	1001	-	-	0/3/5/5	0/0/0/0
2	BET	C	1001	-	-	0/3/5/5	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	BET	C-CA-N	-6.05	109.73	114.74
2	B	1001	BET	C-CA-N	-5.66	110.06	114.74
2	C	1001	BET	C-CA-N	-5.63	110.08	114.74

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	BET	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1001	BET	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	516/566 (91%)	0.64	61 (11%) 5 5	13, 76, 164, 193	0
1	B	485/566 (85%)	0.41	49 (10%) 8 8	5, 64, 113, 211	0
1	C	495/566 (87%)	0.20	31 (6%) 21 21	5, 40, 104, 192	0
All	All	1496/1698 (88%)	0.42	141 (9%) 9 10	5, 60, 138, 211	0

The worst 5 of 141 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	ILE	9.1
1	A	270	ILE	7.4
1	A	275	SER	7.1
1	B	80	PHE	7.1
1	A	490	TRP	6.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	BET	C	1001	8/8	0.82	0.31	1.61	30,78,154,165	0
2	BET	B	1001	8/8	0.82	0.34	0.48	18,69,117,138	0
2	BET	A	1001	8/8	0.77	0.34	-0.02	3,108,128,151	0

6.5 Other polymers [i](#)

There are no such residues in this entry.