



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 14, 2017 – 07:21 AM EST

PDB ID : 4WK0  
Title : Metal Ion and Ligand Binding of Integrin  
Authors : Xia, W.; Springer, T.A.  
Deposited on : unknown  
Resolution : 1.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	7172 (1.80-1.76)
Clashscore	112137	8247 (1.80-1.76)
Ramachandran outliers	110173	8154 (1.80-1.76)
Sidechain outliers	110143	8153 (1.80-1.76)
RSRZ outliers	101464	7262 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	
2	B	445	
3	C	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CA	A	502	-	-	-	X
4	CA	A	503	-	-	-	X
4	CA	A	504	-	-	-	X
5	NAG	A	511	-	-	-	X
5	NAG	A	512	-	-	-	X

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 14311 atoms, of which 6699 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	449	6576	2160	3184	554	670	8	0	2	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	451	VAL	ILE	engineered mutation	UNP P08648

- Molecule 2 is a protein called Integrin beta-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	427	6590	2080	3269	562	655	24	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	195	THR	SER	conflict	UNP P05556


- Molecule 3 is a protein called ARG-GLY-ASP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	C	3	42	12	19	6	5	0	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		
4	A	4	Total	Ca	0	0
			4	4		

- Molecule 5 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	B	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	B	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
5	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
5	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
5	B	1	28	8	14	1	5	0	0

- Molecule 6 is BETA-D-MANNOSE (three-letter code: BMA) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
6	A	1	20	6	9	5	0	0

- Molecule 7 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			21	6	10	5		
7	A	1	Total	C	H	O	0	0
			20	6	9	5		
7	A	1	Total	C	H	O	0	0
			21	6	10	5		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Mg	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	383	Total	O	0	0
			383	383		
9	B	239	Total	O	0	0
			239	239		
9	C	7	Total	O	0	0
			7	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrin alpha-5



- Molecule 2: Integrin beta-1



- Molecule 3: ARG-GLY-ASP



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.06Å 117.07Å 167.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.90 – 1.78 47.94 – 1.78	Depositor EDS
% Data completeness (in resolution range)	98.2 (47.90-1.78) 98.2 (47.94-1.78)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.95 (at 1.78Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: dev_1760)	Depositor
R, $R_{free}$	0.186 , 0.223 0.188 , 0.223	Depositor DCC
$R_{free}$ test set	1994 reflections (1.76%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.3	Xtrriage
Anisotropy	0.226	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 54.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14311	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BMA, CA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3492	0.56	0/4760
2	B	0.35	0/3377	0.52	0/4559
3	C	0.33	0/22	0.57	0/27
All	All	0.36	0/6891	0.54	0/9346

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3392	3184	3194	13	0
2	B	3321	3269	3270	17	0
3	C	23	19	19	1	0
4	A	4	0	0	0	0
4	B	2	0	0	0	0
5	A	112	107	100	0	0
5	B	84	82	76	0	0
6	A	11	9	8	0	0
7	A	33	29	29	1	0
8	B	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	A	383	0	0	3	0
9	B	239	0	0	0	0
9	C	7	0	0	1	0
All	All	7612	6699	6696	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 31 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:30:ASN:ND2	2:B:32:THR:OG1	1.62	1.28
1:A:451:VAL:O	1:A:452:VAL:HG22	1.89	0.72
1:A:27:ARG:HH22	1:A:30:THR:HG22	1.61	0.66
1:A:258:THR:HG21	9:A:938:HOH:O	2.00	0.62
2:B:40:THR:OG1	2:B:41:SER:N	2.36	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/452 (99%)	430 (96%)	16 (4%)	1 (0%)	51	33
2	B	421/445 (95%)	405 (96%)	16 (4%)	0	100	100
3	C	1/3 (33%)	1 (100%)	0	0	100	100
All	All	869/900 (97%)	836 (96%)	32 (4%)	1 (0%)	55	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	270	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	357/358 (100%)	353 (99%)	4 (1%)	78	70
2	B	380/395 (96%)	378 (100%)	2 (0%)	91	89
3	C	2/2 (100%)	2 (100%)	0	100	100
All	All	739/755 (98%)	733 (99%)	6 (1%)	85	80

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	284	TYR
2	B	415	CYS
1	A	413	ASP
1	A	121	TRP
2	B	224	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	221	GLN
2	B	30	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 7 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	505	1,5	14,14,15	0.41	0	15,19,21	0.60	0
5	NAG	A	506	5,6	14,14,15	0.31	0	15,19,21	0.62	0
6	BMA	A	507	5,7	11,11,12	0.94	1 (9%)	13,15,17	0.95	1 (7%)
7	MAN	A	508	6	11,11,12	0.80	1 (9%)	13,15,17	0.99	0
7	MAN	A	509	7,6	11,11,12	0.78	0	13,15,17	0.97	0
7	MAN	A	510	7	11,11,12	0.87	1 (9%)	13,15,17	1.05	1 (7%)
5	NAG	A	511	1	14,14,15	0.37	0	15,19,21	0.53	0
5	NAG	A	512	1,5	14,14,15	0.37	0	15,19,21	0.44	0
5	NAG	A	513	5	14,14,15	0.20	0	15,19,21	0.58	0
5	NAG	A	514	1,5	14,14,15	0.46	0	15,19,21	0.59	0
5	NAG	A	515	5	14,14,15	0.40	0	15,19,21	0.39	0
5	NAG	A	516	1	14,14,15	0.15	0	15,19,21	0.48	0
5	NAG	B	504	2,5	14,14,15	0.81	1 (7%)	15,19,21	0.56	0
5	NAG	B	505	5	14,14,15	0.27	0	15,19,21	0.53	0
5	NAG	B	506	2,5	14,14,15	0.60	1 (7%)	15,19,21	0.62	0
5	NAG	B	507	5	14,14,15	0.30	0	15,19,21	0.50	0
5	NAG	B	508	2	14,14,15	0.16	0	15,19,21	0.55	0
5	NAG	B	509	2	14,14,15	0.41	0	15,19,21	0.81	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	505	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	506	5,6	-	0/6/23/26	0/1/1/1
6	BMA	A	507	5,7	-	0/2/19/22	0/1/1/1
7	MAN	A	508	6	-	0/2/19/22	0/1/1/1
7	MAN	A	509	7,6	-	0/2/19/22	0/1/1/1
7	MAN	A	510	7	-	0/2/19/22	0/1/1/1
5	NAG	A	511	1	-	0/6/23/26	0/1/1/1
5	NAG	A	512	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	513	5	-	0/6/23/26	0/1/1/1
5	NAG	A	514	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	515	5	-	0/6/23/26	0/1/1/1
5	NAG	A	516	1	-	0/6/23/26	0/1/1/1
5	NAG	B	504	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	505	5	-	0/6/23/26	0/1/1/1
5	NAG	B	506	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	507	5	-	0/6/23/26	0/1/1/1
5	NAG	B	508	2	-	0/6/23/26	0/1/1/1
5	NAG	B	509	2	-	0/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	504	NAG	O5-C1	-2.68	1.39	1.43
6	A	507	BMA	O5-C1	-2.52	1.39	1.43
7	A	510	MAN	O5-C1	-2.16	1.40	1.43
5	B	506	NAG	O5-C1	-2.06	1.40	1.43
7	A	508	MAN	O5-C1	-2.06	1.40	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	507	BMA	C1-C2-C3	-2.20	106.86	109.65
7	A	510	MAN	C1-O5-C5	2.23	115.24	112.17
5	B	509	NAG	C1-O5-C5	2.79	116.01	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	509	MAN	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	449/452 (99%)	0.60	44 (9%) 8   8	20, 32, 70, 132	0
2	B	427/445 (95%)	1.51	89 (20%) 1   1	21, 40, 124, 173	0
3	C	3/3 (100%)	0.45	0 100   100	48, 48, 55, 71	0
All	All	879/900 (97%)	1.04	133 (15%) 3   2	20, 35, 107, 173	0

The worst 5 of 133 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	33	PHE	17.5
2	B	34	LEU	17.1
2	B	47	LEU	16.1
2	B	31	SER	14.1
2	B	50	LEU	14.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	CA	A	503	1/1	0.94	0.34	16.75	68,68,68,68	0
4	CA	A	502	1/1	1.00	0.15	10.06	31,31,31,31	0
4	CA	A	504	1/1	0.99	0.21	9.40	45,45,45,45	0
5	NAG	A	511	14/15	0.67	0.24	7.39	65,101,148,164	0
5	NAG	A	512	14/15	0.94	0.14	3.45	37,51,70,76	0
5	NAG	A	516	14/15	0.86	0.34	0.80	68,91,111,113	0
4	CA	B	503	1/1	1.00	0.13	0.73	26,26,26,26	0
5	NAG	A	514	14/15	0.95	0.16	0.06	29,44,60,65	0
4	CA	A	501	1/1	0.99	0.11	0.03	33,33,33,33	0
5	NAG	A	506	14/15	0.95	0.11	0.02	29,44,63,75	0
8	MG	B	501	1/1	0.98	0.10	-0.20	26,26,26,26	0
5	NAG	A	505	14/15	0.97	0.06	-1.93	27,36,40,46	0
4	CA	B	502	1/1	0.98	0.04	-2.48	43,43,43,43	0
7	MAN	A	509	11/12	0.84	0.38	-	75,101,128,128	0
5	NAG	A	515	14/15	0.88	0.39	-	57,84,107,127	0
5	NAG	B	508	14/15	0.58	0.44	-	116,138,163,165	0
5	NAG	B	507	14/15	0.68	0.49	-	94,117,139,140	0
6	BMA	A	507	11/12	0.85	0.19	-	62,75,91,93	0
5	NAG	B	506	14/15	0.93	0.36	-	53,78,93,94	0
7	MAN	A	510	11/12	0.73	0.25	-	71,96,112,119	0
5	NAG	A	513	14/15	0.92	0.30	-	67,89,113,117	0
5	NAG	B	504	14/15	0.88	0.11	-	55,77,95,97	0
5	NAG	B	505	14/15	0.67	0.35	-	91,121,150,158	0
5	NAG	B	509	14/15	0.68	0.49	-	97,140,168,173	0
7	MAN	A	508	11/12	0.76	0.33	-	94,114,134,143	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.