



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:21 am GMT

PDB ID : 1WL4  
Title : Human cytosolic acetoacetyl-CoA thiolase complexed with CoA  
Authors : Kursula, P.; Fukao, T.; Kondo, N.; Wierenga, R.K.  
Deposited on : 2004-06-20  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	<div> <div>5%</div> <div>88%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	3001	-	-	-	X
3	COA	A	1001	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	2001	-	-	-	X
4	GOL	A	2002	-	-	-	X
4	GOL	A	2003	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3577 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called acetyl-Coenzyme A acetyltransferase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	394	2911	1824	509	558	20	0	13	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	92	CSO	CYS	MODIFIED RESIDUE	UNP Q9BWD1
A	169	LYS	THR	CONFLICT	UNP Q9BWD1
A	262	VAL	ALA	CONFLICT	UNP Q9BWD1

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is COENZYME A (three-letter code: COA) (formula: C<sub>21</sub>H<sub>36</sub>N<sub>7</sub>O<sub>16</sub>P<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	585	Total	O	0	0
			585	585		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: acetyl-Coenzyme A acetyltransferase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.86Å 109.83Å 155.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.55 28.75 – 1.55	Depositor EDS
% Data completeness (in resolution range)	96.9 (10.00-1.55) 96.9 (28.75-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.10 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.2.0000	Depositor
R, $R_{free}$	0.106 , 0.140 0.124 , 0.155	Depositor DCC
$R_{free}$ test set	2199 reflections (3.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.8	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 60.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	3577	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, GOL, CSO, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.91	7/3006 (0.2%)	0.88	3/4070 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	371	GLU	CD-OE2	8.53	1.35	1.25
1	A	50	GLU	CD-OE1	6.59	1.32	1.25
1	A	289	GLU	CD-OE1	5.86	1.32	1.25
1	A	302	LYS	CD-CE	-5.50	1.37	1.51
1	A	169	LYS	CD-CE	5.37	1.64	1.51
1	A	396[A]	ARG	CB-CG	-5.29	1.38	1.52
1	A	396[B]	ARG	CB-CG	-5.29	1.38	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	377	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	A	199	ASP	CB-CG-OD2	5.55	123.30	118.30
1	A	223	ARG	NE-CZ-NH2	-5.04	117.78	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	159	HIS	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2911	0	2965	30	0
2	A	15	0	0	0	0
3	A	48	0	32	0	0
4	A	18	0	24	3	0
5	A	585	0	0	15	1
All	All	3577	0	3021	31	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:CSO:SG	1:A:92:CSO:OD	1.94	1.25
1:A:130[B]:TYR:CD1	5:A:3321:HOH:O	2.03	1.12
1:A:130[B]:TYR:CE1	5:A:3321:HOH:O	2.21	0.91
1:A:130[B]:TYR:HD1	5:A:3321:HOH:O	1.49	0.74
1:A:209:THR:HB	1:A:212:GLY:O	1.95	0.67
1:A:170:LYS:NZ	4:A:2003:GOL:H11	2.12	0.65
1:A:129:ALA:HB1	1:A:140:MET:HE3	1.80	0.63
1:A:317[A]:ILE:HD12	1:A:370:LEU:HD23	1.88	0.55
1:A:266:LYS:HE2	5:A:3268:HOH:O	2.06	0.54
1:A:266:LYS:CE	5:A:3268:HOH:O	2.59	0.51
1:A:317[A]:ILE:HD13	1:A:369:THR:HG22	1.92	0.50
1:A:170:LYS:HZ1	4:A:2003:GOL:H11	1.77	0.49
1:A:170:LYS:HZ2	4:A:2003:GOL:H11	1.77	0.49
1:A:159:HIS:CD2	1:A:238:PHE:HE1	2.30	0.48
1:A:307:LYS:NZ	5:A:3376:HOH:O	2.46	0.48
1:A:31[B]:GLN:NE2	5:A:3479:HOH:O	2.46	0.48
1:A:344:ILE:HD11	1:A:373:MET:SD	2.56	0.46
1:A:140:MET:HB2	5:A:3363:HOH:O	2.15	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:LYS:N	5:A:3437:HOH:O	2.47	0.46
1:A:139:GLU:OE2	5:A:3214:HOH:O	2.21	0.46
1:A:151:LEU:HB3	1:A:159:HIS:CE1	2.51	0.46
1:A:159:HIS:CD2	1:A:238:PHE:CE1	3.05	0.45
1:A:59:HIS:HE1	5:A:3004:HOH:O	1.99	0.44
1:A:140:MET:HB2	5:A:3501:HOH:O	2.18	0.43
1:A:305:VAL:HG13	1:A:310:TRP:HB2	2.01	0.43
1:A:190:GLU:O	1:A:194:LYS:HG2	2.19	0.43
1:A:135:VAL:HG13	1:A:135:VAL:O	2.20	0.42
1:A:59:HIS:HD2	5:A:3028:HOH:O	2.03	0.42
1:A:221:PHE:N	1:A:222:PRO:CD	2.83	0.41
1:A:375:ARG:NH2	5:A:3102:HOH:O	2.40	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:3202:HOH:O	5:A:3529:HOH:O[1_655]	1.89	0.31

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/397 (102%)	391 (97%)	12 (3%)	1 (0%)	51	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	318/307 (104%)	311 (98%)	7 (2%)	57 25

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	137	ILE
1	A	169	LYS
1	A	210	ARG
1	A	217	LYS
1	A	293	MET
1	A	295	ILE
1	A	397	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	HIS
1	A	187	ASN
1	A	197	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSO	A	92	1	4,6,7	1.82	1 (25%)	1,6,8	1.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	92	1	-	0/1/5/7	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	CSO	CA-C	3.57	1.54	1.50

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	92	CSO	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	COA	A	1001	-	43,50,50	1.40	5 (11%)	48,75,75	1.85	9 (18%)
4	GOL	A	2001	-	5,5,5	0.52	0	5,5,5	0.49	0
4	GOL	A	2002	-	5,5,5	1.04	0	5,5,5	1.52	1 (20%)
4	GOL	A	2003	-	5,5,5	0.62	0	5,5,5	1.33	0
2	SO4	A	3001	-	4,4,4	0.43	0	6,6,6	0.75	0
2	SO4	A	3002	-	4,4,4	0.53	0	6,6,6	0.43	0
2	SO4	A	3003	-	4,4,4	0.45	0	6,6,6	0.94	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	COA	A	1001	-	-	0/44/64/64	0/3/3/3
4	GOL	A	2001	-	-	0/4/4/4	0/0/0/0
4	GOL	A	2002	-	-	0/4/4/4	0/0/0/0
4	GOL	A	2003	-	-	0/4/4/4	0/0/0/0
2	SO4	A	3001	-	-	0/0/0/0	0/0/0/0
2	SO4	A	3002	-	-	0/0/0/0	0/0/0/0
2	SO4	A	3003	-	-	0/0/0/0	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1001	COA	P1A-O1A	2.03	1.58	1.50
3	A	1001	COA	C2A-N1A	2.27	1.38	1.33
3	A	1001	COA	P3B-O3B	3.03	1.64	1.59
3	A	1001	COA	C2A-N3A	3.29	1.37	1.32
3	A	1001	COA	O9P-C9P	5.31	1.33	1.23

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1001	COA	N3A-C2A-N1A	-7.28	122.52	128.86
3	A	1001	COA	O6A-CCP-CBP	-5.75	101.30	110.55
3	A	1001	COA	C4A-C5A-N7A	-3.15	106.36	109.41
3	A	1001	COA	C4B-O4B-C1B	-3.12	106.45	109.77

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2002	GOL	C3-C2-C1	-2.83	100.26	111.52
3	A	1001	COA	C3P-N4P-C5P	-2.46	118.13	122.84
3	A	1001	COA	C2P-C3P-N4P	-2.40	107.28	112.50
3	A	1001	COA	C5A-C6A-N6A	2.08	124.72	120.47
3	A	1001	COA	O5A-P2A-O4A	2.18	123.58	112.28
3	A	1001	COA	CAP-C9P-N8P	2.33	121.44	116.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2003	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	393/397 (98%)	-0.18	19 (4%)	31 37	8, 13, 31, 68	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	210	ARG	8.7
1	A	4	GLY	7.2
1	A	211	LYS	6.4
1	A	209	THR	6.3
1	A	213	LEU	5.8
1	A	212	GLY	5.2
1	A	135	VAL	4.5
1	A	5	SER	4.5
1	A	208[A]	SER	4.4
1	A	240	THR	4.0
1	A	214	ILE	3.8
1	A	235	LYS	2.9
1	A	241	ASP	2.9
1	A	206	LEU	2.8
1	A	133	THR	2.6
1	A	397	GLU	2.4
1	A	229	GLU	2.2
1	A	242	GLY	2.1
1	A	243	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSO	A	92	7/8	0.94	0.08	-	11,13,22,27	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	A	2001	6/6	0.91	0.22	8.64	29,32,34,35	0
2	SO4	A	3001	5/5	0.99	0.10	3.38	16,16,17,18	0
4	GOL	A	2003	6/6	0.76	0.19	2.92	37,46,47,50	0
4	GOL	A	2002	6/6	0.90	0.17	2.88	27,32,38,40	0
3	COA	A	1001	48/48	0.87	0.12	2.82	14,20,56,59	0
2	SO4	A	3003	5/5	0.96	0.18	-	36,37,38,41	0
2	SO4	A	3002	5/5	0.99	0.12	-	19,21,26,27	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.