



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 03:32 am GMT

PDB ID : 1WNY  
Title : Isoleucyl-tRNA synthetase editing domain  
Authors : Fukunaga, R.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2004-08-11  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	186	
1	B	186	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called isoleucyl-trna synthetase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	180	Total	C	N	O	0	0	0
			1407	919	228	260			
1	B	180	Total	C	N	O	0	0	0
			1407	919	228	260			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	200	MET	-	INITIATING METHIONINE	UNP P56690
A	274	PRO	GLN	CONFLICT	UNP P56690
B	200	MET	-	INITIATING METHIONINE	UNP P56690
B	274	PRO	GLN	CONFLICT	UNP P56690

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	284	Total	O	0	0
			284	284		
2	B	236	Total	O	0	0
			236	236		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: isoleucyl-trna synthetase



- Molecule 1: isoleucyl-trna synthetase



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.72Å 102.72Å 82.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.25 – 1.60	Depositor
% Data completeness (in resolution range)	99.7 (30.25-1.60)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.176 , 0.209	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	3334	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.78	0/1444	0.85	1/1965 (0.1%)
1	B	0.75	0/1444	0.87	0/1965
All	All	0.77	0/2888	0.86	1/3930 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	373	ARG	NE-CZ-NH1	-5.74	117.43	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1407	0	1420	23	3
1	B	1407	0	1420	33	0
2	A	284	0	0	5	5
2	B	236	0	0	10	2
All	All	3334	0	2840	56	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:243:HIS:HD2	1:B:245:GLU:H	1.20	0.89
1:A:243:HIS:HD2	1:A:245:GLU:H	1.22	0.86
1:B:215:LYS:NZ	1:B:215:LYS:HB2	1.97	0.78
1:B:378:LYS:HE3	2:B:496:HOH:O	1.85	0.75
1:A:202:ASP:HA	1:A:203:PRO:C	2.08	0.74
1:B:333:ARG:HE	1:B:333:ARG:HA	1.53	0.73
1:B:213:GLU:HG3	2:B:599:HOH:O	1.95	0.67
1:B:243:HIS:CD2	1:B:245:GLU:H	2.10	0.65
1:B:215:LYS:HZ2	1:B:215:LYS:HB2	1.60	0.65
1:B:367:LEU:HB3	2:B:580:HOH:O	1.97	0.64
1:A:298:GLU:HG3	2:A:429:HOH:O	1.98	0.63
1:A:202:ASP:HA	1:A:204:SER:N	2.16	0.61
1:B:360:ARG:NH1	2:B:407:HOH:O	2.24	0.58
1:B:202:ASP:N	1:B:203:PRO:CD	2.67	0.57
1:B:252:GLN:NE2	1:B:276:LEU:HD11	2.19	0.57
1:A:243:HIS:CD2	1:A:245:GLU:H	2.13	0.57
1:B:252:GLN:HE21	1:B:276:LEU:HD11	1.69	0.57
1:B:202:ASP:N	1:B:203:PRO:HD2	2.22	0.55
1:B:204:SER:OG	1:B:380:GLU:HB3	2.07	0.54
1:A:231:PRO:HD2	1:A:363:ASN:HD21	1.73	0.54
1:A:267:LYS:NZ	1:A:267:LYS:HB3	2.23	0.54
1:A:333:ARG:HD3	2:A:627:HOH:O	2.09	0.53
1:B:339:LEU:HB2	2:B:514:HOH:O	2.09	0.51
1:A:202:ASP:OD1	1:A:380:GLU:O	2.29	0.51
1:A:213:GLU:OE1	1:A:216:LYS:HE2	2.12	0.50
1:B:339:LEU:C	1:B:339:LEU:HD23	2.32	0.50
1:B:233:THR:OG1	1:B:319:HIS:HE1	1.94	0.49
1:B:348:LYS:HE2	1:B:356:GLY:O	2.13	0.49
1:A:226:ILE:C	1:A:226:ILE:HD12	2.34	0.48
1:B:212:LYS:HE3	2:B:593:HOH:O	2.11	0.48
1:B:255:ASP:HB2	2:B:592:HOH:O	2.13	0.48
1:B:243:HIS:HB3	1:B:246:TYR:CD2	2.49	0.48
1:A:215:LYS:HG3	2:A:436:HOH:O	2.14	0.47
1:B:246:TYR:CZ	1:B:311:GLN:HB3	2.49	0.47
1:B:308:TYR:CD2	1:B:308:TYR:C	2.88	0.47
1:B:213:GLU:CD	2:B:516:HOH:O	2.54	0.46
1:B:215:LYS:NZ	1:B:215:LYS:CB	2.74	0.46
1:A:246:TYR:CE2	1:A:311:GLN:HB3	2.52	0.45
1:A:202:ASP:OD1	1:A:381:SER:HA	2.17	0.45
1:B:345:GLU:HG2	1:B:345:GLU:H	1.51	0.45
1:B:213:GLU:OE2	1:B:213:GLU:HA	2.17	0.45
1:B:226:ILE:C	1:B:226:ILE:HD12	2.38	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:LYS:HD2	1:A:305:LEU:HD11	2.00	0.43
1:A:243:HIS:HB3	1:A:246:TYR:CD2	2.54	0.43
1:A:243:HIS:HE1	2:A:649:HOH:O	2.02	0.43
1:A:205:VAL:HG11	1:A:367:LEU:HD21	2.01	0.42
1:A:282:LYS:HD3	2:A:645:HOH:O	2.19	0.42
1:B:213:GLU:HB2	2:B:547:HOH:O	2.19	0.42
1:A:266:ARG:HE	1:A:271:GLU:CD	2.24	0.41
1:B:243:HIS:HD2	1:B:245:GLU:N	2.02	0.41
1:B:237:ASN:OD1	1:B:319:HIS:HD2	2.03	0.41
1:B:308:TYR:CE1	1:B:327:GLU:HB3	2.57	0.40
1:A:205:VAL:HG11	1:A:367:LEU:CD2	2.51	0.40
1:A:308:TYR:OH	1:A:320:GLN:NE2	2.54	0.40
1:A:232:TRP:H	1:A:363:ASN:HD22	1.68	0.40
1:B:341:LYS:HE3	2:B:499:HOH:O	2.21	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:391:HOH:O	2:B:477:HOH:O[5_547]	1.73	0.47
2:A:391:HOH:O	2:B:504:HOH:O[5_547]	1.80	0.40
1:A:368:ARG:NH1	2:A:435:HOH:O[3_545]	1.98	0.22
1:A:368:ARG:NH1	2:A:618:HOH:O[3_545]	2.02	0.18
1:A:368:ARG:NH2	2:A:618:HOH:O[3_545]	2.13	0.07

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	178/186 (96%)	175 (98%)	3 (2%)	0	100	100
1	B	178/186 (96%)	175 (98%)	3 (2%)	0	100	100

Continued on next page...



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	356/372 (96%)	350 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/152 (96%)	140 (96%)	6 (4%)	35	11
1	B	146/152 (96%)	135 (92%)	11 (8%)	16	3
All	All	292/304 (96%)	275 (94%)	17 (6%)	23	5

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	279	PHE
1	A	319	HIS
1	A	328	ASP
1	A	353	PRO
1	A	375	LEU
1	A	380	GLU
1	B	215	LYS
1	B	279	PHE
1	B	308	TYR
1	B	312	GLU
1	B	319	HIS
1	B	328	ASP
1	B	330	GLU
1	B	333	ARG
1	B	353	PRO
1	B	360	ARG
1	B	375	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	237	ASN
1	A	243	HIS
1	A	295	GLN
1	A	320	GLN
1	A	363	ASN
1	B	237	ASN
1	B	243	HIS
1	B	252	GLN
1	B	295	GLN
1	B	311	GLN
1	B	319	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.