



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 11:13 pm GMT

PDB ID : 3WNU
Title : The crystal structure of catalase-peroxidase, KatG, from *Synechococcus* PCC7942
Authors : Tada, T.; Wada, K.; Kamachi, S.
Deposited on : 2013-12-17
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4002 (2.20-2.20)
Clashscore	112137	4730 (2.20-2.20)
Ramachandran outliers	110173	4656 (2.20-2.20)
Sidechain outliers	110143	4657 (2.20-2.20)
RSRZ outliers	101464	4033 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	720	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>...</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6149 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Catalase-peroxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	710	5582	3523	980	1060	19	0	0	0

- Molecule 2 is HEME B/C (three-letter code: HEB) (formula: $C_{34}H_{34}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	522	Total 522	O 522	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Catalase-peroxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	108.74Å 108.74Å 202.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.39 – 2.20 34.39 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.39-2.20) 99.9 (34.39-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.168 , 0.206 0.173 , 0.208	Depositor DCC
R_{free} test set	3159 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	34.3	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6149	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, HEB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	6/5729 (0.1%)	1.12	39/7803 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	327	ARG	NE-CZ	9.67	1.45	1.33
1	A	287	GLU	CD-OE1	7.77	1.34	1.25
1	A	327	ARG	CZ-NH2	6.15	1.41	1.33
1	A	321	TRP	CB-CG	-5.92	1.39	1.50
1	A	76	ASP	CB-CG	5.77	1.63	1.51
1	A	327	ARG	CD-NE	5.64	1.56	1.46

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	327	ARG	NE-CZ-NH2	17.98	129.29	120.30
1	A	576	ARG	NE-CZ-NH2	-17.08	111.76	120.30
1	A	242	ARG	NE-CZ-NH1	15.87	128.24	120.30
1	A	242	ARG	NE-CZ-NH2	-14.01	113.30	120.30
1	A	576	ARG	NE-CZ-NH1	12.35	126.48	120.30
1	A	327	ARG	CG-CD-NE	10.33	133.50	111.80
1	A	327	ARG	NH1-CZ-NH2	-9.74	108.69	119.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	502	ARG	NE-CZ-NH1	9.28	124.94	120.30
1	A	502	ARG	NE-CZ-NH2	-9.16	115.72	120.30
1	A	414	MET	CG-SD-CE	8.65	114.04	100.20
1	A	370	VAL	CB-CA-C	-8.26	95.71	111.40
1	A	567	VAL	CG1-CB-CG2	7.40	122.73	110.90
1	A	124	ASP	CB-CG-OD2	7.31	124.88	118.30
1	A	385	ARG	NE-CZ-NH2	-7.11	116.74	120.30
1	A	628	THR	CB-CA-C	-7.04	92.59	111.60
1	A	412	ARG	NE-CZ-NH1	7.04	123.82	120.30
1	A	99	THR	N-CA-CB	-7.02	96.97	110.30
1	A	643	THR	N-CA-CB	-6.89	97.20	110.30
1	A	608	LEU	CB-CG-CD1	6.48	122.02	111.00
1	A	165	LEU	CA-CB-CG	6.46	130.15	115.30
1	A	310	LEU	CB-CG-CD2	6.38	121.84	111.00
1	A	538	ARG	NE-CZ-NH1	6.35	123.48	120.30
1	A	353	GLU	CB-CA-C	-6.29	97.82	110.40
1	A	242	ARG	CD-NE-CZ	5.90	131.86	123.60
1	A	424	ASP	CB-CG-OD2	5.84	123.56	118.30
1	A	310	LEU	CA-CB-CG	5.80	128.64	115.30
1	A	576	ARG	CD-NE-CZ	5.72	131.60	123.60
1	A	115	ARG	NE-CZ-NH2	5.69	123.15	120.30
1	A	165	LEU	CB-CG-CD2	5.62	120.56	111.00
1	A	366	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	285	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	150	ASP	CB-CG-OD1	5.45	123.20	118.30
1	A	412	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	A	429	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	370	VAL	CG1-CB-CG2	5.26	119.31	110.90
1	A	48	ASP	CB-CG-OD2	-5.22	113.61	118.30
1	A	628	THR	N-CA-CB	5.15	120.09	110.30
1	A	288	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	488	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	210	GLU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5582	0	5365	76	0
2	A	43	0	32	3	0
3	A	2	0	0	0	0
4	A	522	0	0	11	1
All	All	6149	0	5397	79	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:TRP:CH2	1:A:222:TYR:HE2	1.00	1.65
1:A:94:TRP:HH2	1:A:222:TYR:CE2	0.93	1.58
1:A:94:TRP:CH2	1:A:222:TYR:CE2	1.83	1.31
1:A:94:TRP:HH2	1:A:222:TYR:CD2	1.83	0.95
1:A:99:THR:HG22	1:A:108:GLY:H	1.36	0.88
1:A:94:TRP:CZ3	1:A:222:TYR:HE2	1.88	0.88
1:A:591:LEU:O	1:A:595:THR:HG23	1.77	0.84
1:A:94:TRP:CH2	1:A:222:TYR:CD2	2.64	0.81
1:A:106:ARG:HH21	1:A:596:GLN:HE21	1.32	0.78
1:A:114:GLN:HE21	1:A:132:ARG:HH11	1.28	0.76
1:A:94:TRP:CZ3	1:A:222:TYR:CE2	2.70	0.73
1:A:99:THR:CG2	1:A:108:GLY:H	2.02	0.71
1:A:38:HIS:H	1:A:179:HIS:HE1	1.42	0.67
1:A:179:HIS:HD2	1:A:180:PRO:O	1.78	0.66
1:A:198:ASN:OD1	1:A:199:PRO:O	2.14	0.66
1:A:171:ALA:H	1:A:411:HIS:HE1	1.42	0.65
1:A:251:ASN:C	1:A:251:ASN:HD22	2.00	0.65
1:A:628:THR:HB	1:A:630:ARG:H	1.66	0.60
1:A:492:GLN:HG2	1:A:495:TRP:CZ3	2.36	0.60
1:A:251:ASN:ND2	1:A:254:GLU:H	2.00	0.59
1:A:639:PHE:O	1:A:643:THR:HB	2.02	0.59
1:A:114:GLN:HE21	1:A:132:ARG:NH1	2.00	0.58
1:A:242:ARG:HD3	1:A:252:ASP:OD1	2.03	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:TRP:HZ3	1:A:222:TYR:HH	1.49	0.57
1:A:106:ARG:HH21	1:A:596:GLN:NE2	2.00	0.56
1:A:502:ARG:HD3	4:A:1327:HOH:O	2.06	0.55
1:A:628:THR:HG21	4:A:1254:HOH:O	2.06	0.55
1:A:99:THR:HG21	4:A:1053:HOH:O	2.05	0.55
1:A:285:ASP:CB	4:A:1344:HOH:O	2.54	0.54
2:A:801:HEB:HHC	2:A:801:HEB:HBB3	1.90	0.54
1:A:595:THR:HG21	1:A:605:MET:SD	2.47	0.54
1:A:440:ASN:ND2	1:A:442:ASN:H	2.06	0.53
1:A:224:ASN:HD22	1:A:224:ASN:C	2.12	0.53
1:A:38:HIS:H	1:A:179:HIS:CE1	2.25	0.53
1:A:445:VAL:HG13	1:A:527:LEU:HD21	1.91	0.53
1:A:591:LEU:O	1:A:595:THR:CG2	2.55	0.53
1:A:171:ALA:H	1:A:411:HIS:CE1	2.26	0.52
1:A:366:ARG:NH1	4:A:1199:HOH:O	2.43	0.51
1:A:648:LEU:HD12	1:A:648:LEU:C	2.31	0.51
1:A:310:LEU:HD22	1:A:343:ALA:HB1	1.93	0.50
1:A:224:ASN:HD22	1:A:226:GLU:H	1.58	0.49
1:A:628:THR:CG2	1:A:637:ASP:OD2	2.60	0.49
1:A:618:ASN:OD1	1:A:622:THR:HG22	2.12	0.49
1:A:113:ASN:HD22	1:A:114:GLN:N	2.11	0.48
1:A:120:ASN:HD22	1:A:121:SER:N	2.12	0.47
1:A:385:ARG:NH2	1:A:389:GLU:OE1	2.38	0.47
1:A:507:LEU:O	1:A:511:GLU:HB2	2.15	0.47
1:A:224:ASN:ND2	1:A:226:GLU:H	2.14	0.46
1:A:618:ASN:ND2	4:A:1046:HOH:O	2.48	0.46
1:A:655:ASN:ND2	1:A:673:ARG:H	2.14	0.46
1:A:561:ASP:OD1	1:A:561:ASP:C	2.54	0.46
1:A:217:THR:HG22	1:A:220:LEU:HD12	1.97	0.45
1:A:611:GLY:HA3	1:A:704:PHE:CE1	2.51	0.45
1:A:514:ALA:HB1	1:A:519:ALA:O	2.17	0.45
1:A:68:GLN:HE21	1:A:68:GLN:HB2	1.62	0.45
1:A:179:HIS:HB2	1:A:180:PRO:HD2	1.99	0.44
1:A:48:ASP:N	4:A:1421:HOH:O	2.16	0.44
1:A:238:ALA:HB2	1:A:377:ALA:HB1	2.00	0.44
1:A:492:GLN:HG2	1:A:495:TRP:CH2	2.52	0.44
2:A:801:HEB:HMB1	2:A:801:HEB:HAB2	1.78	0.44
1:A:127:ASN:HA	1:A:129:ASP:OD1	2.18	0.43
1:A:94:TRP:CZ2	1:A:222:TYR:CD2	3.07	0.43
1:A:25:TRP:HB2	1:A:26:TRP:CE3	2.53	0.43
1:A:317:HIS:HE1	4:A:998:HOH:O	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:285:ASP:HB3	4:A:1344:HOH:O	2.16	0.42
1:A:412:ARG:NH1	4:A:1387:HOH:O	2.44	0.42
1:A:92:LEU:HD23	1:A:114:GLN:HE22	1.84	0.42
1:A:312:GLY:C	1:A:370:VAL:HG22	2.39	0.42
1:A:622:THR:HG23	1:A:624:HIS:H	1.84	0.42
1:A:628:THR:HG22	1:A:637:ASP:OD2	2.20	0.41
1:A:602:ALA:HB3	1:A:603:PRO:HD3	2.01	0.41
1:A:224:ASN:HD22	1:A:225:PRO:N	2.19	0.41
1:A:439:GLY:HA3	1:A:552:ARG:O	2.20	0.41
1:A:502:ARG:CD	4:A:1327:HOH:O	2.67	0.41
1:A:488:ARG:O	1:A:493:LYS:HD3	2.21	0.41
1:A:628:THR:HG23	1:A:637:ASP:OD2	2.21	0.41
2:A:801:HEB:HMC1	2:A:801:HEB:HBC1	2.03	0.41
1:A:37:GLN:NE2	1:A:182:LYS:H	2.18	0.41
1:A:251:ASN:HD21	1:A:254:GLU:H	1.68	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1332:HOH:O	4:A:1334:HOH:O[7_554]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	708/720 (98%)	687 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	573/581 (99%)	528 (92%)	45 (8%)	14	14

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	THR
1	A	22	THR
1	A	24	GLU
1	A	28	LYS
1	A	32	LEU
1	A	40	ARG
1	A	53	GLU
1	A	65	GLN
1	A	67	LEU
1	A	68	GLN
1	A	92	LEU
1	A	99	THR
1	A	113	ASN
1	A	120	ASN
1	A	124	ASP
1	A	154	TYR
1	A	165	LEU
1	A	204	THR
1	A	206	ASP
1	A	207	ARG
1	A	224	ASN
1	A	251	ASN
1	A	298	GLN
1	A	310	LEU
1	A	327	ARG
1	A	370	VAL
1	A	373	ASP
1	A	427	GLN
1	A	442	ASN
1	A	511	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	527	LEU
1	A	544	ILE
1	A	563	GLU
1	A	567	VAL
1	A	582	ASP
1	A	595	THR
1	A	596	GLN
1	A	608	LEU
1	A	622	THR
1	A	628	THR
1	A	643	THR
1	A	656	LEU
1	A	663	LYS
1	A	719	LEU
1	A	720	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	37	GLN
1	A	68	GLN
1	A	113	ASN
1	A	114	GLN
1	A	120	ASN
1	A	125	ASN
1	A	179	HIS
1	A	224	ASN
1	A	251	ASN
1	A	317	HIS
1	A	411	HIS
1	A	440	ASN
1	A	535	GLN
1	A	572	HIS
1	A	596	GLN
1	A	618	ASN
1	A	655	ASN
1	A	666	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEB	A	801	1	31,50,50	5.61	21 (67%)	20,82,82	4.68	13 (65%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEB	A	801	1	-	0/8/94/94	0/0/8/8

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEB	C4B-NB	-13.41	1.31	1.49
2	A	801	HEB	C1D-ND	-11.41	1.34	1.49
2	A	801	HEB	C1B-NB	-11.07	1.34	1.49
2	A	801	HEB	CHD-C1D	-8.23	1.37	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEB	C4D-ND	-8.16	1.38	1.49
2	A	801	HEB	CHA-C4D	-8.06	1.37	1.53
2	A	801	HEB	CHB-C1B	-7.45	1.39	1.53
2	A	801	HEB	CHC-C4B	-7.31	1.39	1.53
2	A	801	HEB	CHA-C1A	-6.46	1.35	1.51
2	A	801	HEB	CHD-C4C	-5.53	1.38	1.51
2	A	801	HEB	CHC-C1C	-5.18	1.39	1.51
2	A	801	HEB	CHB-C4A	-4.73	1.40	1.51
2	A	801	HEB	C3C-C2C	-4.70	1.34	1.40
2	A	801	HEB	CBB-CAB	-4.06	1.32	1.51
2	A	801	HEB	CAB-C3B	-3.47	1.44	1.51
2	A	801	HEB	CMA-C3A	-2.74	1.46	1.51
2	A	801	HEB	CMC-C2C	-2.25	1.47	1.51
2	A	801	HEB	C4A-C3A	2.72	1.41	1.38
2	A	801	HEB	CMB-C2B	2.80	1.55	1.50
2	A	801	HEB	C1A-C2A	3.25	1.42	1.38
2	A	801	HEB	C1C-C2C	4.23	1.44	1.38

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	HEB	CAA-C2A-C1A	-3.82	124.61	127.30
2	A	801	HEB	CHD-C4C-C3C	-3.32	125.09	129.55
2	A	801	HEB	CAA-CBA-CGA	-3.09	107.37	112.66
2	A	801	HEB	CHB-C4A-C3A	-2.52	125.09	129.45
2	A	801	HEB	CHC-C1C-C2C	-2.03	125.94	129.45
2	A	801	HEB	CMC-C2C-C3C	2.65	129.81	124.89
2	A	801	HEB	CAD-C3D-C4D	3.13	128.91	122.52
2	A	801	HEB	CBD-CAD-C3D	4.51	122.53	114.28
2	A	801	HEB	CBB-CAB-C3B	5.95	121.98	112.95
2	A	801	HEB	CHD-C1D-ND	7.24	124.41	110.75
2	A	801	HEB	CHA-C4D-ND	8.35	126.50	110.75
2	A	801	HEB	CHB-C1B-NB	9.49	128.65	110.75
2	A	801	HEB	CHC-C4B-NB	9.91	129.44	110.75

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	HEB	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	710/720 (98%)	-0.38	19 (2%) 55 52	23, 33, 57, 97	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	199	PRO	4.8
1	A	11	MET	4.7
1	A	206	ASP	3.7
1	A	23	ALA	3.4
1	A	19	ASN	3.0
1	A	204	THR	3.0
1	A	211	ASN	3.0
1	A	200	ASN	2.9
1	A	197	THR	2.8
1	A	18	VAL	2.8
1	A	48	ASP	2.6
1	A	541	GLY	2.5
1	A	20	ILE	2.3
1	A	442	ASN	2.2
1	A	454	ALA	2.2
1	A	720	ASP	2.2
1	A	584	ALA	2.1
1	A	92	LEU	2.1
1	A	94	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	HEB	A	801	43/43	0.98	0.20	1.15	22,26,29,33	0
3	NA	A	802	1/1	0.99	0.09	-1.57	27,27,27,27	0
3	NA	A	803	1/1	0.99	0.04	-2.02	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.