



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2017 – 05:40 AM EDT

PDB ID : 2WOV
Title : Trypanosoma brucei trypanothione reductase with bound NADP.
Authors : Alphey, M.S.; Fairlamb, A.H.
Deposited on : 2009-07-30
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3846 (2.50-2.50)
Clashscore	112137	4554 (2.50-2.50)
Ramachandran outliers	110173	4463 (2.50-2.50)
Sidechain outliers	110143	4465 (2.50-2.50)
RSRZ outliers	101464	3876 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 15%, green 83%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 83% 15% .. </div> </div>
1	B	495	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 16%, green 81%, grey 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 81% 16% .. </div> </div>
1	C	495	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 17%, green 80%, orange 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 80% 17% .. </div> </div>
1	D	495	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 16%, green 80%, grey 4%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 100% 80% 16% .. </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NA	D	1489	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 15706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRYPANOTHIONE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	0	2	0
			3735	2375	637	703	20			
1	B	486	Total	C	N	O	S	0	0	0
			3691	2348	627	697	19			
1	C	490	Total	C	N	O	S	0	2	0
			3733	2374	634	705	20			
1	D	487	Total	C	N	O	S	0	0	0
			3696	2351	628	698	19			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q389T8
A	-1	SER	-	expression tag	UNP Q389T8
A	0	HIS	-	expression tag	UNP Q389T8
B	-2	GLY	-	expression tag	UNP Q389T8
B	-1	SER	-	expression tag	UNP Q389T8
B	0	HIS	-	expression tag	UNP Q389T8
C	-2	GLY	-	expression tag	UNP Q389T8
C	-1	SER	-	expression tag	UNP Q389T8
C	0	HIS	-	expression tag	UNP Q389T8
D	-2	GLY	-	expression tag	UNP Q389T8
D	-1	SER	-	expression tag	UNP Q389T8
D	0	HIS	-	expression tag	UNP Q389T8

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Na	0	0
			2	2		
4	A	1	Total	Na	0	0
			1	1		
4	D	2	Total	Na	0	0
			2	2		
4	C	2	Total	Na	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	121	Total	O	0	0
			121	121		
5	B	99	Total	O	0	0
			99	99		
5	C	128	Total	O	0	0
			128	128		
5	D	92	Total	O	0	0
			92	92		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TRYPANOTHIONE REDUCTASE



• Molecule 1: TRYPANOTHIONE REDUCTASE




• Molecule 1: TRYPANOTHIONE REDUCTASE





● Molecule 1: TRYPANOTHIONE REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	100.87Å 63.45Å 169.35Å 90.00° 97.85° 90.00°	Depositor
Resolution (Å)	19.85 – 2.50 19.85 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.85-2.50) 98.1 (19.85-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.50Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.179 , 0.251 0.179 , 0.250	Depositor DCC
R_{free} test set	3628 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15706	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, FAD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/3820	0.80	1/5181 (0.0%)
1	B	0.75	0/3768	0.78	2/5111 (0.0%)
1	C	0.83	0/3818	0.82	2/5179 (0.0%)
1	D	0.81	0/3773	0.80	4/5118 (0.1%)
All	All	0.80	0/15179	0.80	9/20589 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	222	ARG	NE-CZ-NH2	-7.32	116.64	120.30
1	A	290	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	C	121	ASP	CB-CG-OD2	-6.14	112.77	118.30
1	D	260	LEU	CB-CG-CD2	6.14	121.45	111.00
1	B	260	LEU	CB-CG-CD1	-5.95	100.88	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	214	GLY	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	C	352	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3735	0	3753	54	0
1	B	3691	0	3703	55	0
1	C	3733	0	3746	56	0
1	D	3696	0	3704	44	0
2	A	48	0	26	1	0
2	B	48	0	26	0	0
2	C	48	0	26	0	0
2	D	48	0	26	1	0
3	A	53	0	31	1	0
3	B	53	0	31	2	0
3	C	53	0	31	2	0
3	D	53	0	31	1	0
4	A	1	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
4	D	2	0	0	0	0
5	A	121	0	0	1	0
5	B	99	0	0	4	0
5	C	128	0	0	2	0
5	D	92	0	0	2	0
All	All	15706	0	15134	196	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:486:LYS:O	1:B:487:LEU:HB2	1.54	1.06
1:A:292[B]:ASN:H	1:A:292[B]:ASN:ND2	1.61	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:52:CYS:HG	1:B:57:CYS:HG	1.03	0.93
1:A:73:LEU:HD23	1:B:73:LEU:HD23	1.58	0.86
1:A:440:ALA:HB3	1:B:440:ALA:HB3	1.57	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	490/495 (99%)	474 (97%)	14 (3%)	2 (0%)	38	59
1	B	484/495 (98%)	459 (95%)	19 (4%)	6 (1%)	15	27
1	C	490/495 (99%)	470 (96%)	18 (4%)	2 (0%)	38	59
1	D	485/495 (98%)	469 (97%)	12 (2%)	4 (1%)	22	39
All	All	1949/1980 (98%)	1872 (96%)	63 (3%)	14 (1%)	25	43

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	132	LYS
1	D	132	LYS
1	B	320	PRO
1	A	48	LEU
1	B	45	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/407 (100%)	388 (96%)	17 (4%)	34	59
1	B	399/407 (98%)	386 (97%)	13 (3%)	43	70
1	C	405/407 (100%)	382 (94%)	23 (6%)	24	44
1	D	399/407 (98%)	372 (93%)	27 (7%)	18	34
All	All	1608/1628 (99%)	1528 (95%)	80 (5%)	29	51

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	236	GLU
1	C	323	TYR
1	D	306	LYS
1	C	266	LYS
1	C	292[B]	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	359	HIS
1	C	152	GLN
1	D	310	GLN
1	C	68	GLN
1	C	115	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 15 ligands modelled in this entry, 7 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	A	800	-	43,52,52	1.69	11 (25%)	49,80,80	1.61	4 (8%)
3	FAD	A	998	-	51,58,58	1.38	8 (15%)	54,89,89	2.00	9 (16%)
2	NDP	B	800	-	43,52,52	1.54	8 (18%)	49,80,80	1.67	5 (10%)
3	FAD	B	998	-	51,58,58	1.48	6 (11%)	54,89,89	2.08	10 (18%)
2	NDP	C	800	-	43,52,52	1.67	8 (18%)	49,80,80	1.53	6 (12%)
3	FAD	C	998	-	51,58,58	1.62	10 (19%)	54,89,89	1.93	9 (16%)
2	NDP	D	800	-	43,52,52	1.50	6 (13%)	49,80,80	1.80	5 (10%)
3	FAD	D	998	-	51,58,58	1.58	9 (17%)	54,89,89	2.15	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	800	-	-	0/30/77/77	0/5/5/5
3	FAD	A	998	-	-	0/28/50/50	0/6/6/6
2	NDP	B	800	-	-	0/30/77/77	0/5/5/5
3	FAD	B	998	-	-	0/28/50/50	0/6/6/6
2	NDP	C	800	-	-	0/30/77/77	0/5/5/5
3	FAD	C	998	-	-	0/28/50/50	0/6/6/6
2	NDP	D	800	-	-	0/30/77/77	0/5/5/5
3	FAD	D	998	-	-	0/28/50/50	0/6/6/6

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	800	NDP	C4N-C5N	-4.05	1.40	1.49
2	B	800	NDP	C4N-C5N	-3.77	1.40	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	800	NDP	C4N-C5N	-3.74	1.41	1.49
2	D	800	NDP	C4N-C5N	-3.47	1.41	1.49
2	B	800	NDP	P2B-O3X	-2.73	1.43	1.54

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	998	FAD	N3A-C2A-N1A	-11.55	118.80	128.86
3	A	998	FAD	N3A-C2A-N1A	-9.53	120.56	128.86
2	D	800	NDP	N3A-C2A-N1A	-8.62	121.35	128.86
3	B	998	FAD	N3A-C2A-N1A	-8.47	121.48	128.86
2	A	800	NDP	N3A-C2A-N1A	-7.58	122.26	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	800	NDP	1	0
3	A	998	FAD	1	0
3	B	998	FAD	2	0
3	C	998	FAD	2	0
2	D	800	NDP	1	0
3	D	998	FAD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	490/495 (98%)	-0.45	4 (0%) 86 86	13, 24, 41, 55	2 (0%)
1	B	486/495 (98%)	-0.33	5 (1%) 82 83	16, 29, 51, 66	0
1	C	490/495 (98%)	-0.49	2 (0%) 92 92	14, 24, 38, 50	0
1	D	487/495 (98%)	-0.38	6 (1%) 79 80	15, 26, 47, 59	1 (0%)
All	All	1953/1980 (98%)	-0.41	17 (0%) 84 85	13, 25, 47, 66	3 (0%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	SER	5.2
1	B	352	ASN	3.8
1	B	320	PRO	3.3
1	B	354	PRO	3.3
1	A	352	ASN	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NA	D	1489	1/1	0.94	0.21	4.05	35,35,35,35	0
4	NA	D	1490	1/1	0.78	0.18	1.03	58,58,58,58	0
4	NA	A	1489	1/1	0.89	0.13	0.49	40,40,40,40	0
4	NA	B	1488	1/1	0.94	0.12	0.21	32,32,32,32	0
3	FAD	C	998	53/53	0.98	0.10	-0.37	9,16,20,23	0
3	FAD	B	998	53/53	0.97	0.11	-0.41	10,22,30,31	0
2	NDP	C	800	48/48	0.97	0.10	-0.59	15,23,28,29	0
2	NDP	B	800	48/48	0.97	0.10	-0.59	16,26,29,30	0
3	FAD	A	998	53/53	0.98	0.09	-0.68	6,13,19,22	0
2	NDP	D	800	48/48	0.97	0.10	-0.85	20,26,28,29	0
4	NA	C	1489	1/1	0.88	0.09	-0.85	34,34,34,34	0
3	FAD	D	998	53/53	0.98	0.10	-0.87	5,20,25,26	0
2	NDP	A	800	48/48	0.98	0.08	-1.14	15,24,30,33	0
4	NA	C	1490	1/1	0.92	0.06	-2.06	23,23,23,23	0
4	NA	B	1489	1/1	0.92	0.09	-2.15	41,41,41,41	0

6.5 Other polymers

There are no such residues in this entry.