



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 4, 2017 – 05:55 AM EDT

PDB ID : 4WP2
Title : Chaetomium Mex67 UBA domain
Authors : Aibara, S.; Stewart, M.
Deposited on : unknown
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3453 (1.70-1.70)
Clashscore	112137	3876 (1.70-1.70)
Ramachandran outliers	110173	3815 (1.70-1.70)
Sidechain outliers	110143	3815 (1.70-1.70)
RSRZ outliers	101464	3491 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	<div> <div>7%</div> <div>98%</div> <div>•</div> </div>
1	B	61	<div> <div>28%</div> <div>79%</div> <div>16%</div> <div>5%</div> </div>
1	C	61	<div> <div>11%</div> <div>93%</div> <div>7%</div> </div>
1	D	61	<div> <div>31%</div> <div>82%</div> <div>13%</div> <div>5%</div> </div>
1	E	61	<div> <div>11%</div> <div>97%</div> <div>•</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	61	
1	G	61	
1	H	61	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7657 atoms, of which 3623 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative mRNA export protein.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	B	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			
1	C	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	D	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			
1	E	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	F	56	Total	As	C	H	N	O	S	0	0	0
			865	1	272	430	75	84	3			
1	G	61	Total	As	C	H	N	O	S	0	0	0
			934	1	292	463	82	93	3			
1	H	58	Total	As	C	H	N	O	S	0	0	0
			899	1	282	447	78	88	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	SER	-	expression tag	UNP G0SET4
A	598	ASN	-	expression tag	UNP G0SET4
A	599	ALA	-	expression tag	UNP G0SET4
B	597	SER	-	expression tag	UNP G0SET4
B	598	ASN	-	expression tag	UNP G0SET4
B	599	ALA	-	expression tag	UNP G0SET4
C	597	SER	-	expression tag	UNP G0SET4
C	598	ASN	-	expression tag	UNP G0SET4
C	599	ALA	-	expression tag	UNP G0SET4
D	597	SER	-	expression tag	UNP G0SET4
D	598	ASN	-	expression tag	UNP G0SET4
D	599	ALA	-	expression tag	UNP G0SET4
E	597	SER	-	expression tag	UNP G0SET4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	598	ASN	-	expression tag	UNP G0SET4
E	599	ALA	-	expression tag	UNP G0SET4
F	597	SER	-	expression tag	UNP G0SET4
F	598	ASN	-	expression tag	UNP G0SET4
F	599	ALA	-	expression tag	UNP G0SET4
G	597	SER	-	expression tag	UNP G0SET4
G	598	ASN	-	expression tag	UNP G0SET4
G	599	ALA	-	expression tag	UNP G0SET4
H	597	SER	-	expression tag	UNP G0SET4
H	598	ASN	-	expression tag	UNP G0SET4
H	599	ALA	-	expression tag	UNP G0SET4

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	46	Total O 46 46	0	0
2	B	41	Total O 41 41	0	0
2	C	38	Total O 38 38	0	0
2	D	56	Total O 56 56	0	0
2	E	49	Total O 49 49	0	0
2	F	47	Total O 47 47	0	0
2	G	40	Total O 40 40	0	0
2	H	42	Total O 42 42	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative mRNA export protein





- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein




- Molecule 1: Putative mRNA export protein



• Molecule 1: Putative mRNA export protein



• Molecule 1: Putative mRNA export protein



GLOBAL-STATISTICS INFOmissingINFO

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/468	0.46	0/632
1	B	0.31	0/449	0.49	0/606
1	C	0.29	0/468	0.47	0/632
1	D	0.32	0/449	0.50	0/606
1	E	0.32	0/468	0.47	0/632
1	F	0.29	0/432	0.47	0/584
1	G	0.30	0/468	0.47	0/632
1	H	0.32	0/449	0.50	0/606
All	All	0.31	0/3651	0.48	0/4930

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	471	463	457	1	0
1	B	452	447	441	8	0
1	C	471	463	457	3	1
1	D	452	447	441	7	0
1	E	471	463	457	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	435	430	424	3	0
1	G	471	463	457	2	1
1	H	452	447	441	2	0
2	A	46	0	0	1	1
2	B	41	0	0	2	0
2	C	38	0	0	1	0
2	D	56	0	0	1	0
2	E	49	0	0	0	0
2	F	47	0	0	0	0
2	G	40	0	0	0	1
2	H	42	0	0	1	0
All	All	4034	3623	3575	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 23 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:GLU:OE2	2:A:701:HOH:O	2.11	0.69
1:E:621:GLN:OE1	1:F:604:GLN:NE2	2.29	0.66
1:B:615:MET:CE	1:B:619:TYR:HB3	2.29	0.63
1:B:615:MET:HE3	1:B:619:TYR:HB3	1.81	0.62
1:H:600:ALA:N	2:H:703:HOH:O	2.33	0.62

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:745:HOH:O	2:G:734:HOH:O[5_665]	2.09	0.11
1:C:643:LYS:HZ3	1:G:597:SER:O[4_455]	1.57	0.03

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	B	55/61 (90%)	54 (98%)	1 (2%)	0	100	100
1	C	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	D	55/61 (90%)	55 (100%)	0	0	100	100
1	E	58/61 (95%)	58 (100%)	0	0	100	100
1	F	53/61 (87%)	53 (100%)	0	0	100	100
1	G	58/61 (95%)	57 (98%)	1 (2%)	0	100	100
1	H	55/61 (90%)	55 (100%)	0	0	100	100
All	All	450/488 (92%)	446 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/49 (100%)	49 (100%)	0	100	100
1	B	47/49 (96%)	47 (100%)	0	100	100
1	C	49/49 (100%)	49 (100%)	0	100	100
1	D	47/49 (96%)	47 (100%)	0	100	100
1	E	49/49 (100%)	49 (100%)	0	100	100
1	F	45/49 (92%)	45 (100%)	0	100	100
1	G	49/49 (100%)	49 (100%)	0	100	100
1	H	47/49 (96%)	46 (98%)	1 (2%)	59	40
All	All	382/392 (97%)	381 (100%)	1 (0%)	94	91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	637	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	604	GLN
1	B	645	ASN

4.3.3 RNA ⓘ

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	623	1	6,8,9	1.92	2 (33%)	2,9,11	1.27	0
1	CAS	B	623	1	6,8,9	1.86	1 (16%)	2,9,11	1.44	0
1	CAS	C	623	1	6,8,9	1.92	1 (16%)	2,9,11	1.40	0
1	CAS	D	623	1	6,8,9	1.98	3 (50%)	2,9,11	1.17	0
1	CAS	E	623	1	6,8,9	1.87	1 (16%)	2,9,11	1.44	1 (50%)
1	CAS	F	623	1	6,8,9	2.04	2 (33%)	2,9,11	1.27	0
1	CAS	G	623	1	6,8,9	1.73	1 (16%)	2,9,11	1.28	0
1	CAS	H	623	1	6,8,9	1.95	2 (33%)	2,9,11	1.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	623	1	-	0/0/7/9	0/0/0/0
1	CAS	B	623	1	-	0/0/7/9	0/0/0/0
1	CAS	C	623	1	-	0/0/7/9	0/0/0/0
1	CAS	D	623	1	-	0/0/7/9	0/0/0/0
1	CAS	E	623	1	-	0/0/7/9	0/0/0/0
1	CAS	F	623	1	-	0/0/7/9	0/0/0/0
1	CAS	G	623	1	-	0/0/7/9	0/0/0/0
1	CAS	H	623	1	-	0/0/7/9	0/0/0/0

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	623	CAS	AS-CE2	-2.37	1.89	1.96
1	A	623	CAS	O-C	2.05	1.28	1.19
1	F	623	CAS	O-C	2.07	1.28	1.19
1	H	623	CAS	O-C	2.14	1.28	1.19
1	D	623	CAS	O-C	2.16	1.29	1.19

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	623	CAS	O-C-CA	-2.04	119.38	125.02

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	623	CAS	1	0
1	D	623	CAS	2	0
1	H	623	CAS	1	0

4.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

4.6 Ligand geometry [i](#)

There are no ligands in this entry.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	60/61 (98%)	0.67	4 (6%) 19 22	20, 29, 54, 65	0
1	B	57/61 (93%)	1.77	17 (29%) 1 0	21, 39, 87, 100	0
1	C	60/61 (98%)	0.86	7 (11%) 5 6	19, 33, 71, 89	0
1	D	57/61 (93%)	2.39	19 (33%) 0 0	17, 30, 76, 92	0
1	E	60/61 (98%)	1.03	7 (11%) 5 6	18, 27, 47, 99	0
1	F	55/61 (90%)	1.83	17 (30%) 0 0	18, 35, 82, 108	0
1	G	60/61 (98%)	0.87	6 (10%) 8 9	18, 32, 68, 80	0
1	H	57/61 (93%)	1.27	13 (22%) 1 1	17, 31, 80, 104	0
All	All	466/488 (95%)	1.32	90 (19%) 1 1	17, 31, 74, 108	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	657	VAL	14.8
1	D	648	PRO	9.4
1	B	600	ALA	9.3
1	B	657	VAL	9.1
1	F	600	ALA	8.9

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CAS	H	623	9/10	0.94	0.11	-	19,28,67,67	0
1	CAS	A	623	9/10	0.96	0.11	-	19,23,47,47	0
1	CAS	C	623	9/10	0.97	0.10	-	20,27,43,43	0
1	CAS	B	623	9/10	0.93	0.13	-	21,32,52,52	0
1	CAS	E	623	9/10	0.86	0.14	-	18,23,51,51	0
1	CAS	D	623	9/10	0.85	0.20	-	17,24,83,83	0
1	CAS	G	623	9/10	0.95	0.12	-	18,27,69,69	0
1	CAS	F	623	9/10	0.89	0.15	-	20,30,68,68	0

5.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.4 Ligands [i](#)

There are no ligands in this entry.

5.5 Other polymers [i](#)

There are no such residues in this entry.