



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 07:46 pm GMT

PDB ID : 1WR5  
Title : Three dimensional Structure of the E41K mutant of Tetraheme Cytochrome c3 from *Desulfovibrio vulgaris* Miyazaki F  
Authors : Tomimoto, Y.; Ogata, H.; Higuchi, Y.  
Deposited on : 2004-10-11  
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	1411 (1.40-1.40)
Ramachandran outliers	110173	1373 (1.40-1.40)
Sidechain outliers	110143	1372 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	108	 84%14%.

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EOH	A	2002	-	-	X	-
3	EOH	A	2005	-	-	X	-
3	EOH	A	2007	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	108	Total	C	N	O	S	0	0	0
			806	493	152	150	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	LYS	GLU	ENGINEERED	UNP P00132

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is ETHANOL (three-letter code: EOH) (formula: C<sub>2</sub>H<sub>6</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		
3	A	1	Total	C	O	0	0
			3	2	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	244	Total	O	0	0
			244	244		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Cytochrome c3

Chain A:  84% 14% .



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.12Å 67.51Å 34.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.40	Depositor
% Data completeness (in resolution range)	91.9 (10.00-1.40)	Depositor
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.134 , 0.225	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1264	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, EOH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	1/823 (0.1%)	1.20	3/1095 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	107	SER	C-OXT	5.01	1.32	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	46	CYS	C-N-CA	5.64	135.79	121.70
1	A	9	LEU	CA-CB-CG	5.29	127.47	115.30
1	A	46	CYS	O-C-N	-5.14	114.48	122.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	806	0	792	12	0
2	A	172	0	120	2	0
3	A	42	0	84	11	0
4	A	244	0	0	10	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	1264	0	996	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2010:EOH:H21	4:A:3098:HOH:O	1.91	0.71
1:A:85:GLU:O	3:A:2014:EOH:H12	1.92	0.70
1:A:63:LYS:HD3	4:A:3069:HOH:O	1.95	0.67
1:A:15:LYS:N	1:A:15:LYS:HD2	2.13	0.63
2:A:1001:HEM:O2D	3:A:2001:EOH:H22	2.03	0.58
1:A:29:LYS:HG3	4:A:3122:HOH:O	2.11	0.50
1:A:53:ASP:OD1	1:A:63:LYS:HE3	2.14	0.48
1:A:45:LYS:CB	3:A:2002:EOH:H12	2.44	0.47
1:A:13:LYS:NZ	3:A:2002:EOH:H22	2.30	0.47
1:A:101:LYS:NZ	4:A:3099:HOH:O	2.49	0.45
1:A:72:LYS:NZ	4:A:3213:HOH:O	2.49	0.44
1:A:71:ASP:OD1	3:A:2011:EOH:H21	2.17	0.44
3:A:2007:EOH:H22	4:A:3171:HOH:O	2.20	0.42
3:A:2007:EOH:H23	4:A:3081:HOH:O	2.20	0.42
3:A:2005:EOH:H22	4:A:3209:HOH:O	2.19	0.41
3:A:2007:EOH:H11	4:A:3112:HOH:O	2.20	0.41
1:A:96:GLU:CG	1:A:104:LYS:HG3	2.50	0.41
3:A:2005:EOH:H11	4:A:3133:HOH:O	2.19	0.41
1:A:5:PRO:HD3	2:A:1001:HEM:C4A	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	106/108 (98%)	104 (98%)	1 (1%)	1 (1%)	20 4

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	16	GLN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	85/85 (100%)	83 (98%)	2 (2%)	54 18

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LYS
1	A	29	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	HEM	A	1001	1	28,50,50	1.28	5 (17%)	17,82,82	2.57	9 (52%)
2	HEM	A	1002	1	28,50,50	1.33	5 (17%)	17,82,82	2.31	8 (47%)
2	HEM	A	1003	1	28,50,50	1.42	6 (21%)	17,82,82	2.08	7 (41%)
2	HEM	A	1004	1	28,50,50	1.30	4 (14%)	17,82,82	1.99	7 (41%)
3	EOH	A	2001	-	2,2,2	0.31	0	1,1,1	1.07	0
3	EOH	A	2002	-	2,2,2	0.58	0	1,1,1	0.37	0
3	EOH	A	2003	-	2,2,2	0.53	0	1,1,1	0.48	0
3	EOH	A	2004	-	2,2,2	0.48	0	1,1,1	0.28	0
3	EOH	A	2005	-	2,2,2	0.52	0	1,1,1	0.08	0
3	EOH	A	2006	-	2,2,2	0.46	0	1,1,1	0.09	0
3	EOH	A	2007	-	2,2,2	0.52	0	1,1,1	0.12	0
3	EOH	A	2008	-	2,2,2	0.51	0	1,1,1	0.39	0
3	EOH	A	2009	-	2,2,2	0.51	0	1,1,1	0.19	0
3	EOH	A	2010	-	2,2,2	0.55	0	1,1,1	0.40	0
3	EOH	A	2011	-	2,2,2	0.54	0	1,1,1	0.05	0
3	EOH	A	2012	-	2,2,2	0.44	0	1,1,1	0.00	0
3	EOH	A	2013	-	2,2,2	0.51	0	1,1,1	0.09	0
3	EOH	A	2014	-	2,2,2	0.55	0	1,1,1	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1001	1	-	0/6/54/54	0/0/8/8

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1002	1	-	0/6/54/54	0/0/8/8
2	HEM	A	1003	1	-	0/6/54/54	0/0/8/8
2	HEM	A	1004	1	-	0/6/54/54	0/0/8/8
3	EOH	A	2001	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2002	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2003	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2004	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2005	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2006	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2007	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2008	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2009	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2010	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2011	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2012	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2013	-	-	0/0/0/0	0/0/0/0
3	EOH	A	2014	-	-	0/0/0/0	0/0/0/0

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1003	HEM	C3B-C2B	-3.73	1.35	1.40
2	A	1004	HEM	C3B-CAB	-3.28	1.41	1.47
2	A	1002	HEM	C3B-CAB	-2.86	1.42	1.47
2	A	1001	HEM	C3C-CAC	-2.76	1.42	1.47
2	A	1004	HEM	C3B-C2B	-2.58	1.36	1.40
2	A	1003	HEM	C4D-ND	-2.56	1.33	1.36
2	A	1002	HEM	C3B-C2B	-2.55	1.37	1.40
2	A	1002	HEM	C3C-CAC	-2.42	1.42	1.47
2	A	1001	HEM	C3B-C2B	-2.28	1.37	1.40
2	A	1001	HEM	C3B-CAB	-2.27	1.43	1.47
2	A	1002	HEM	C1C-NC	-2.22	1.34	1.36
2	A	1003	HEM	C3C-CAC	-2.15	1.43	1.47
2	A	1003	HEM	C1B-NB	-2.04	1.34	1.36
2	A	1001	HEM	C4A-NA	2.11	1.40	1.36
2	A	1003	HEM	C4A-NA	2.11	1.40	1.36
2	A	1004	HEM	CBC-CAC	2.19	1.44	1.28
2	A	1003	HEM	CBB-CAB	2.53	1.46	1.28
2	A	1004	HEM	CBB-CAB	2.57	1.46	1.28
2	A	1002	HEM	CBB-CAB	2.71	1.47	1.28
2	A	1001	HEM	CBB-CAB	2.87	1.49	1.28

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	HEM	C1D-C2D-C3D	-5.74	103.00	107.00
2	A	1001	HEM	CMA-C3A-C4A	-5.38	120.20	128.46
2	A	1002	HEM	CMA-C3A-C4A	-4.96	120.84	128.46
2	A	1004	HEM	CMA-C3A-C4A	-3.91	122.46	128.46
2	A	1003	HEM	CMA-C3A-C4A	-3.61	122.92	128.46
2	A	1002	HEM	C4C-C3C-C2C	-3.31	104.59	106.90
2	A	1003	HEM	CMD-C2D-C1D	-2.96	123.91	128.46
2	A	1002	HEM	CBA-CAA-C2A	-2.75	107.22	112.48
2	A	1001	HEM	CAD-CBD-CGD	-2.62	108.18	112.66
2	A	1004	HEM	C1D-C2D-C3D	-2.37	105.34	107.00
2	A	1003	HEM	C1D-C2D-C3D	-2.28	105.41	107.00
2	A	1002	HEM	C1D-C2D-C3D	-2.25	105.43	107.00
2	A	1004	HEM	CMD-C2D-C1D	-2.14	125.17	128.46
2	A	1002	HEM	CMD-C2D-C1D	-2.14	125.18	128.46
2	A	1001	HEM	C3C-C4C-NC	-2.07	107.03	110.94
2	A	1001	HEM	CMB-C2B-C3B	2.01	128.62	124.89
2	A	1002	HEM	CMB-C2B-C3B	2.02	128.65	124.89
2	A	1002	HEM	CMC-C2C-C3C	2.19	128.96	124.89
2	A	1001	HEM	CMD-C2D-C3D	2.34	129.36	124.94
2	A	1001	HEM	CMC-C2C-C3C	2.36	129.27	124.89
2	A	1001	HEM	CMA-C3A-C2A	2.38	129.42	124.94
2	A	1003	HEM	CMA-C3A-C2A	2.38	129.43	124.94
2	A	1004	HEM	CMD-C2D-C3D	2.41	129.48	124.94
2	A	1001	HEM	CAA-CBA-CGA	2.61	117.11	112.66
2	A	1004	HEM	CMC-C2C-C3C	2.93	130.32	124.89
2	A	1003	HEM	CMD-C2D-C3D	2.95	130.50	124.94
2	A	1004	HEM	CMB-C2B-C3B	2.99	130.45	124.89
2	A	1004	HEM	CMA-C3A-C2A	3.12	130.83	124.94
2	A	1003	HEM	CMB-C2B-C3B	3.18	130.80	124.89
2	A	1002	HEM	CMA-C3A-C2A	3.54	131.61	124.94
2	A	1003	HEM	CMC-C2C-C3C	4.08	132.45	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	HEM	2	0
3	A	2001	EOH	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2002	EOH	2	0
3	A	2005	EOH	2	0
3	A	2007	EOH	3	0
3	A	2010	EOH	1	0
3	A	2011	EOH	1	0
3	A	2014	EOH	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.