



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 13, 2017 – 07:32 AM EST

PDB ID : 3WRM  
Title : Crystal structure of P450cam  
Authors : Kishimoto, A.; Takagi, K.; Amano, A.; Sakurai, K.; Mizushima, T.; Shimada, H.  
Deposited on : unknown  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	421	 2% 86% 10% •
1	F	421	 3% 88% 7% ••

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6961 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Camphor 5-monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	405	3223	2045	562	599	17	0	3	0
1	F	405	3236	2055	566	597	18	0	5	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	415	HIS	-	EXPRESSION TAG	UNP P00183
A	416	HIS	-	EXPRESSION TAG	UNP P00183
A	417	HIS	-	EXPRESSION TAG	UNP P00183
A	418	HIS	-	EXPRESSION TAG	UNP P00183
A	419	HIS	-	EXPRESSION TAG	UNP P00183
A	420	HIS	-	EXPRESSION TAG	UNP P00183
F	415	HIS	-	EXPRESSION TAG	UNP P00183
F	416	HIS	-	EXPRESSION TAG	UNP P00183
F	417	HIS	-	EXPRESSION TAG	UNP P00183
F	418	HIS	-	EXPRESSION TAG	UNP P00183
F	419	HIS	-	EXPRESSION TAG	UNP P00183
F	420	HIS	-	EXPRESSION TAG	UNP P00183

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	F	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		
3	F	1	Total	K	0	0
			1	1		

- Molecule 4 is CAMPHOR (three-letter code: CAM) (formula: C<sub>10</sub>H<sub>16</sub>O).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 11 10 1	0	0
4	F	1	Total C O 11 10 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	212	Total O 212 212	0	0
5	F	180	Total O 180 180	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Camphor 5-monooxygenase



- Molecule 1: Camphor 5-monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.09Å 62.54Å 95.49Å 90.00° 90.88° 90.00°	Depositor
Resolution (Å)	39.18 – 1.95 39.18 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (39.18-1.95) 99.7 (39.18-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.30 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.177 , 0.233 0.178 , 0.235	Depositor DCC
$R_{free}$ test set	2921 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.2	Xtrriage
Anisotropy	0.064	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6961	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, K, CAM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.68	0/3312	0.72	0/4501
1	F	0.68	1/3330 (0.0%)	0.73	3/4523 (0.1%)
All	All	0.68	1/6642 (0.0%)	0.72	3/9024 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	91	GLU	CG-CD	5.03	1.59	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	277	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	F	277	ARG	NE-CZ-NH2	-5.73	117.44	120.30
1	F	165	LEU	CA-CB-CG	5.43	127.80	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3223	0	3173	27	0
1	F	3236	0	3206	27	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	43	0	30	3	0
2	F	43	0	30	5	0
3	A	1	0	0	0	0
3	F	1	0	0	0	0
4	A	11	0	16	0	0
4	F	11	0	16	1	0
5	A	212	0	0	1	0
5	F	180	0	0	6	0
All	All	6961	0	6471	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:163:PHE:CE2	1:F:246:LEU:HD23	1.75	1.18
1:F:163:PHE:HE2	1:F:246:LEU:HD23	1.19	0.97
1:F:163:PHE:HE2	1:F:246:LEU:CD2	1.78	0.94
1:F:163:PHE:CE2	1:F:246:LEU:CD2	2.56	0.88
1:F:400:GLN:HA	1:F:400:GLN:HE21	1.41	0.85
1:A:343:GLN:H	1:A:343:GLN:CD	1.85	0.80
1:F:277:ARG:HG3	5:F:746:HOH:O	1.88	0.73
1:F:95:ALA:O	1:F:197:LYS:HE2	1.87	0.73
1:F:13:PRO:HD2	5:F:690:HOH:O	1.89	0.72
1:F:56:THR:O	1:F:61:GLY:HA2	1.93	0.68
2:F:501:HEM:HMC1	2:F:501:HEM:HBC2	1.76	0.67
1:A:171:GLU:OE1	1:F:277:ARG:HD3	1.97	0.64
1:F:33:ASN:HB2	5:F:696:HOH:O	1.96	0.64
2:A:501:HEM:HBC2	2:A:501:HEM:HMC1	1.83	0.61
1:A:200:LEU:HD11	1:A:246:LEU:HD13	1.84	0.58
1:A:368:ILE:O	1:A:372:LYS:HG3	2.08	0.54
1:A:179:TYR:OH	1:A:190:SER:OG	2.26	0.53
1:A:177:LEU:O	1:A:181[A]:THR:HG23	2.09	0.53
1:F:114:LEU:HD23	1:F:233:ILE:HD12	1.91	0.52
2:F:501:HEM:CMC	2:F:501:HEM:HBC2	2.38	0.52
1:F:33:ASN:ND2	5:F:695:HOH:O	2.43	0.51
1:A:141:SER:O	1:A:145:GLN:NE2	2.43	0.51
1:A:206:PRO:HG2	5:A:808:HOH:O	2.11	0.50
1:A:179:TYR:HH	1:A:190:SER:HG	1.59	0.50
1:F:17:HIS:CD2	1:F:313:LYS:HG2	2.46	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:TYR:CD2	1:A:239:LYS:HE3	2.48	0.49
1:F:179:TYR:OH	1:F:190:SER:OG	2.30	0.48
1:A:67:ARG:HH12	1:A:329[B]:GLU:CD	2.16	0.48
1:F:344:LYS:HA	1:F:344:LYS:HD2	1.62	0.48
1:A:143:ARG:HB3	1:A:144:PRO:CD	2.43	0.48
1:A:98:PHE:HB3	1:A:244:LEU:HB2	1.95	0.47
1:A:214:LYS:NZ	1:A:214:LYS:HB2	2.29	0.47
1:F:121:MET:HB3	1:F:122:PRO:HD3	1.96	0.47
1:F:98:PHE:HB3	1:F:244:LEU:HB2	1.96	0.47
1:F:400:GLN:CA	1:F:400:GLN:HE21	2.20	0.46
1:F:400:GLN:HA	1:F:400:GLN:NE2	2.21	0.46
1:A:344:LYS:HB2	1:A:344:LYS:HE2	1.69	0.46
1:F:186:ARG:HD2	1:F:392:LYS:HG3	1.98	0.45
1:A:49:ASN:N	1:A:49:ASN:OD1	2.40	0.45
2:A:501:HEM:HBC2	2:A:501:HEM:CMC	2.46	0.45
1:A:212:ARG:HG3	1:A:224:ALA:HB1	1.98	0.45
1:A:121:MET:HB3	1:A:122:PRO:HD3	2.00	0.44
2:F:501:HEM:HBB2	2:F:501:HEM:HMB2	2.00	0.44
2:F:501:HEM:C1A	4:F:503:CAM:H4	2.53	0.44
1:F:303:SER:HA	1:F:314:LYS:HB2	1.99	0.43
1:F:372:LYS:HG3	5:F:743:HOH:O	2.19	0.43
1:A:161:ARG:CZ	1:F:277:ARG:HD2	2.49	0.42
2:A:501:HEM:HMB2	2:A:501:HEM:HBB2	2.01	0.42
1:A:99:ILE:HG22	1:A:241:MET:HG2	2.02	0.42
1:F:91:GLU:HB3	5:F:666:HOH:O	2.20	0.42
1:A:201:TYR:CE2	1:A:239:LYS:HE3	2.55	0.41
1:A:72:ARG:O	1:A:76:GLU:HG3	2.20	0.41
1:A:143:ARG:N	1:A:144:PRO:HD2	2.35	0.41
2:F:501:HEM:HBB2	2:F:501:HEM:CMB	2.51	0.41
1:A:75:TYR:CZ	1:A:320:LEU:HB2	2.56	0.41
1:F:163:PHE:HE2	1:F:246:LEU:HD21	1.75	0.41
1:A:343:GLN:N	1:A:343:GLN:CD	2.63	0.41
1:F:276:GLU:C	1:F:277:ARG:HG2	2.40	0.40
1:A:162:ILE:HA	1:A:162:ILE:HD13	1.85	0.40
1:A:91:GLU:H	1:A:91:GLU:CD	2.24	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/421 (96%)	400 (98%)	6 (2%)	0	100	100
1	F	408/421 (97%)	396 (97%)	12 (3%)	0	100	100
All	All	814/842 (97%)	796 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	352/365 (96%)	343 (97%)	9 (3%)	51	40
1	F	354/365 (97%)	345 (98%)	9 (2%)	53	42
All	All	706/730 (97%)	688 (98%)	18 (2%)	51	42

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	ASN
1	A	70	LEU
1	A	126	LYS
1	A	134	LEU
1	A	276	GLU
1	A	277	ARG
1	A	343	GLN
1	A	382	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	412	LYS
1	F	70	LEU
1	F	84	GLU
1	F	91	GLU
1	F	126	LYS
1	F	165	LEU
1	F	209	GLU
1	F	285	CYS
1	F	400	GLN
1	F	412	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	ASN
1	F	33	ASN
1	F	110	GLN
1	F	117	GLN
1	F	129	ASN
1	F	213	GLN
1	F	229	ASN
1	F	400	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	501	1	28,50,50	2.12	9 (32%)	17,82,82	1.69	5 (29%)
4	CAM	A	503	-	12,12,12	3.05	2 (16%)	20,21,21	1.62	5 (25%)
2	HEM	F	501	1	28,50,50	2.16	9 (32%)	17,82,82	1.66	3 (17%)
4	CAM	F	503	-	12,12,12	3.24	1 (8%)	20,21,21	0.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	1	-	0/6/54/54	0/0/8/8
4	CAM	A	503	-	-	0/0/31/31	0/0/2/2
2	HEM	F	501	1	-	0/6/54/54	0/0/8/8
4	CAM	F	503	-	-	0/0/31/31	0/0/2/2

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	501	HEM	C3C-C2C	-4.34	1.34	1.40
2	F	501	HEM	C3B-C2B	-4.15	1.34	1.40
2	A	501	HEM	C3B-C2B	-3.87	1.35	1.40
4	A	503	CAM	C1-C2	-3.64	1.48	1.53
2	A	501	HEM	C3C-C2C	-3.47	1.35	1.40
2	F	501	HEM	CMA-C3A	2.07	1.55	1.51
2	A	501	HEM	C4A-NA	2.30	1.41	1.36
2	F	501	HEM	C4D-ND	2.39	1.39	1.36
2	A	501	HEM	C1B-NB	2.40	1.39	1.36
2	A	501	HEM	C4D-ND	2.61	1.39	1.36
2	F	501	HEM	CAA-C2A	2.68	1.56	1.52
2	A	501	HEM	CMA-C3A	2.80	1.57	1.51
2	F	501	HEM	C1B-NB	2.85	1.40	1.36
2	F	501	HEM	C3C-CAC	3.11	1.53	1.47
2	A	501	HEM	C3C-CAC	3.33	1.54	1.47

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	501	HEM	C3B-CAB	3.47	1.54	1.47
2	A	501	HEM	C3B-CAB	3.70	1.55	1.47
2	A	501	HEM	C3D-C2D	4.94	1.52	1.37
2	F	501	HEM	C3D-C2D	5.17	1.53	1.37
4	A	503	CAM	O-C2	9.77	1.38	1.21
4	F	503	CAM	O-C2	10.76	1.40	1.21

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	CAD-CBD-CGD	-3.65	106.42	112.66
2	F	501	HEM	CAD-CBD-CGD	-3.39	106.88	112.66
4	A	503	CAM	C4-C3-C2	-3.07	96.69	102.00
2	F	501	HEM	C1D-C2D-C3D	-3.06	104.87	107.00
4	A	503	CAM	O-C2-C3	-2.62	121.06	126.51
4	A	503	CAM	C5-C4-C3	-2.61	99.22	106.44
4	F	503	CAM	C4-C3-C2	-2.43	97.79	102.00
4	A	503	CAM	C5-C6-C1	-2.32	99.94	104.73
2	F	501	HEM	CMA-C3A-C4A	-2.15	125.16	128.46
2	A	501	HEM	C3B-C4B-NB	-2.12	106.47	109.21
2	A	501	HEM	CMA-C3A-C4A	-2.05	125.31	128.46
2	A	501	HEM	CMC-C2C-C3C	2.14	128.86	124.89
2	A	501	HEM	CMB-C2B-C3B	2.36	129.27	124.89
4	A	503	CAM	C6-C5-C4	2.41	106.95	103.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	3	0
2	F	501	HEM	5	0
4	F	503	CAM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	405/421 (96%)	-0.05	8 (1%) 65 75	6, 16, 31, 44	17 (4%)
1	F	405/421 (96%)	0.01	11 (2%) 55 65	6, 18, 33, 44	14 (3%)
All	All	810/842 (96%)	-0.02	19 (2%) 61 71	6, 17, 32, 44	31 (3%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	414	VAL	4.4
1	A	413	ALA	4.0
1	F	119	VAL	3.9
1	A	88	ILE	3.7
1	F	246	LEU	3.6
1	A	119	VAL	3.4
1	A	277	ARG	3.3
1	F	24	PHE	3.0
1	F	49	ASN	2.9
1	F	22	LEU	2.9
1	F	44	VAL	2.8
1	F	88	ILE	2.7
1	A	49	ASN	2.6
1	F	309	GLY	2.5
1	A	412	LYS	2.4
1	A	414	VAL	2.4
1	F	20	GLU	2.4
1	A	385	PRO	2.1
1	F	14	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CAM	F	503	11/11	0.97	0.18	2.00	9,10,13,15	0
4	CAM	A	503	11/11	0.97	0.16	1.73	4,7,10,11	0
2	HEM	F	501	43/43	0.98	0.14	0.29	8,11,14,19	0
2	HEM	A	501	43/43	0.98	0.13	0.23	4,8,12,19	0
3	K	A	502	1/1	0.99	0.07	-0.87	10,10,10,10	0
3	K	F	502	1/1	0.99	0.07	-0.88	16,16,16,16	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.