



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 10, 2017 – 04:35 PM EDT

PDB ID : 5WTA
Title : Crystal Structure of Staphylococcus aureus SdrE apo form
Authors : Wu, M.; Zhang, Y.; Hang, T.; Wang, C.; Yang, Y.; Zang, J.; Zhang, M.;
Zhang, X.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	338	<div> <div> <div></div> <div>83%</div> <div>9%</div> <div>7%</div> </div> </div>
1	B	338	<div> <div> <div>79%</div> <div>13%</div> <div>7%</div> </div> </div>
1	C	338	<div> <div> <div>80%</div> <div>12%</div> <div>7%</div> </div> </div>
1	D	338	<div> <div> <div>79%</div> <div>12%</div> <div>7%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10565 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine-aspartate repeat-containing protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	0	0
			2478	1546	401	527	4			
1	B	315	Total	C	N	O	S	0	1	0
			2481	1548	401	528	4			
1	C	314	Total	C	N	O	S	0	0	0
			2472	1543	400	525	4			
1	D	314	Total	C	N	O	S	0	3	0
			2487	1552	402	529	4			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	489	MET	ILE	engineered mutation	UNP Q932F7
A	600	LEU	-	expression tag	UNP Q932F7
A	601	GLU	-	expression tag	UNP Q932F7
A	602	HIS	-	expression tag	UNP Q932F7
A	603	HIS	-	expression tag	UNP Q932F7
A	604	HIS	-	expression tag	UNP Q932F7
A	605	HIS	-	expression tag	UNP Q932F7
A	606	HIS	-	expression tag	UNP Q932F7
A	607	HIS	-	expression tag	UNP Q932F7
B	489	MET	ILE	engineered mutation	UNP Q932F7
B	600	LEU	-	expression tag	UNP Q932F7
B	601	GLU	-	expression tag	UNP Q932F7
B	602	HIS	-	expression tag	UNP Q932F7
B	603	HIS	-	expression tag	UNP Q932F7
B	604	HIS	-	expression tag	UNP Q932F7
B	605	HIS	-	expression tag	UNP Q932F7
B	606	HIS	-	expression tag	UNP Q932F7
B	607	HIS	-	expression tag	UNP Q932F7
C	489	MET	ILE	engineered mutation	UNP Q932F7
C	600	LEU	-	expression tag	UNP Q932F7
C	601	GLU	-	expression tag	UNP Q932F7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	602	HIS	-	expression tag	UNP Q932F7
C	603	HIS	-	expression tag	UNP Q932F7
C	604	HIS	-	expression tag	UNP Q932F7
C	605	HIS	-	expression tag	UNP Q932F7
C	606	HIS	-	expression tag	UNP Q932F7
C	607	HIS	-	expression tag	UNP Q932F7
D	489	MET	ILE	engineered mutation	UNP Q932F7
D	600	LEU	-	expression tag	UNP Q932F7
D	601	GLU	-	expression tag	UNP Q932F7
D	602	HIS	-	expression tag	UNP Q932F7
D	603	HIS	-	expression tag	UNP Q932F7
D	604	HIS	-	expression tag	UNP Q932F7
D	605	HIS	-	expression tag	UNP Q932F7
D	606	HIS	-	expression tag	UNP Q932F7
D	607	HIS	-	expression tag	UNP Q932F7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	157	Total O 157 157	0	0
2	B	167	Total O 167 167	0	0
2	C	147	Total O 147 147	0	0
2	D	176	Total O 176 176	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine-aspartate repeat-containing protein E




- Molecule 1: Serine-aspartate repeat-containing protein E



- Molecule 1: Serine-aspartate repeat-containing protein E



- Molecule 1: Serine-aspartate repeat-containing protein E





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.81Å 61.73Å 139.61Å 80.89° 89.83° 73.34°	Depositor
Resolution (Å)	38.93 – 2.30 38.63 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.5 (38.93-2.30) 90.0 (38.63-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.194 , 0.252 0.196 , 0.251	Depositor DCC
R_{free} test set	2821 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10565	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2518	0.75	0/3418
1	B	0.67	0/2524	0.75	0/3426
1	C	0.66	0/2512	0.74	1/3410 (0.0%)
1	D	0.69	0/2536	0.75	0/3443
All	All	0.68	0/10090	0.74	1/13697 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	303	ASP	CB-CG-OD1	5.20	122.98	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	339	ASN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2478	0	2392	18	0
1	B	2481	0	2397	24	0
1	C	2472	0	2387	19	0
1	D	2487	0	2403	24	0
2	A	157	0	0	3	0
2	B	167	0	0	2	0
2	C	147	0	0	3	1
2	D	176	0	0	8	1
All	All	10565	0	9579	83	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:527:ASN:HB2	2:A:837:HOH:O	1.72	0.87
1:D:340:ASP:HB2	2:D:723:HOH:O	1.74	0.86
1:B:315:LYS:NZ	1:D:521:ASN:O	2.12	0.81
1:A:438:GLU:HG2	2:A:747:HOH:O	1.86	0.73
1:A:336:THR:OG1	1:A:337:ASP:N	2.27	0.65
1:D:372:VAL:O	2:D:701:HOH:O	2.11	0.65
1:B:313:ASP:OD1	1:B:315:LYS:HE3	1.97	0.64
2:A:706:HOH:O	1:B:282:THR:HG21	1.97	0.63
1:B:488:GLU:HB2	2:B:834:HOH:O	1.99	0.62
1:C:406:ALA:O	1:C:408:LYS:HE2	2.00	0.61
1:C:554:ASP:HB3	1:C:556:GLU:HG2	1.82	0.61
1:D:379:LYS:NZ	2:D:703:HOH:O	2.22	0.61
1:C:499:GLN:HG2	2:C:780:HOH:O	2.00	0.60
1:D:370:ASP:OD1	2:D:702:HOH:O	2.16	0.60
1:D:340:ASP:CB	2:D:723:HOH:O	2.39	0.57
1:C:435:LYS:CD	2:C:845:HOH:O	2.53	0.56
1:D:468:VAL:HG13	1:D:472:GLY:HA2	1.88	0.56
1:D:309:GLU:HG2	1:D:381:ARG:HB2	1.88	0.55
1:A:432:ILE:HA	1:A:581:ASN:ND2	2.23	0.54
1:C:489:MET:HE1	1:C:532:LEU:HD11	1.89	0.53
1:C:419:GLN:HE22	1:C:432:ILE:HG23	1.73	0.53
1:C:435:LYS:HD2	2:C:845:HOH:O	2.07	0.53
1:B:456:ALA:HB1	1:B:567:THR:HG21	1.90	0.52
1:D:519:PHE:O	1:D:523:LYS:HG3	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:436:LEU:HD11	1:A:438:GLU:HG3	1.92	0.51
1:D:557:LEU:N	1:D:585:THR:OG1	2.42	0.51
1:B:334:ASP:OD2	1:B:390:LYS:NZ	2.42	0.51
1:C:299:HIS:CE1	1:C:387:TYR:HE2	2.28	0.51
1:B:508:ASP:OD1	1:B:510:SER:OG	2.29	0.51
1:A:419:GLN:HE22	1:A:432:ILE:HG23	1.76	0.51
1:D:408:LYS:HD2	2:D:802:HOH:O	2.09	0.51
1:B:324:ILE:HD12	1:B:382:LEU:HD23	1.93	0.50
1:D:505:ARG:NH1	2:D:706:HOH:O	2.30	0.49
1:D:325:ASN:HB2	1:D:404:ALA:HB3	1.94	0.49
1:A:436:LEU:CD1	1:A:438:GLU:HG3	2.42	0.49
1:C:277:ASP:OD1	1:C:277:ASP:N	2.44	0.49
1:A:494:VAL:HG21	1:A:500:LEU:HD21	1.94	0.48
1:C:442:THR:HG22	1:C:548:LYS:HD3	1.95	0.48
1:C:556:GLU:HB3	1:C:585:THR:O	2.14	0.48
1:A:325:ASN:HB2	1:A:404:ALA:HB3	1.96	0.47
1:B:325:ASN:HB2	1:B:404:ALA:HB3	1.97	0.47
1:B:311:THR:HG22	2:B:710:HOH:O	2.13	0.47
1:B:351:VAL:HG12	1:B:369:THR:HG21	1.97	0.47
1:A:302:LYS:HE2	1:A:417:ASP:HB2	1.98	0.46
1:C:448:TYR:CZ	1:C:504:ASN:HA	2.50	0.46
1:C:467:GLN:HG2	1:C:468:VAL:N	2.31	0.46
1:A:277:ASP:N	1:A:277:ASP:OD1	2.49	0.46
1:B:283:LYS:O	1:B:308:THR:HA	2.16	0.46
1:C:552:THR:OG1	1:C:554:ASP:HB2	2.17	0.45
1:D:495:ASN:HB2	1:D:498:GLN:HG3	1.99	0.44
1:A:489:MET:HE2	1:A:532:LEU:HD11	1.99	0.44
1:A:521:ASN:OD1	1:A:521:ASN:N	2.50	0.44
1:A:514:ASP:OD1	1:A:514:ASP:C	2.57	0.43
1:D:317:LYS:NZ	2:D:707:HOH:O	2.34	0.43
1:D:456:ALA:HB1	1:D:567:THR:HG21	2.00	0.43
1:A:459:THR:HA	1:A:566:ARG:O	2.19	0.43
1:B:402:THR:OG1	1:B:411:SER:OG	2.22	0.43
1:D:370:ASP:N	1:D:370:ASP:OD1	2.45	0.43
1:A:435:LYS:HB2	1:A:435:LYS:HE3	1.84	0.42
1:B:495:ASN:HB2	1:B:498:GLN:HG3	2.00	0.42
1:B:397:THR:OG1	1:B:398:SER:N	2.52	0.42
1:D:391:GLN:HB2	1:D:507:TYR:CZ	2.54	0.42
1:B:461:VAL:HA	1:B:564:SER:O	2.20	0.42
1:D:275:VAL:HB	1:D:278:LEU:HD12	2.01	0.42
1:C:299:HIS:NE2	1:C:387:TYR:HE2	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:400[A]:ASN:OD1	1:D:413:ASN:HB2	2.20	0.42
1:B:557:LEU:O	1:B:584:VAL:HA	2.20	0.42
1:B:324:ILE:HD12	1:B:382:LEU:CD2	2.49	0.41
1:C:426:ASP:HA	1:C:454:LYS:HD3	2.01	0.41
1:C:519:PHE:O	1:C:523:LYS:HG3	2.20	0.41
1:A:357:PHE:HE2	1:A:359:LYS:HG2	1.85	0.41
1:B:278:LEU:HB3	1:B:313:ASP:HB3	2.03	0.41
1:B:432:ILE:HD13	1:B:504:ASN:HB3	2.03	0.41
1:B:435:LYS:HB3	1:B:444:GLU:HB3	2.03	0.41
1:D:462:ASP:HA	1:D:530:ALA:O	2.20	0.41
1:C:374:LYS:HG3	1:C:374:LYS:H	1.63	0.41
1:A:463:ILE:HD13	1:A:545:VAL:HG21	2.02	0.41
1:B:481:THR:HG23	1:B:560:ALA:O	2.21	0.41
1:D:440:LYS:O	1:D:441:GLN:HG2	2.21	0.41
1:B:435:LYS:HB2	1:B:435:LYS:HE3	1.91	0.40
1:B:399:LEU:HD23	1:B:399:LEU:N	2.36	0.40
1:D:497:ASN:N	1:D:497:ASN:OD1	2.54	0.40
1:C:295:VAL:CG2	1:D:471:TYR:CD1	3.05	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:746:HOH:O	2:D:795:HOH:O[1_655]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/338 (93%)	307 (98%)	6 (2%)	0	100	100
1	B	314/338 (93%)	306 (98%)	8 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	312/338 (92%)	304 (97%)	8 (3%)	0	100	100
1	D	315/338 (93%)	302 (96%)	13 (4%)	0	100	100
All	All	1254/1352 (93%)	1219 (97%)	35 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/301 (94%)	269 (95%)	14 (5%)	29	39
1	B	284/301 (94%)	270 (95%)	14 (5%)	29	39
1	C	282/301 (94%)	263 (93%)	19 (7%)	19	24
1	D	285/301 (95%)	269 (94%)	16 (6%)	25	33
All	All	1134/1204 (94%)	1071 (94%)	63 (6%)	25	33

All (63) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	282	THR
1	A	292	LYS
1	A	331	ILE
1	A	336	THR
1	A	337	ASP
1	A	348	SER
1	A	399	LEU
1	A	489	MET
1	A	504	ASN
1	A	521	ASN
1	A	522	LYS
1	A	553	SER
1	A	579	TYR
1	A	580	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	282	THR
1	B	311	THR
1	B	331	ILE
1	B	338	LYS
1	B	346	ASP
1	B	379	LYS
1	B	428	ASN
1	B	455	THR
1	B	504	ASN
1	B	522	LYS
1	B	553	SER
1	B	579	TYR
1	B	580	SER
1	B	583	ILE
1	C	273	ASN
1	C	315	LYS
1	C	328	LYS
1	C	374	LYS
1	C	379	LYS
1	C	381	ARG
1	C	399	LEU
1	C	408	LYS
1	C	415	SER
1	C	417	ASP
1	C	439	ASN
1	C	476	LEU
1	C	499	GLN
1	C	504	ASN
1	C	526	SER
1	C	527	ASN
1	C	539	SER
1	C	553	SER
1	C	579	TYR
1	D	283	LYS
1	D	292	LYS
1	D	295	VAL
1	D	317	LYS
1	D	328	LYS
1	D	354	LYS
1	D	381	ARG
1	D	399	LEU
1	D	408	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	441	GLN
1	D	485	GLN
1	D	504	ASN
1	D	511	GLN
1	D	539	SER
1	D	553	SER
1	D	579	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	315/338 (93%)	-0.28	2 (0%) 89 92	12, 20, 33, 65	0
1	B	315/338 (93%)	-0.30	0 100 100	12, 20, 37, 51	0
1	C	314/338 (92%)	-0.20	4 (1%) 77 81	11, 21, 39, 53	0
1	D	314/338 (92%)	-0.22	1 (0%) 93 96	10, 20, 40, 70	0
All	All	1258/1352 (93%)	-0.25	7 (0%) 89 92	10, 20, 38, 70	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	338	LYS	3.6
1	D	553	SER	2.8
1	A	338	LYS	2.7
1	C	586	SER	2.6
1	C	339	ASN	2.2
1	A	272	SER	2.2
1	C	340	ASP	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.