



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 07:10 pm GMT

PDB ID : 1WWS
Title : Crystal structure of ttk003001566 from *Thermus Thermophilus* HB8
Authors : Wang, H.; Murayama, K.; Terada, T.; Shirouzu, M.; Kuramitsu, S.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2005-01-18
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949

i

X-RAY DIFFRACTION

A.

Metric	Percentile Banks	Value
--------	------------------	-------



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div></div> <div>79%</div> <div>20%</div> </div>
1	B	148	<div> <div></div> <div>78%</div> <div>22%</div> </div>
1	C	148	<div> <div>5%</div> <div></div> <div>72%</div> <div>28%</div> </div>
1	D	148	<div> <div>4%</div> <div></div> <div>76%</div> <div>24%</div> </div>
1	E	148	<div> <div>2%</div> <div></div> <div>89%</div> <div>11%</div> </div>
1	F	148	<div> <div></div> <div>78%</div> <div>20%</div> </div>



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	148	 3% 82% 17%
1	H	148	 2% 76% 23%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	B	149	-	-	-	X
2	CA	F	149	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein TTHA1479.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	B	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	C	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	D	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	E	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	F	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	G	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			
1	H	148	Total	C	N	O	Se	0	0	0
			1188	758	208	219	3			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
A	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
A	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
B	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
B	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
B	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
C	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
C	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
C	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
D	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
D	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
D	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
E	1	MSE	MET	MODIFIED RESIDUE	GB 55981448

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
E	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
F	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
F	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
F	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
G	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
G	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
G	76	MSE	MET	MODIFIED RESIDUE	GB 55981448
H	1	MSE	MET	MODIFIED RESIDUE	GB 55981448
H	3	MSE	MET	MODIFIED RESIDUE	GB 55981448
H	76	MSE	MET	MODIFIED RESIDUE	GB 55981448

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	114	Total O 114 114	0	0
3	B	104	Total O 104 104	0	0
3	C	67	Total O 67 67	0	0
3	D	66	Total O 66 66	0	0
3	E	116	Total O 116 116	0	0
3	F	85	Total O 85 85	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	67	Total	O	0	0
			67	67		
3	H	90	Total	O	0	0
			90	90		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: hypothetical protein TTHA1479




- Molecule 1: hypothetical protein TTHA1479



- Molecule 1: hypothetical protein TTHA1479





- Molecule 1: hypothetical protein TTHA1479








- Molecule 1: hypothetical protein TTHA1479




- Molecule 1: hypothetical protein TTHA1479



- Molecule 1: hypothetical protein TTHA1479



- Molecule 1: hypothetical protein TTHA1479



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	90.77Å 70.58Å 97.90Å 90.00° 106.60° 90.00°	Depositor
Resolution (Å)	44.00 – 1.90 44.00 – 1.89	Depositor EDS
% Data completeness (in resolution range)	96.9 (44.00-1.90) 95.6 (44.00-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 1.88Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.208 , 0.254 0.207 , 0.254	Depositor DCC
R_{free} test set	4564 reflections (5.03%)	DCC
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10219	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1204	0.53	0/1624
1	B	0.30	0/1204	0.53	0/1624
1	C	0.27	0/1204	0.47	0/1624
1	D	0.28	0/1204	0.48	0/1624
1	E	0.30	0/1204	0.53	0/1624
1	F	0.29	0/1204	0.54	0/1624
1	G	0.28	0/1204	0.48	0/1624
1	H	0.29	0/1204	0.48	0/1624
All	All	0.29	0/9632	0.51	0/12992

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1188	0	1227	26	0
1	B	1188	0	1227	26	0
1	C	1188	0	1227	36	0
1	D	1188	0	1227	42	0
1	E	1188	0	1227	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1188	0	1227	21	0
1	G	1188	0	1227	26	0
1	H	1188	0	1227	31	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
3	A	114	0	0	0	0
3	B	104	0	0	2	0
3	C	67	0	0	2	0
3	D	66	0	0	2	0
3	E	116	0	0	5	0
3	F	85	0	0	1	0
3	G	67	0	0	0	0
3	H	90	0	0	1	0
All	All	10219	0	9816	208	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 208 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:GLU:HG2	1:A:104:ARG:HH22	1.26	0.98
1:C:71:GLN:HE21	1:C:75:ARG:HH22	1.15	0.91
1:H:82:LEU:HD11	1:H:86:LEU:HD22	1.53	0.89
1:B:71:GLN:NE2	1:B:74:ARG:HH11	1.74	0.86
1:C:125:LEU:HD23	1:D:125:LEU:HD23	1.60	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/148 (99%)	144 (99%)	2 (1%)	0	100	100
1	B	146/148 (99%)	143 (98%)	3 (2%)	0	100	100
1	C	146/148 (99%)	140 (96%)	6 (4%)	0	100	100
1	D	146/148 (99%)	142 (97%)	4 (3%)	0	100	100
1	E	146/148 (99%)	143 (98%)	3 (2%)	0	100	100
1	F	146/148 (99%)	144 (99%)	2 (1%)	0	100	100
1	G	146/148 (99%)	144 (99%)	2 (1%)	0	100	100
1	H	146/148 (99%)	141 (97%)	5 (3%)	0	100	100
All	All	1168/1184 (99%)	1141 (98%)	27 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/123 (102%)	125 (99%)	1 (1%)	85	85
1	B	126/123 (102%)	123 (98%)	3 (2%)	54	47
1	C	126/123 (102%)	124 (98%)	2 (2%)	68	65
1	D	126/123 (102%)	126 (100%)	0	100	100
1	E	126/123 (102%)	125 (99%)	1 (1%)	85	85
1	F	126/123 (102%)	122 (97%)	4 (3%)	44	34
1	G	126/123 (102%)	125 (99%)	1 (1%)	85	85
1	H	126/123 (102%)	124 (98%)	2 (2%)	68	65
All	All	1008/984 (102%)	994 (99%)	14 (1%)	71	69

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	129	LEU
1	F	33	ARG
1	G	82	LEU
1	C	66	LEU
1	F	144	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	14	GLN
1	D	145	ASN
1	G	145	ASN
1	D	71	GLN
1	E	14	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	145/148 (97%)	-0.21	2 (1%) 75 78	13, 24, 50, 60	0
1	B	145/148 (97%)	-0.32	1 (0%) 87 89	13, 25, 45, 69	0
1	C	145/148 (97%)	0.14	8 (5%) 26 29	16, 32, 72, 80	0
1	D	145/148 (97%)	0.04	6 (4%) 38 42	17, 33, 66, 73	0
1	E	145/148 (97%)	-0.18	3 (2%) 64 67	14, 25, 48, 64	0
1	F	145/148 (97%)	-0.26	0 100 100	15, 27, 51, 67	0
1	G	145/148 (97%)	-0.01	5 (3%) 46 49	16, 32, 64, 77	0
1	H	145/148 (97%)	-0.04	3 (2%) 64 67	17, 30, 58, 70	0
All	All	1160/1184 (97%)	-0.10	28 (2%) 59 63	13, 29, 59, 80	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	129	LEU	3.8
1	C	133	GLN	3.5
1	G	17	GLY	3.5
1	C	22	LYS	3.4
1	D	125	LEU	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	CA	F	149	1/1	0.99	0.14	7.00	18,18,18,18	0
2	CA	B	149	1/1	0.99	0.15	5.67	22,22,22,22	0
2	CA	D	149	1/1	1.00	0.15	-	15,15,15,15	0
2	CA	G	149	1/1	1.00	0.17	-	11,11,11,11	0
2	CA	A	149	1/1	1.00	0.18	-	12,12,12,12	0
2	CA	E	149	1/1	1.00	0.17	-	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.