



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 15, 2017 – 06:43 PM EST

PDB ID : 4WWR
Title : Crystal Structure of Bag6-Ubl4A Dimerization Domain
Authors : Mock, J.Y.; Chartron, J.W.; Clemons Jr., W.M.
Deposited on : unknown
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	47	
1	D	47	
1	F	47	
1	H	47	
2	A	53	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	C	53	<div><div></div><div>8%</div><div>92%</div><div>8%</div></div>
2	E	53	<div><div></div><div>19%</div><div>83%</div><div>13%</div><div></div><div></div></div>
2	G	53	<div><div></div><div>6%</div><div>85%</div><div>13%</div><div></div><div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6530 atoms, of which 3234 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin-like protein 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	46	Total	C	H	N	O	0	0	0
			770	239	388	72	71			
1	H	45	Total	C	H	N	O	0	0	0
			756	234	381	71	70			
1	D	46	Total	C	H	N	O	0	0	0
			766	237	386	72	71			
1	F	46	Total	C	H	N	O	0	2	0
			786	244	395	73	74			

- Molecule 2 is a protein called Large proline-rich protein BAG6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	53	Total	C	H	N	O	0	0	0
			839	252	425	79	82			
2	G	52	Total	C	H	N	O	0	0	0
			823	247	417	78	81			
2	C	53	Total	C	H	N	O	0	0	0
			839	252	425	79	82			
2	E	52	Total	C	H	N	O	0	0	0
			822	247	417	77	80			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1053	MET	-	initiating methionine	UNP P46379
G	1053	MET	-	initiating methionine	UNP P46379
C	1053	MET	-	initiating methionine	UNP P46379
E	1053	MET	-	initiating methionine	UNP P46379

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	20	Total 20	O 20	0	0
3	A	27	Total 27	O 27	0	0
3	H	15	Total 15	O 15	0	0
3	G	11	Total 11	O 11	0	0
3	D	10	Total 10	O 10	0	0
3	C	27	Total 27	O 27	0	0
3	F	4	Total 4	O 4	0	0
3	E	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ubiquitin-like protein 4A





- Molecule 1: Ubiquitin-like protein 4A



- Molecule 1: Ubiquitin-like protein 4A




- Molecule 1: Ubiquitin-like protein 4A




- Molecule 2: Large proline-rich protein BAG6




- Molecule 2: Large proline-rich protein BAG6



• Molecule 2: Large proline-rich protein BAG6



• Molecule 2: Large proline-rich protein BAG6



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.23Å 56.49Å 75.55Å 90.00° 96.23° 90.00°	Depositor
Resolution (Å)	19.65 – 2.00 19.65 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.5 (19.65-2.00) 98.5 (19.65-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.98 (at 2.01Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.225 , 0.269 0.226 , 0.270	Depositor DCC
R_{free} test set	1334 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	34.1	Xtriage
Anisotropy	0.371	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6530	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.34	0/387	0.58	0/521
1	D	0.31	0/384	0.55	0/517
1	F	0.35	0/405	0.60	0/545
1	H	0.40	0/379	0.61	0/510
2	A	0.42	0/417	0.58	0/560
2	C	0.34	0/417	0.51	0/560
2	E	0.35	0/408	0.55	0/548
2	G	0.32	0/409	0.50	0/550
All	All	0.36	0/3206	0.56	0/4311

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	382	388	388	2	2
1	D	380	386	385	6	0
1	F	391	395	386	1	2
1	H	375	381	380	1	0
2	A	414	425	425	4	0
2	C	414	425	425	1	0
2	E	405	417	417	3	0
2	G	406	417	416	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	27	0	0	1	0
3	B	20	0	0	0	0
3	C	27	0	0	0	0
3	D	10	0	0	2	0
3	E	15	0	0	0	0
3	F	4	0	0	0	0
3	G	11	0	0	1	0
3	H	15	0	0	1	0
All	All	3296	3234	3222	19	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 19 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1053:MET:HE2	2:A:1100:ILE:HD11	1.72	0.71
2:G:1075:SER:OG	2:G:1077:GLU:OE1	2.16	0.64
1:H:94:GLN:N	3:H:215:HOH:O	2.29	0.64
1:D:122:ASP:OD1	1:D:125:ARG:NH2	2.33	0.61
1:F:109:ALA:O	1:F:113:SER:OG	2.17	0.61

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:128:SER:O	1:F:129:ARG:NH1[1_455]	2.10	0.10
1:B:128:SER:O	1:F:129:ARG:HH11[1_455]	1.59	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	44/47 (94%)	44 (100%)	0	0	100	100
1	D	44/47 (94%)	44 (100%)	0	0	100	100
1	F	46/47 (98%)	46 (100%)	0	0	100	100
1	H	43/47 (92%)	43 (100%)	0	0	100	100
2	A	51/53 (96%)	50 (98%)	1 (2%)	0	100	100
2	C	51/53 (96%)	49 (96%)	2 (4%)	0	100	100
2	E	50/53 (94%)	47 (94%)	3 (6%)	0	100	100
2	G	50/53 (94%)	48 (96%)	1 (2%)	1 (2%)	9	3
All	All	379/400 (95%)	371 (98%)	7 (2%)	1 (0%)	44	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	G	1055	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	42/42 (100%)	42 (100%)	0	100	100
1	D	41/42 (98%)	40 (98%)	1 (2%)	54	56
1	F	44/42 (105%)	42 (96%)	2 (4%)	32	27
1	H	41/42 (98%)	40 (98%)	1 (2%)	54	56
2	A	45/45 (100%)	43 (96%)	2 (4%)	33	28
2	C	45/45 (100%)	42 (93%)	3 (7%)	19	13
2	E	44/45 (98%)	38 (86%)	6 (14%)	4	2
2	G	44/45 (98%)	44 (100%)	0	100	100
All	All	346/348 (99%)	331 (96%)	15 (4%)	33	29

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	1055	LEU
1	F	103	LEU
2	E	1102	LYS
2	C	1054	GLN
2	E	1101	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	46/47 (97%)	0.12	1 (2%) 62 61	30, 40, 69, 89	0
1	D	46/47 (97%)	0.53	3 (6%) 20 20	37, 58, 94, 104	0
1	F	46/47 (97%)	1.02	8 (17%) 2 2	42, 69, 95, 192	0
1	H	45/47 (95%)	0.32	2 (4%) 35 35	26, 45, 71, 102	0
2	A	53/53 (100%)	-0.13	0 100 100	22, 33, 52, 76	0
2	C	53/53 (100%)	0.35	4 (7%) 15 15	32, 45, 73, 97	0
2	E	52/53 (98%)	1.06	10 (19%) 1 1	36, 60, 98, 216	0
2	G	52/53 (98%)	0.26	3 (5%) 24 24	27, 53, 79, 96	0
All	All	393/400 (98%)	0.44	31 (7%) 13 13	22, 50, 89, 216	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	1081	ARG	5.4
2	E	1104	LEU	4.7
2	E	1071	ARG	4.3
1	F	121	ARG	3.9
2	C	1073	LEU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.