



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 29, 2017 – 09:15 AM EDT

PDB ID : 2X08
Title : cytochrome c peroxidase: ascorbate bound to the engineered ascorbate binding site
Authors : Murphy, E.J.; Metcalfe, C.L.; Gumiero, A.; Raven, E.L.; Moody, P.C.E.
Deposited on : 2009-12-07
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029077
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029077


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ASC	A	1253	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	1	0
			2319	1480	387	446	6			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	36	ALA	TYR	engineered mutation	UNP P00431
A	184	ARG	ASN	engineered mutation	UNP P00431
A	191	PHE	TRP	engineered mutation	UNP P00431
A	257	ARG	LYS	conflict	UNP P00431

- Molecule 2 is ASCORBIC ACID (three-letter code: ASC) (formula: $C_6H_8O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	226	Total	O	0	0
			226	226		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYTOCHROME C PEROXIDASE, MITOCHONDRIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.00 Å 74.49 Å 106.47 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.92 – 2.01 43.31 – 2.02	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.92-2.01) 94.2 (43.31-2.02)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.56 (at 2.01 Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.201 , (Not available) 0.336 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 82.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	2600	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ASC, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2390	0.81	5/3243 (0.2%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	ARG	NE-CZ-NH2	-12.44	114.08	120.30
1	A	166	ARG	CG-CD-NE	-6.15	98.88	111.80
1	A	166	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	165	ASP	CB-CG-OD1	5.98	123.68	118.30
1	A	165	ASP	CB-CG-OD2	-5.42	113.42	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2319	0	2136	29	0
2	A	12	0	7	2	0
3	A	43	0	30	0	0
4	A	226	0	0	7	0
All	All	2600	0	2173	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:ASP:HB2	4:A:2163:HOH:O	1.35	1.27
1:A:184:ARG:HH22	2:A:1253:ASC:H61	1.40	0.87
1:A:257:ARG:NE	4:A:2193:HOH:O	2.19	0.74
1:A:166:ARG:NH2	1:A:250:GLU:OE1	2.21	0.74
1:A:216:ASN:HD22	1:A:222:GLN:HE21	1.42	0.68
1:A:166:ARG:HH22	1:A:250:GLU:CD	1.97	0.66
1:A:20:GLN:HE22	1:A:287:LYS:H	1.45	0.64
1:A:44:PRO:HD2	2:A:1253:ASC:O1	1.98	0.63
1:A:216:ASN:HD22	1:A:222:GLN:NE2	1.98	0.62
1:A:210:ASP:CG	4:A:2165:HOH:O	2.39	0.60
1:A:98:GLU:OE2	4:A:2084:HOH:O	2.17	0.57
1:A:6:HIS:HE1	4:A:2059:HOH:O	1.88	0.57
1:A:257:ARG:NH2	4:A:2193:HOH:O	2.38	0.56
1:A:257:ARG:CZ	4:A:2193:HOH:O	2.55	0.54
1:A:188:GLU:H	1:A:222:GLN:HE22	1.56	0.54
1:A:53:THR:HG22	1:A:71:TYR:HB2	1.90	0.53
1:A:29:LYS:HE2	1:A:91:PHE:O	2.11	0.50
1:A:99:PHE:O	1:A:102:ILE:HG22	2.12	0.49
1:A:84:GLY:HA3	1:A:184:ARG:HD3	1.97	0.47
1:A:20:GLN:HE22	1:A:287:LYS:N	2.13	0.46
1:A:67:TYR:HA	1:A:130:ARG:HG2	1.98	0.46
1:A:58:ASP:OD1	1:A:60:HIS:HD2	1.99	0.45
1:A:127:ARG:HG3	1:A:271:GLU:OE1	2.17	0.45
1:A:57:TRP:CD2	1:A:272:ASN:HB3	2.53	0.43
1:A:201:GLU:HG2	1:A:205:ASN:ND2	2.35	0.42
1:A:215:LYS:HD3	1:A:219:ASN:OD1	2.19	0.42
1:A:58:ASP:OD1	1:A:60:HIS:CD2	2.73	0.42
1:A:200:ASN:HD22	1:A:200:ASN:H	1.69	0.41
1:A:277:PRO:HD2	1:A:280:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/293 (100%)	287 (98%)	4 (1%)	1 (0%)	44	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/251 (94%)	231 (98%)	4 (2%)	66	70

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	TRP
1	A	77	PHE
1	A	200	ASN
1	A	257	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	159	GLN
1	A	200	ASN
1	A	208	ASN
1	A	222	GLN
1	A	292	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ASC	A	1253	-	12,12,12	1.89	1 (8%)	17,17,17	2.41	8 (47%)
3	HEM	A	1254	1,4	28,50,50	2.18	8 (28%)	17,82,82	1.82	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ASC	A	1253	-	-	0/6/22/22	0/1/1/1
3	HEM	A	1254	1,4	-	0/6/54/54	0/0/8/8

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1254	HEM	C3B-C2B	-3.70	1.35	1.40
3	A	1254	HEM	C3C-C2C	-2.60	1.36	1.40
3	A	1254	HEM	CMA-C3A	2.09	1.55	1.51
3	A	1254	HEM	C4B-NB	2.21	1.40	1.36
3	A	1254	HEM	C4D-ND	4.14	1.41	1.36
3	A	1254	HEM	C3B-CAB	4.22	1.56	1.47
3	A	1254	HEM	C3C-CAC	4.27	1.56	1.47
3	A	1254	HEM	C3D-C2D	4.83	1.52	1.37
2	A	1253	ASC	O4-C1	5.43	1.44	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1253	ASC	C1-C2-C3	-5.00	100.29	107.77
3	A	1254	HEM	CAA-CBA-CGA	-4.35	105.23	112.66
2	A	1253	ASC	O4-C4-C3	-3.41	101.48	104.03
3	A	1254	HEM	CMA-C3A-C4A	-3.29	123.41	128.46
2	A	1253	ASC	O3-C3-C2	-2.84	124.35	132.45
2	A	1253	ASC	C4-O4-C1	-2.59	106.57	109.25
2	A	1253	ASC	C5-C4-C3	2.28	117.24	114.23
3	A	1254	HEM	CMA-C3A-C2A	2.36	129.40	124.94
2	A	1253	ASC	O2-C2-C1	2.52	128.32	121.87
2	A	1253	ASC	O4-C1-C2	3.11	112.50	109.91
2	A	1253	ASC	C4-C3-C2	3.65	115.78	109.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1253	ASC	2	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/293 (100%)	1.59	84 (28%) 1 1	17, 26, 39, 53	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	THR	6.6
1	A	99	PHE	5.9
1	A	16	TYR	5.5
1	A	101	TRP	5.1
1	A	247	ILE	4.8
1	A	95	ILE	4.5
1	A	71	TYR	4.5
1	A	14	ARG	4.3
1	A	3	PRO	4.1
1	A	280	ALA	3.9
1	A	15	SER	3.9
1	A	206	LEU	3.9
1	A	182	LEU	3.9
1	A	8	ALA	3.8
1	A	194	ALA	3.7
1	A	10	VAL	3.7
1	A	293	GLY	3.5
1	A	265	ALA	3.5
1	A	252	ALA	3.3
1	A	285	ILE	3.3
1	A	218	ALA	3.2
1	A	96	HIS	3.2
1	A	256	ASP	3.2
1	A	23	TYR	3.2
1	A	207	LEU	3.2
1	A	217	ASP	3.2
1	A	77	PHE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	198	PHE	3.1
1	A	286	PHE	3.1
1	A	128	CYS	3.1
1	A	39	TYR	3.1
1	A	251	TYR	3.1
1	A	294	LEU	3.1
1	A	19	PHE	3.1
1	A	281	PRO	3.0
1	A	73	PHE	3.0
1	A	258	PHE	3.0
1	A	98	GLU	2.9
1	A	213	LEU	2.8
1	A	133	THR	2.8
1	A	100	PRO	2.8
1	A	277	PRO	2.8
1	A	13	GLY	2.8
1	A	169	VAL	2.7
1	A	291	GLU	2.7
1	A	276	PHE	2.7
1	A	259	PHE	2.7
1	A	244	TYR	2.6
1	A	180	THR	2.6
1	A	103	SER	2.6
1	A	22	VAL	2.6
1	A	170	ALA	2.5
1	A	131	VAL	2.4
1	A	20	GLN	2.4
1	A	116	VAL	2.4
1	A	246	SER	2.4
1	A	287	LYS	2.4
1	A	102	ILE	2.4
1	A	88	GLY	2.4
1	A	7	VAL	2.4
1	A	38	ASN	2.4
1	A	248	VAL	2.3
1	A	94	PRO	2.3
1	A	254	ASP	2.3
1	A	245	LEU	2.3
1	A	255	GLN	2.3
1	A	18	ASP	2.3
1	A	214	GLU	2.3
1	A	219	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	263	SER	2.2
1	A	132	ASP	2.2
1	A	203	TYR	2.2
1	A	70	THR	2.2
1	A	233	PRO	2.2
1	A	9	SER	2.1
1	A	211	TRP	2.1
1	A	26	ILE	2.1
1	A	134	PRO	2.1
1	A	72	ARG	2.0
1	A	126	TRP	2.0
1	A	220	ASN	2.0
1	A	257	ARG	2.0
1	A	283	PRO	2.0
1	A	173	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	ASC	A	1253	12/12	0.48	0.47	10.66	13,25,34,35	12
3	HEM	A	1254	43/43	0.91	0.16	-0.71	17,23,25,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.