



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 02:53 am GMT

PDB ID : 1X15
Title : Crystal structure of E. coli transhydrogenase domain I with bound NADH
Authors : Johansson, T.; Oswald, C.; Pedersen, A.; Tornroth, S.; Okvist, M.; Karlsson, B.G.; Rydstrom, J.; Krengel, U.
Deposited on : 2005-03-31
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1316 (2.04-2.04)
Clashscore	112137	1394 (2.04-2.04)
Ramachandran outliers	110173	1383 (2.04-2.04)
Sidechain outliers	110143	1383 (2.04-2.04)
RSRZ outliers	101464	1319 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	<div> <div>4%</div> <div> <div></div> <div>80%</div> <div>12%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	401	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>•</div> <div>10%</div> </div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD(P) transhydrogenase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	370	Total	C	N	O	S	0	0	0
			2764	1754	467	531	12			
1	B	360	Total	C	N	O	S	0	0	0
			2693	1707	458	517	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	994	MET	-	EXPRESSION TAG	UNP P07001
A	995	HIS	-	EXPRESSION TAG	UNP P07001
A	996	HIS	-	EXPRESSION TAG	UNP P07001
A	997	HIS	-	EXPRESSION TAG	UNP P07001
A	998	HIS	-	EXPRESSION TAG	UNP P07001
A	999	HIS	-	EXPRESSION TAG	UNP P07001
A	1000	HIS	-	EXPRESSION TAG	UNP P07001
A	1001	GLY	-	EXPRESSION TAG	UNP P07001
B	994	MET	-	EXPRESSION TAG	UNP P07001
B	995	HIS	-	EXPRESSION TAG	UNP P07001
B	996	HIS	-	EXPRESSION TAG	UNP P07001
B	997	HIS	-	EXPRESSION TAG	UNP P07001
B	998	HIS	-	EXPRESSION TAG	UNP P07001
B	999	HIS	-	EXPRESSION TAG	UNP P07001
B	1000	HIS	-	EXPRESSION TAG	UNP P07001
B	1001	GLY	-	EXPRESSION TAG	UNP P07001

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	104	Total	O	0	0
			104	104		
3	B	125	Total	O	0	0
			125	125		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD(P) transhydrogenase subunit alpha



- Molecule 1: NAD(P) transhydrogenase subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.73Å 67.09Å 76.84Å 67.22° 80.19° 81.60°	Depositor
Resolution (Å)	30.00 – 2.04 29.15 – 2.04	Depositor EDS
% Data completeness (in resolution range)	96.5 (30.00-2.04) 84.5 (29.15-2.04)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.04Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.186 , 0.241 0.195 , 0.243	Depositor DCC
R_{free} test set	2132 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 41.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5730	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/2807	0.90	11/3812 (0.3%)
1	B	0.83	1/2735 (0.0%)	0.88	10/3716 (0.3%)
All	All	0.81	1/5542 (0.0%)	0.89	21/7528 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1113	MET	CG-SD	-7.03	1.62	1.81

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1318	ARG	NE-CZ-NH2	-9.23	115.69	120.30
1	A	1047	ASP	CB-CG-OD2	7.81	125.33	118.30
1	A	1250	ASP	CB-CG-OD2	7.09	124.69	118.30
1	A	1078	ASP	CB-CG-OD2	7.04	124.64	118.30
1	A	1195	ASP	CB-CG-OD2	6.84	124.45	118.30
1	B	1343	ASP	CB-CG-OD2	6.69	124.32	118.30
1	A	1343	ASP	CB-CG-OD2	6.62	124.26	118.30
1	A	1318	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	A	1215	ASP	CB-CG-OD2	6.21	123.89	118.30
1	B	1047	ASP	CB-CG-OD2	6.19	123.88	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1351	ASP	CB-CG-OD2	6.16	123.85	118.30
1	A	1232	ASP	CB-CG-OD2	5.84	123.56	118.30
1	B	1116	ASP	CB-CG-OD2	5.76	123.49	118.30
1	B	1195	ASP	CB-CG-OD2	5.76	123.48	118.30
1	B	1269	ARG	NE-CZ-NH1	5.57	123.09	120.30
1	B	1197	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	B	1128	ASP	CB-CG-OD2	5.45	123.21	118.30
1	A	1077	ASP	CB-CG-OD2	5.33	123.10	118.30
1	B	1351	ASP	CB-CG-OD2	5.16	122.94	118.30
1	B	1250	ASP	CB-CG-OD2	5.15	122.93	118.30
1	B	1127	LEU	CB-CA-C	-5.08	100.54	110.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	1260	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2764	0	2817	15	0
1	B	2693	0	2746	23	0
2	B	44	0	26	0	0
3	A	104	0	0	3	0
3	B	125	0	0	5	0
All	All	5730	0	5589	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1151:ARG:NH2	3:B:149:HOH:O	1.90	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1151:ARG:CZ	3:B:149:HOH:O	2.10	0.98
1:B:1258:ILE:CG2	1:B:1258:ILE:O	2.39	0.71
1:B:1254:THR:HG22	1:B:1283:ASP:HA	1.79	0.65
1:B:1025:GLN:OE1	3:B:306:HOH:O	2.15	0.63
1:B:1258:ILE:O	1:B:1258:ILE:HG23	2.02	0.59
1:A:1276:LYS:NZ	3:A:129:HOH:O	2.35	0.57
1:B:1246:ALA:O	1:B:1275:MET:HG2	2.05	0.57
1:B:1100:GLU:CD	1:B:1100:GLU:H	2.08	0.55
1:B:1061:ASN:HB3	1:B:1082:LEU:HD13	1.89	0.54
1:B:1113:MET:HG2	1:B:1358:VAL:HG13	1.90	0.52
1:B:1061:ASN:ND2	3:B:37:HOH:O	2.43	0.52
1:A:1135:ASN:OD1	1:B:1156:GLN:NE2	2.44	0.51
1:B:1080:ILE:HD11	1:B:1101:LEU:HD12	1.94	0.49
1:B:1141:ALA:HB2	1:B:1284:LEU:HD11	1.94	0.48
1:B:1148:GLU:OE1	1:B:1308:LYS:NZ	2.44	0.48
1:B:1258:ILE:O	1:B:1259:PRO:C	2.52	0.48
1:A:1113:MET:HB3	1:A:1358:VAL:CG1	2.43	0.48
1:A:1064:TRP:HA	1:A:1069:ILE:HD11	1.95	0.47
1:A:1326:LEU:HD13	3:A:203:HOH:O	2.15	0.47
1:A:1079:GLU:HA	1:A:1082:LEU:HD12	1.97	0.47
1:A:1170:VAL:O	1:A:1193:ALA:HA	2.15	0.47
1:A:1287:GLN:NE2	3:A:107:HOH:O	2.48	0.46
1:B:1256:ALA:H	1:B:1291:ASN:HD21	1.64	0.45
1:A:1090:VAL:HG11	1:A:1331:LEU:HD22	1.98	0.44
1:B:1315:LEU:N	1:B:1316:PRO:CD	2.80	0.43
1:B:1254:THR:HG23	1:B:1291:ASN:HD22	1.83	0.43
1:A:1252:ILE:HG21	1:A:1267:ILE:HD11	2.01	0.43
1:A:1257:LEU:CD1	1:A:1288:ASN:HB2	2.49	0.43
1:A:1113:MET:SD	1:A:1338:LEU:HD21	2.59	0.43
1:A:1235:ILE:O	1:A:1239:MET:HG2	2.19	0.42
1:A:1080:ILE:O	1:A:1108:ARG:HD2	2.19	0.42
1:B:1235:ILE:O	1:B:1239:MET:HG2	2.21	0.41
1:A:1136:ILE:HD11	1:A:1326:LEU:HD12	2.02	0.41
1:B:1257:LEU:HD11	1:B:1288:ASN:OD1	2.21	0.41
1:B:1258:ILE:HG22	1:B:1258:ILE:O	2.19	0.41
1:B:1318:ARG:NH1	3:B:292:HOH:O	2.34	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/401 (91%)	357 (98%)	8 (2%)	1 (0%)	44	35
1	B	356/401 (89%)	346 (97%)	9 (2%)	1 (0%)	44	35
All	All	722/802 (90%)	703 (97%)	17 (2%)	2 (0%)	44	35

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1215	ASP
1	B	1260	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/317 (92%)	275 (94%)	18 (6%)	22	12
1	B	286/317 (90%)	270 (94%)	16 (6%)	25	15
All	All	579/634 (91%)	545 (94%)	34 (6%)	23	13

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1062	SER
1	A	1068	ILE
1	A	1101	LEU
1	A	1108	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1120	ARG
1	A	1153	PHE
1	A	1201	LYS
1	A	1202	GLU
1	A	1216	PHE
1	A	1228	LYS
1	A	1241	LEU
1	A	1308	LYS
1	A	1338	LEU
1	A	1341	GLU
1	A	1342	LYS
1	A	1354	VAL
1	A	1362	ARG
1	A	1365	GLU
1	B	1021	LYS
1	B	1028	LYS
1	B	1041	GLN
1	B	1061	ASN
1	B	1097	GLN
1	B	1100	GLU
1	B	1120	ARG
1	B	1162	LYS
1	B	1202	GLU
1	B	1232	ASP
1	B	1235	ILE
1	B	1258	ILE
1	B	1276	LYS
1	B	1287	GLN
1	B	1345	ASN
1	B	1351	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1135	ASN
1	A	1330	ASN
1	B	1012	ASN
1	B	1061	ASN
1	B	1156	GLN
1	B	1288	ASN
1	B	1291	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	B	1395	-	41,48,48	1.43	3 (7%)	43,73,73	1.91	7 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	1395	-	-	0/22/62/62	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1395	NAD	PA-O1A	2.20	1.59	1.50
2	B	1395	NAD	O4D-C1D	3.59	1.46	1.41
2	B	1395	NAD	C4A-N3A	5.89	1.44	1.35

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1395	NAD	N3A-C2A-N1A	-8.01	121.88	128.86
2	B	1395	NAD	C4A-C5A-N7A	-4.71	104.86	109.41
2	B	1395	NAD	C1B-N9A-C4A	-3.13	121.22	126.64
2	B	1395	NAD	C3N-C7N-N7N	2.20	120.29	117.77
2	B	1395	NAD	C2D-C3D-C4D	2.42	107.33	102.62
2	B	1395	NAD	C2A-N1A-C6A	2.85	123.76	118.77
2	B	1395	NAD	C2B-C3B-C4B	3.01	108.48	102.62

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	370/401 (92%)	0.10	15 (4%) 38 42	22, 37, 54, 70	0
1	B	360/401 (89%)	-0.05	12 (3%) 47 52	20, 33, 54, 72	0
All	All	730/802 (91%)	0.03	27 (3%) 42 47	20, 35, 54, 72	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1376	SER	5.2
1	B	1260	GLY	4.7
1	A	1215	ASP	4.6
1	A	1216	PHE	4.4
1	B	1258	ILE	4.4
1	B	1215	ASP	4.3
1	A	1375	VAL	3.8
1	B	1259	PRO	3.7
1	A	1374	GLN	3.1
1	A	1232	ASP	3.1
1	A	1122	SER	3.1
1	A	1060	GLY	2.8
1	B	1232	ASP	2.8
1	A	1090	VAL	2.7
1	A	1120	ARG	2.7
1	B	1231	SER	2.6
1	B	1257	LEU	2.5
1	A	1070	LEU	2.4
1	B	1233	ALA	2.4
1	B	1289	GLY	2.4
1	A	1121	ILE	2.3
1	B	1090	VAL	2.2
1	B	1373	ILE	2.2
1	A	1331	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1260	GLY	2.1
1	B	1120	ARG	2.1
1	A	1236	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAD	B	1395	44/44	0.88	0.18	0.63	54,61,83,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.