



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 14, 2017 – 11:19 am GMT

PDB ID : 3X1K
Title : crystal structure of Phosphoapantetheine adenylyltransferase PPAT/CoaD with AMP-PNP from Pseudomonas aeruginosa
Authors : Chatterjee, R.; Datta, S.
Deposited on : 2014-11-20
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4993 (2.58-2.50)
Clashscore	112137	5755 (2.58-2.50)
Ramachandran outliers	110173	5652 (2.58-2.50)
Sidechain outliers	110143	5654 (2.58-2.50)
RSRZ outliers	101464	5026 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	<div> <div>79%</div> <div>19%</div> <div>.</div> </div>
1	B	159	<div> <div>79%</div> <div>18%</div> <div>..</div> </div>
1	C	159	<div> <div>79%</div> <div>19%</div> <div>..</div> </div>
1	D	159	<div> <div>82%</div> <div>14%</div> <div>..</div> </div>
1	E	159	<div> <div>82%</div> <div>15%</div> <div>..</div> </div>
1	F	159	<div> <div>83%</div> <div>15%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DMS	B	1002	-	-	-	X
3	DMS	D	1003	-	-	-	X
3	DMS	E	1002	-	-	-	X
3	DMS	E	1004	-	-	-	X
3	DMS	E	1006	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7865 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphopantetheine adenylyltransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	157	Total	C	N	O	S	0	1	0
			1244	800	216	226	2			
1	B	157	Total	C	N	O	S	0	1	0
			1244	800	216	226	2			
1	C	157	Total	C	N	O	S	0	0	0
			1236	796	214	224	2			
1	D	158	Total	C	N	O	S	0	0	0
			1244	801	215	225	3			
1	E	157	Total	C	N	O	S	0	0	0
			1236	796	214	224	2			
1	F	158	Total	C	N	O	S	0	1	0
			1254	807	218	226	3			

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	C	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	E	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	F	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		
3	B	1	Total	C	O	S	0	0
			4	2	1	1		
3	D	1	Total	C	O	S	0	0
			4	2	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	O	S	0	0
			4	2	1	1		
3	E	1	Total	C	O	S	0	0
			4	2	1	1		
3	D	1	Total	C	O	S	0	0
			4	2	1	1		
3	E	1	Total	C	O	S	0	0
			4	2	1	1		
3	F	1	Total	C	O	S	0	0
			4	2	1	1		
3	F	1	Total	C	O	S	0	0
			4	2	1	1		
3	E	1	Total	C	O	S	0	0
			4	2	1	1		
3	E	1	Total	C	O	S	0	0
			4	2	1	1		
3	F	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		
4	D	1	Total	C	O	0	0
			3	1	2		

- Molecule 5 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		
5	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			6	3	3		
6	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	23	Total	O	0	0
			23	23		
7	B	31	Total	O	0	0
			31	31		
7	C	19	Total	O	0	0
			19	19		
7	D	21	Total	O	0	0
			21	21		
7	E	14	Total	O	0	0
			14	14		
7	F	9	Total	O	0	0
			9	9		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphopantetheine adenylyltransferase




- Molecule 1: Phosphopantetheine adenylyltransferase




- Molecule 1: Phosphopantetheine adenylyltransferase




- Molecule 1: Phosphopantetheine adenylyltransferase



- Molecule 1: Phosphopantetheine adenylyltransferase



- Molecule 1: Phosphopantetheine adenylyltransferase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.33Å 89.79Å 80.31Å 90.00° 98.27° 90.00°	Depositor
Resolution (Å)	39.74 – 2.55 39.74 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.2 (39.74-2.55) 91.0 (39.74-2.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.06 (at 2.54Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.190 , 0.245 0.190 , 0.246	Depositor DCC
R_{free} test set	2012 reflections (5.52%)	DCC
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.206	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 16.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.477 for l,-k,h	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7865	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FMT, DMS, ANP, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1271	0.64	0/1722
1	B	0.51	0/1271	0.63	0/1722
1	C	0.49	0/1263	0.64	0/1710
1	D	0.46	0/1271	0.63	1/1720 (0.1%)
1	E	0.46	0/1263	0.62	0/1710
1	F	0.47	0/1282	0.60	0/1736
All	All	0.48	0/7621	0.63	1/10320 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	153	LEU	CA-CB-CG	5.86	128.78	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1244	0	1255	19	0
1	B	1244	0	1255	19	0
1	C	1236	0	1249	21	0
1	D	1244	0	1261	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1236	0	1249	16	0
1	F	1254	0	1268	18	0
2	A	31	0	11	1	0
2	B	31	0	11	0	0
2	C	31	0	11	1	0
2	D	31	0	10	1	0
2	E	31	0	11	0	0
2	F	31	0	11	0	0
3	A	8	0	12	2	0
3	B	12	0	18	1	0
3	C	4	0	6	2	0
3	D	8	0	12	0	0
3	E	20	0	30	1	0
3	F	12	0	18	2	0
4	A	3	0	1	0	0
4	B	12	0	4	0	0
4	D	3	0	1	0	0
5	A	5	0	0	0	0
5	C	5	0	0	0	0
6	C	12	0	16	1	0
7	A	23	0	0	0	0
7	B	31	0	0	0	0
7	C	19	0	0	0	0
7	D	21	0	0	0	0
7	E	14	0	0	0	0
7	F	9	0	0	0	0
All	All	7865	0	7720	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:VAL:HB	1:E:96:GLU:OE1	1.83	0.77
1:F:57:VAL:HG13	1:F:146[B]:HIS:HD2	1.51	0.76
1:B:92:VAL:HB	1:F:96:GLU:OE1	1.91	0.71
1:F:127:SER:HB3	1:F:130:LEU:HB2	1.73	0.70
1:C:40:LYS:NZ	1:F:133:GLU:OE2	2.22	0.66

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	156/159 (98%)	154 (99%)	0	2 (1%)	14	24
1	B	156/159 (98%)	153 (98%)	1 (1%)	2 (1%)	14	24
1	C	155/159 (98%)	154 (99%)	1 (1%)	0	100	100
1	D	156/159 (98%)	155 (99%)	1 (1%)	0	100	100
1	E	155/159 (98%)	151 (97%)	4 (3%)	0	100	100
1	F	157/159 (99%)	155 (99%)	2 (1%)	0	100	100
All	All	935/954 (98%)	922 (99%)	9 (1%)	4 (0%)	51	57

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42[A]	ASN
1	A	42[B]	ASN
1	B	42[A]	ASN
1	B	42[B]	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	134/135 (99%)	127 (95%)	7 (5%)	27	47
1	B	134/135 (99%)	128 (96%)	6 (4%)	32	54
1	C	133/135 (98%)	127 (96%)	6 (4%)	32	54

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	134/135 (99%)	126 (94%)	8 (6%)	22	39
1	E	133/135 (98%)	125 (94%)	8 (6%)	22	39
1	F	135/135 (100%)	129 (96%)	6 (4%)	33	55
All	All	803/810 (99%)	762 (95%)	41 (5%)	28	48

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	158	LYS
1	D	92	VAL
1	F	53	LEU
1	D	14	THR
1	D	27	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

32 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ANP	A	1001	-	29,33,33	4.07	17 (58%)	28,52,52	2.86	7 (25%)
3	DMS	A	1002	-	3,3,3	0.61	0	3,3,3	0.12	0
4	FMT	A	1003	-	0,2,2	0.00	-	0,1,1	0.00	-
5	PO4	A	1004	-	4,4,4	0.74	0	6,6,6	0.58	0
3	DMS	A	1005	-	3,3,3	0.62	0	3,3,3	0.54	0
2	ANP	B	1001	-	29,33,33	4.13	17 (58%)	28,52,52	2.95	6 (21%)
3	DMS	B	1002	-	3,3,3	0.62	0	3,3,3	0.69	0
3	DMS	B	1003	-	3,3,3	0.66	0	3,3,3	0.31	0
3	DMS	B	1004	-	3,3,3	0.67	0	3,3,3	0.54	0
4	FMT	B	1005	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1006	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1007	-	0,2,2	0.00	-	0,1,1	0.00	-
4	FMT	B	1008	-	0,2,2	0.00	-	0,1,1	0.00	-
2	ANP	C	1001	-	29,33,33	3.95	16 (55%)	28,52,52	3.03	8 (28%)
3	DMS	C	1002	-	3,3,3	0.59	0	3,3,3	1.32	0
5	PO4	C	1003	-	4,4,4	0.80	0	6,6,6	0.74	0
6	GOL	C	1004	-	5,5,5	0.66	0	5,5,5	1.18	0
6	GOL	C	1005	-	5,5,5	0.40	0	5,5,5	0.43	0
2	ANP	D	1001	-	29,33,33	4.30	16 (55%)	28,52,52	2.96	8 (28%)
3	DMS	D	1002	-	3,3,3	0.71	0	3,3,3	0.23	0
3	DMS	D	1003	-	3,3,3	0.61	0	3,3,3	0.71	0
4	FMT	D	1004	-	0,2,2	0.00	-	0,1,1	0.00	-
2	ANP	E	1001	-	29,33,33	4.20	15 (51%)	28,52,52	3.02	10 (35%)
3	DMS	E	1002	-	3,3,3	0.59	0	3,3,3	0.42	0
3	DMS	E	1003	-	3,3,3	0.65	0	3,3,3	0.67	0
3	DMS	E	1004	-	3,3,3	0.59	0	3,3,3	0.75	0
3	DMS	E	1005	-	3,3,3	0.71	0	3,3,3	0.82	0
3	DMS	E	1006	-	3,3,3	0.69	0	3,3,3	0.40	0
2	ANP	F	1001	-	29,33,33	4.10	16 (55%)	28,52,52	2.96	10 (35%)
3	DMS	F	1002	-	3,3,3	0.59	0	3,3,3	0.35	0
3	DMS	F	1003	-	3,3,3	0.62	0	3,3,3	0.46	0
3	DMS	F	1004	-	3,3,3	0.52	0	3,3,3	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ANP	A	1001	-	-	0/13/38/38	0/3/3/3
3	DMS	A	1002	-	-	0/0/0/0	0/0/0/0
4	FMT	A	1003	-	-	0/0/0/0	0/0/0/0
5	PO4	A	1004	-	-	0/0/0/0	0/0/0/0
3	DMS	A	1005	-	-	0/0/0/0	0/0/0/0
2	ANP	B	1001	-	-	0/13/38/38	0/3/3/3
3	DMS	B	1002	-	-	0/0/0/0	0/0/0/0
3	DMS	B	1003	-	-	0/0/0/0	0/0/0/0
3	DMS	B	1004	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1005	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1006	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1007	-	-	0/0/0/0	0/0/0/0
4	FMT	B	1008	-	-	0/0/0/0	0/0/0/0
2	ANP	C	1001	-	-	0/13/38/38	0/3/3/3
3	DMS	C	1002	-	-	0/0/0/0	0/0/0/0
5	PO4	C	1003	-	-	0/0/0/0	0/0/0/0
6	GOL	C	1004	-	-	0/4/4/4	0/0/0/0
6	GOL	C	1005	-	-	0/4/4/4	0/0/0/0
2	ANP	D	1001	-	-	0/13/38/38	0/3/3/3
3	DMS	D	1002	-	-	0/0/0/0	0/0/0/0
3	DMS	D	1003	-	-	0/0/0/0	0/0/0/0
4	FMT	D	1004	-	-	0/0/0/0	0/0/0/0
2	ANP	E	1001	-	-	1/13/38/38	0/3/3/3
3	DMS	E	1002	-	-	0/0/0/0	0/0/0/0
3	DMS	E	1003	-	-	0/0/0/0	0/0/0/0
3	DMS	E	1004	-	-	0/0/0/0	0/0/0/0
3	DMS	E	1005	-	-	0/0/0/0	0/0/0/0
3	DMS	E	1006	-	-	0/0/0/0	0/0/0/0
2	ANP	F	1001	-	-	0/13/38/38	0/3/3/3
3	DMS	F	1002	-	-	0/0/0/0	0/0/0/0
3	DMS	F	1003	-	-	0/0/0/0	0/0/0/0
3	DMS	F	1004	-	-	0/0/0/0	0/0/0/0

The worst 5 of 97 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1001	ANP	C2'-C3'	-12.00	1.21	1.53
2	F	1001	ANP	C2'-C3'	-11.98	1.21	1.53
2	B	1001	ANP	C2'-C3'	-11.68	1.22	1.53
2	C	1001	ANP	C2'-C3'	-11.65	1.22	1.53
2	A	1001	ANP	C2'-C3'	-11.52	1.22	1.53

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	ANP	C1'-N9-C4	-10.15	109.09	126.64
2	E	1001	ANP	C1'-N9-C4	-10.02	109.32	126.64
2	C	1001	ANP	C1'-N9-C4	-9.86	109.60	126.64
2	B	1001	ANP	C1'-N9-C4	-9.65	109.96	126.64
2	D	1001	ANP	C1'-N9-C4	-9.26	110.64	126.64

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	1001	ANP	O1B-PB-N3B-PG

There are no ring outliers.

10 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	ANP	1	0
3	A	1002	DMS	1	0
3	A	1005	DMS	1	0
3	B	1004	DMS	1	0
2	C	1001	ANP	1	0
3	C	1002	DMS	2	0
6	C	1004	GOL	1	0
2	D	1001	ANP	1	0
3	E	1003	DMS	1	0
3	F	1004	DMS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	157/159 (98%)	-0.52	0 100 100	19, 30, 43, 52	0
1	B	157/159 (98%)	-0.51	0 100 100	20, 29, 44, 52	0
1	C	157/159 (98%)	-0.46	0 100 100	23, 33, 48, 65	0
1	D	158/159 (99%)	-0.51	0 100 100	21, 34, 54, 63	0
1	E	157/159 (98%)	-0.46	0 100 100	25, 36, 50, 70	0
1	F	158/159 (99%)	-0.46	0 100 100	24, 36, 51, 69	0
All	All	944/954 (98%)	-0.49	0 100 100	19, 33, 48, 70	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	DMS	E	1002	4/4	0.94	0.29	5.96	30,37,38,38	4
3	DMS	B	1002	4/4	0.98	0.20	4.58	31,33,37,43	0
3	DMS	D	1003	4/4	0.95	0.19	3.38	41,47,51,54	0
3	DMS	E	1004	4/4	0.99	0.17	2.40	38,39,40,59	0
3	DMS	E	1006	4/4	0.99	0.18	2.33	32,37,48,50	0
3	DMS	F	1003	4/4	0.97	0.17	1.95	30,31,36,45	0
3	DMS	E	1005	4/4	0.97	0.17	1.86	41,43,43,54	0
3	DMS	B	1003	4/4	0.98	0.15	1.64	27,37,42,43	0
3	DMS	D	1002	4/4	0.99	0.15	1.37	31,32,33,38	0
3	DMS	E	1003	4/4	0.98	0.22	1.19	33,50,52,57	0
3	DMS	F	1002	4/4	1.00	0.18	1.17	33,37,43,44	0
4	FMT	A	1003	3/3	0.90	0.17	0.63	32,32,45,53	0
3	DMS	A	1002	4/4	0.96	0.16	0.46	38,45,50,52	0
2	ANP	E	1001	31/31	0.93	0.13	0.30	31,40,63,72	0
2	ANP	D	1001	31/31	0.94	0.13	0.30	26,32,51,70	0
6	GOL	C	1004	6/6	0.93	0.14	0.21	25,30,42,43	0
3	DMS	F	1004	4/4	0.97	0.14	0.19	42,46,56,63	0
3	DMS	C	1002	4/4	0.97	0.14	0.14	36,39,46,54	0
2	ANP	F	1001	31/31	0.94	0.12	0.06	32,39,59,66	0
2	ANP	C	1001	31/31	0.94	0.13	-0.06	26,33,52,62	0
2	ANP	B	1001	31/31	0.96	0.12	-0.12	25,31,51,56	0
5	PO4	C	1003	5/5	0.99	0.14	-0.19	39,43,51,60	0
4	FMT	B	1005	3/3	0.97	0.12	-0.23	39,39,42,48	0
3	DMS	B	1004	4/4	0.99	0.12	-0.43	35,36,52,53	0
3	DMS	A	1005	4/4	0.97	0.12	-0.60	40,46,50,51	0
4	FMT	B	1006	3/3	0.96	0.13	-0.68	37,37,43,50	0
6	GOL	C	1005	6/6	0.87	0.12	-0.77	47,49,53,55	0
5	PO4	A	1004	5/5	0.96	0.12	-0.81	48,48,63,75	0
2	ANP	A	1001	31/31	0.97	0.10	-0.97	25,31,51,55	0
4	FMT	B	1007	3/3	0.97	0.08	-1.49	43,43,47,49	0
4	FMT	D	1004	3/3	0.94	0.10	-	49,49,50,53	0
4	FMT	B	1008	3/3	0.98	0.10	-	33,33,34,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.