



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2017 – 03:23 PM EDT

PDB ID : 5X2D  
Title : Crystal structure of DLC like domain of CsTAL3 (83-177aa)  
Authors : Jo, C.H.; Hwang, K.Y.  
Deposited on : unknown  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2542 (2.60-2.60)
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	95	<div> <div>9%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>
1	B	95	<div> <div>4%</div> <div>73%</div> <div>26%</div> <div>.</div> </div>
1	C	95	<div> <div>18%</div> <div>78%</div> <div>21%</div> <div>.</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tegumental protein 20.8 kDa.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	95	Total	C	N	O	S	0	0	0
			760	491	120	145	4			
1	B	95	Total	C	N	O	S	0	0	0
			760	491	120	145	4			
1	C	95	Total	C	N	O	S	0	0	0
			760	491	120	145	4			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	O	0	0
			6	6		
2	B	5	Total	O	0	0
			5	5		
2	C	8	Total	O	0	0
			8	8		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Tegumental protein 20.8 kDa



- Molecule 1: Tegumental protein 20.8 kDa



- Molecule 1: Tegumental protein 20.8 kDa



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.40Å 154.08Å 61.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 2.60 48.13 – 2.60	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.13-2.60) 93.6 (48.13-2.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.8.3_1479	Depositor
R, $R_{free}$	0.224 , 0.267 0.223 , 0.270	Depositor DCC
$R_{free}$ test set	1277 reflections (10.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	61.9	Xtriage
Anisotropy	0.644	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 66.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.458 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.439 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2299	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	96.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/780	0.73	0/1061
1	B	0.50	0/780	0.75	2/1061 (0.2%)
1	C	0.48	0/780	0.73	0/1061
All	All	0.48	0/2340	0.74	2/3183 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	116	GLN	CA-CB-CG	-5.51	101.28	113.40
1	B	116	GLN	N-CA-C	5.24	125.13	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	117	THR	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	760	0	739	11	0
1	B	760	0	739	13	0
1	C	760	0	739	15	0
2	A	6	0	0	0	1
2	B	5	0	0	0	0
2	C	8	0	0	0	1
All	All	2299	0	2217	37	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:GLN:HE21	1:A:116:GLN:HA	1.38	0.87
1:C:123:ARG:O	1:C:127:ASN:ND2	2.15	0.80
1:A:115:ALA:HB1	1:A:120:GLU:HB3	1.66	0.76
1:A:152:THR:HG21	1:C:129:LYS:HB2	1.72	0.71
1:A:129:LYS:HB2	1:C:152:THR:HG21	1.74	0.68
1:C:115:ALA:HA	1:C:120:GLU:HG2	1.79	0.65
1:A:116:GLN:HE21	1:A:116:GLN:CA	2.13	0.57
1:A:125:THR:HG22	1:A:144:VAL:HG23	1.89	0.55
1:C:129:LYS:HG3	1:C:140:TRP:O	2.07	0.54
1:B:129:LYS:HG3	1:B:140:TRP:O	2.09	0.53
1:C:117:THR:N	1:C:120:GLU:OE1	2.40	0.53
1:B:101:GLN:HG2	1:B:161:PHE:CZ	2.43	0.52
1:A:129:LYS:HG3	1:A:140:TRP:O	2.10	0.52
1:A:116:GLN:N	1:A:120:GLU:OE1	2.39	0.51
1:C:125:THR:HG22	1:C:144:VAL:HG23	1.93	0.50
1:A:117:THR:N	1:A:120:GLU:OE1	2.41	0.48
1:B:117:THR:O	1:B:120:GLU:HG2	2.12	0.48
1:B:112:ALA:HB1	1:B:170:TYR:OH	2.14	0.48
1:B:115:ALA:HB1	1:B:120:GLU:HG3	1.97	0.47
1:C:116:GLN:H	1:C:120:GLU:CD	2.17	0.47
1:B:89:ILE:HG23	1:B:163:PHE:HB2	1.98	0.46
1:B:138:GLY:HA3	1:B:176:SER:O	2.17	0.45
1:A:138:GLY:HA3	1:A:176:SER:O	2.17	0.44
1:B:158:ASN:HB2	1:B:174:GLN:O	2.18	0.44
1:B:89:ILE:HD13	1:B:105:ALA:HB1	2.00	0.44
1:C:123:ARG:HA	1:C:126:GLU:HB2	2.00	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:102:ILE:O	1:C:105:ALA:HB3	2.18	0.43
1:B:109:ILE:HA	1:B:165:LEU:HD21	2.00	0.43
1:C:121:LEU:HA	1:C:121:LEU:HD23	1.80	0.43
1:C:88:GLU:H	1:C:88:GLU:HG2	1.59	0.43
1:C:151:ILE:HG22	1:C:152:THR:O	2.19	0.42
1:B:96:MET:HE2	1:B:159:MET:HB3	2.02	0.41
1:B:143:MET:CE	1:B:171:LEU:HD22	2.50	0.41
1:A:107:GLU:HG2	1:A:131:PHE:CG	2.56	0.41
1:B:127:ASN:HA	1:B:130:SER:HB3	2.01	0.41
1:C:107:GLU:HG2	1:C:131:PHE:CG	2.56	0.40
1:C:92:ILE:N	1:C:162:GLN:O	2.48	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:202:HOH:O	2:C:202:HOH:O[6_445]	2.06	0.14

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	93/95 (98%)	92 (99%)	1 (1%)	0	100	100
1	B	93/95 (98%)	92 (99%)	1 (1%)	0	100	100
1	C	93/95 (98%)	90 (97%)	3 (3%)	0	100	100
All	All	279/285 (98%)	274 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/84 (100%)	82 (98%)	2 (2%)	54	80
1	B	84/84 (100%)	83 (99%)	1 (1%)	75	91
1	C	84/84 (100%)	83 (99%)	1 (1%)	75	91
All	All	252/252 (100%)	248 (98%)	4 (2%)	68	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	106	ARG
1	A	116	GLN
1	B	98	LEU
1	C	120	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	C	127	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	95/95 (100%)	0.57	9 (9%) <b>9</b> <b>5</b>	68, 91, 127, 178	0
1	B	95/95 (100%)	0.39	4 (4%) <b>37</b> <b>29</b>	64, 90, 146, 175	0
1	C	95/95 (100%)	0.89	17 (17%) <b>2</b> <b>1</b>	62, 90, 136, 178	0
All	All	285/285 (100%)	0.62	30 (10%) <b>7</b> <b>4</b>	62, 91, 139, 178	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	112	ALA	6.8
1	A	170	TYR	6.5
1	C	111	ILE	5.5
1	B	106	ARG	4.1
1	C	113	PRO	3.9
1	C	136	PHE	3.9
1	B	122	GLY	3.5
1	C	145	VAL	3.5
1	C	170	TYR	3.3
1	A	146	ASP	3.0
1	A	145	VAL	3.0
1	C	146	ASP	2.9
1	C	109	ILE	2.8
1	C	166	TYR	2.7
1	C	115	ALA	2.6
1	C	108	THR	2.6
1	A	172	PHE	2.6
1	C	148	SER	2.5
1	B	85	LEU	2.4
1	C	106	ARG	2.4
1	A	163	PHE	2.4
1	C	158	ASN	2.4
1	A	144	VAL	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	113	PRO	2.3
1	C	169	ALA	2.2
1	A	168	ARG	2.1
1	B	170	TYR	2.0
1	C	98	LEU	2.0
1	A	149	TYR	2.0
1	C	172	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.