



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 16, 2018 – 04:34 PM EST

PDB ID : 2X3E  
Title : Crystal structure of 3-oxoacyl-(acyl carrier protein) synthase III, FabH from *Pseudomonas aeruginosa* PAO1  
Authors : Oke, M.; Carter, L.G.; Johnson, K.A.; Liu, H.; McMahon, S.A.; White, M.F.; Naismith, J.H.  
Deposited on : 2010-01-24  
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5868 (1.84-1.80)
Clashscore	112137	6856 (1.84-1.80)
Ramachandran outliers	110173	6780 (1.84-1.80)
Sidechain outliers	110143	6780 (1.84-1.80)
RSRZ outliers	101464	5947 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>9%</div> <div></div> </div> </div>
1	B	331	<div> <div>3%</div> <div> <div></div> <div>89%</div> <div>7%</div> <div></div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5195 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C	N	O	S	0	14	0
			2415	1510	430	465	10			
1	B	317	Total	C	N	O	S	0	4	0
			2316	1451	417	439	9			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q9HYR2
A	111	ALA	CYS	engineered mutation	UNP Q9HYR2
B	0	GLY	-	expression tag	UNP Q9HYR2
B	111	ALA	CYS	engineered mutation	UNP Q9HYR2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	241	Total	O	0	0
			241	241		
2	B	223	Total	O	0	0
			223	223		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 3



#### • Molecule 1: 3-OXOACYL-[ACYL-CARRIER-PROTEIN] SYNTHASE 3



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.24Å 69.24Å 212.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	213.20 – 1.81 28.99 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.2 (213.20-1.81) 99.2 (28.99-1.81)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0090	Depositor
R, $R_{free}$	0.169 , 0.204 0.194 , 0.230	Depositor DCC
$R_{free}$ test set	2647 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.0	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.076 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5195	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/2494	0.56	1/3392 (0.0%)
1	B	0.35	0/2364	0.55	0/3217
All	All	0.34	0/4858	0.56	1/6609 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	103	LEU	CA-CB-CG	5.07	126.97	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2415	0	2432	20	0
1	B	2316	0	2326	15	0
2	A	241	0	0	0	0
2	B	223	0	0	3	0
All	All	5195	0	4758	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:PRO:HG2	1:A:256:LEU:HD22	1.77	0.67
1:A:243:LEU:HD21	1:A:305:VAL:HG23	1.80	0.62
1:A:115:VAL:HG13	1:A:318[A]:VAL:HG23	1.81	0.62
1:A:225[B]:MET:HE2	1:A:307:THR:HG23	1.83	0.61
1:A:248:PRO:HG2	1:A:256:LEU:CD2	2.31	0.60
1:B:77:LEU:HD22	1:B:136:VAL:HB	1.85	0.57
1:A:179:PHE:CD1	1:A:318[A]:VAL:HG22	2.41	0.55
1:B:192:MET:HG3	1:B:193:THR:N	2.26	0.51
1:A:225[B]:MET:CE	1:A:307:THR:HG23	2.41	0.50
1:B:267:VAL:HG13	2:B:2189:HOH:O	2.11	0.50
1:A:44:ARG:N	1:A:44:ARG:HD2	2.26	0.50
1:A:77:LEU:HD22	1:A:136:VAL:HB	1.93	0.49
1:A:225[A]:MET:HE2	1:A:259:VAL:HG21	1.95	0.49
1:B:179:PHE:CD1	1:B:318:VAL:HG22	2.48	0.49
1:B:230:ARG:HH22	1:B:262:GLN:HG3	1.78	0.49
1:A:129:LEU:HD22	1:B:123:SER:OG	2.13	0.48
1:B:208:ASN:N	2:B:2143:HOH:O	2.48	0.47
1:B:252:ASN:HA	1:B:274:ILE:CD1	2.45	0.47
1:B:220:GLN:HA	1:B:223:THR:HG22	1.96	0.47
1:A:192:MET:HE3	1:A:194:PRO:HG3	1.98	0.46
1:A:225[A]:MET:HE1	1:A:248:PRO:HB2	1.97	0.46
1:B:115:VAL:HG13	1:B:318:VAL:HG23	1.98	0.46
1:A:35:ILE:HG23	1:A:154:ALA:HB1	1.98	0.45
1:B:326:VAL:HG23	2:B:2218:HOH:O	2.16	0.44
1:A:192:MET:HE1	1:A:211:ARG:NH2	2.33	0.44
1:A:36:SER:O	1:A:40:GLY:N	2.46	0.44
1:A:179:PHE:CE1	1:A:318[A]:VAL:HG22	2.53	0.44
1:B:179:PHE:CE1	1:B:318:VAL:HG22	2.54	0.43
1:B:225:MET:HE2	1:B:259:VAL:HG21	2.01	0.43
1:B:312:GLY:N	1:B:313:LEU:HA	2.35	0.41
1:A:223:THR:HG22	1:A:227:ASP:OD2	2.22	0.40
1:B:230:ARG:NH2	1:B:262:GLN:HG3	2.37	0.40
1:A:69:LEU:HD12	1:A:164:VAL:HG11	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/331 (102%)	329 (98%)	7 (2%)	1 (0%)	44	28
1	B	317/331 (96%)	309 (98%)	7 (2%)	1 (0%)	44	28
All	All	654/662 (99%)	638 (98%)	14 (2%)	2 (0%)	44	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	281	VAL
1	B	281	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/241 (104%)	247 (99%)	3 (1%)	75	69
1	B	234/241 (97%)	232 (99%)	2 (1%)	82	77
All	All	484/482 (100%)	479 (99%)	5 (1%)	78	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ARG
1	A	84	PHE
1	A	205	GLN
1	B	44	ARG
1	B	232	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	325/331 (98%)	0.15	18 (5%) 26 21	13, 19, 40, 61	0
1	B	317/331 (95%)	-0.01	10 (3%) 48 43	12, 18, 37, 49	0
All	All	642/662 (96%)	0.07	28 (4%) 35 29	12, 19, 39, 61	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	326	VAL	7.1
1	B	327	PRO	6.9
1	A	34	TRP	5.6
1	A	36	SER	5.5
1	A	203	SER	5.3
1	A	202	SER	4.9
1	A	33	ALA	4.4
1	A	199	ALA	3.9
1	A	204	GLY	3.9
1	A	205	GLN	3.7
1	A	35	ILE	3.4
1	A	38	ARG	3.4
1	A	32	ASP	3.2
1	A	218	PHE	3.1
1	A	198	ARG	3.1
1	B	199	ALA	3.0
1	B	208	ASN	3.0
1	A	37	SER	2.8
1	B	325	ILE	2.6
1	B	38	ARG	2.5
1	A	42	ARG	2.4
1	A	215	LYS	2.3
1	A	327	PRO	2.1
1	A	169[A]	ASP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	42	ARG	2.1
1	B	33	ALA	2.1
1	B	254	ARG	2.0
1	B	150	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.