



# Full wwPDB X-ray Structure Validation Report i

Nov 15, 2017 – 09:35 PM EST

PDB ID : 4X7R  
Title : Crystal structure of S. aureus TarM G117R mutant in complex with Fondaparinux, alpha-GlcNAc-glycerol and UDP  
Authors : Worrall, L.J.; Sobhanifar, S.; Strynadka, N.C.  
Deposited on : unknown  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	3YW	A	503	-	-	-	X
4	3YW	B	503	-	-	-	X

## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 8717 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TarM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	493	4040	2585	684	752	19	0	0	0
1	B	493	4040	2585	684	752	19	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	117	ARG	GLY	engineered mutation	UNP H0AM96
B	117	ARG	GLY	engineered mutation	UNP H0AM96

- Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: C<sub>9</sub>H<sub>14</sub>N<sub>2</sub>O<sub>12</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	25	9	2	12	2	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	25	9	2	12	2	0	0

- Molecule 3 is TRISULFOAMINO HEPARIN PENTASACCHARIDE (three-letter code: NTO) (formula: C<sub>31</sub>H<sub>53</sub>N<sub>3</sub>O<sub>49</sub>S<sub>8</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	91	31	3	49	8	0	0
3	B	1	91	31	3	49	8	0	0

- Molecule 4 is (2S)-2,3-dihydroxypropyl 2-(acetylamino)-2-deoxy-alpha-D-glucopyranoside (three-letter code: 3YW) (formula: C<sub>11</sub>H<sub>21</sub>NO<sub>8</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			20	11	1	8		

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	183	Total	O			0	0
			183	183				


  

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	182	Total	O			0	0
			182	182				


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TarM



- Molecule 1: TarM



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.47 Å    92.14 Å    96.11 Å 65.89°    83.66°    84.12°	Depositor
Resolution (Å)	87.48 – 2.15 87.48 – 2.15	Depositor EDS
% Data completeness (in resolution range)	97.9 (87.48-2.15) 85.1 (87.48-2.15)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.93 (at 2.14 Å)	Xtriage
Refinement program	BUSTER	Depositor
$R$ , $R_{free}$	0.179 , 0.200 0.181 , 0.202	Depositor DCC
$R_{free}$ test set	3707 reflections (5.44%)	DCC
Wilson B-factor (Å <sup>2</sup> )	38.4	Xtriage
Anisotropy	0.351	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 54.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.106 for -h,-l,-k	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8717	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: UDP, 3YW, NTO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.49	0/4117	0.66	0/5525
1	B	0.49	0/4117	0.66	0/5525
All	All	0.49	0/8234	0.66	0/11050

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4040	0	4067	16	0
1	B	4040	0	4067	17	0
2	A	25	0	11	1	0
2	B	25	0	11	0	0
3	A	91	0	51	1	0
3	B	91	0	51	2	0
4	A	20	0	21	3	0
4	B	20	0	21	2	0
5	A	183	0	0	1	0
5	B	182	0	0	2	0
All	All	8717	0	8300	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:VAL:HG21	1:B:449:MET:HE2	1.57	0.87
1:A:339:VAL:HG21	1:A:449:MET:HE2	1.57	0.87
1:A:343:VAL:HG12	1:A:350:VAL:HG11	1.72	0.71
1:B:343:VAL:HG12	1:B:350:VAL:HG11	1.72	0.69
1:B:18:MET:HB2	4:B:503:3YW:H2	1.79	0.63
1:B:306:VAL:CG2	1:B:414:ILE:HD11	2.29	0.63
1:A:306:VAL:CG2	1:A:414:ILE:HD11	2.31	0.60
1:B:395:LEU:HD22	1:B:456:LEU:HB3	1.89	0.54
1:A:395:LEU:HD22	1:A:456:LEU:HB3	1.89	0.53
1:B:249:HIS:O	4:B:503:3YW:H19	2.09	0.53
1:A:229:GLY:HA3	3:A:502:NTO:O3D	2.11	0.51
1:B:395:LEU:HD21	1:B:457:VAL:HG13	1.95	0.49
1:A:395:LEU:HD21	1:A:457:VAL:HG13	1.94	0.48
1:B:123:ILE:HG12	1:B:132:ILE:HG12	1.98	0.46
1:A:323:HIS:HE1	5:A:610:HOH:O	1.99	0.45
1:A:18:MET:HB2	4:A:503:3YW:C4	2.47	0.45
1:B:229:GLY:HA3	3:B:502:NTO:O3D	2.17	0.44
1:B:323:HIS:HE1	5:B:619:HOH:O	2.00	0.44
1:B:306:VAL:HG22	1:B:414:ILE:HD11	2.00	0.44
1:B:416:LYS:HE3	1:B:471:ARG:HD2	1.99	0.43
1:A:249:HIS:O	4:A:503:3YW:H19	2.18	0.43
1:A:441:ILE:HG21	1:A:449:MET:HB2	2.01	0.43
1:B:113:ASP:O	1:B:117:ARG:HA	2.19	0.43
1:B:77:SER:HB2	1:B:195:VAL:HG11	1.99	0.43
1:A:77:SER:HB2	1:A:195:VAL:HG11	2.01	0.42
1:B:233:LYS:NZ	5:B:604:HOH:O	2.52	0.42
1:B:441:ILE:HG21	1:B:449:MET:HB2	2.01	0.42
1:A:156:THR:HB	1:A:165:TYR:HB3	2.00	0.42
1:A:306:VAL:HG22	1:A:414:ILE:HD11	1.99	0.42
1:A:391:GLU:HA	1:A:417:ARG:HH21	1.85	0.42
1:A:116:THR:HG23	1:A:118:THR:HG22	2.02	0.42
2:A:501:UDP:O1B	4:A:503:3YW:O1	2.39	0.41
1:B:203:ASN:HB3	3:B:502:NTO:O4E	2.20	0.41
1:A:59:LYS:NZ	1:A:382:TYR:OH	2.53	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	491/493 (100%)	473 (96%)	17 (4%)	1 (0%)	51 50
1	B	491/493 (100%)	473 (96%)	17 (4%)	1 (0%)	51 50
All	All	982/986 (100%)	946 (96%)	34 (4%)	2 (0%)	51 50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	298	ASN
1	A	298	ASN

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	450/450 (100%)	446 (99%)	4 (1%)	82 87
1	B	450/450 (100%)	445 (99%)	5 (1%)	78 82
All	All	900/900 (100%)	891 (99%)	9 (1%)	80 84

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	TYR
1	A	137	ASP
1	A	264	LYS
1	A	468	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	47	TYR
1	B	117	ARG
1	B	118	THR
1	B	264	LYS
1	B	363	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	HIS
1	A	203	ASN
1	B	203	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	UDP	A	501	-	21,26,26	0.94	2 (9%)	22,40,40	3.36	2 (9%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NTO	A	502	-	86,95,95	3.83	21 (24%)	109,150,150	1.13	8 (7%)
4	3YW	A	503	-	20,20,20	0.24	0	25,27,27	0.90	2 (8%)
2	UDP	B	501	-	21,26,26	0.99	2 (9%)	22,40,40	3.31	2 (9%)
3	NTO	B	502	-	86,95,95	3.73	22 (25%)	109,150,150	1.26	12 (11%)
4	3YW	B	503	-	20,20,20	0.32	0	25,27,27	0.89	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	501	-	-	0/12/32/32	0/2/2/2
3	NTO	A	502	-	-	0/58/169/169	0/5/5/5
4	3YW	A	503	-	-	0/13/33/33	0/1/1/1
2	UDP	B	501	-	-	0/12/32/32	0/2/2/2
3	NTO	B	502	-	-	0/58/169/169	0/5/5/5
4	3YW	B	503	-	-	0/13/33/33	0/1/1/1

All (47) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	NTO	O4E-S4B	2.04	1.52	1.45
3	A	502	NTO	O1A-S17	2.07	1.53	1.45
3	B	502	NTO	C50-C51	2.09	1.58	1.52
3	B	502	NTO	O52-C5L	2.10	1.47	1.41
3	B	502	NTO	O3E-S3B	2.10	1.53	1.45
3	B	502	NTO	O19-S17	2.12	1.53	1.45
3	B	502	NTO	O4D-S4B	2.13	1.53	1.45
3	A	502	NTO	O3E-S3B	2.16	1.53	1.45
3	B	502	NTO	O36-S35	2.17	1.53	1.45
3	A	502	NTO	O38-S35	2.19	1.53	1.45
3	B	502	NTO	O38-S35	2.19	1.53	1.45
3	A	502	NTO	C26-C20	2.21	1.58	1.52
3	A	502	NTO	O36-S35	2.23	1.53	1.45
3	A	502	NTO	O58-S55	2.25	1.53	1.45
3	A	502	NTO	O4D-S4B	2.26	1.53	1.45
3	B	502	NTO	O1A-S17	2.29	1.53	1.45
3	B	502	NTO	O56-S55	2.30	1.53	1.45
3	A	502	NTO	O56-S55	2.33	1.54	1.45
3	A	502	NTO	O19-S17	2.34	1.54	1.45

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	UDP	C6-N1	2.35	1.39	1.35
3	B	502	NTO	O58-S55	2.37	1.54	1.45
3	B	502	NTO	C20-C21	2.58	1.57	1.52
3	A	502	NTO	C20-C21	2.61	1.57	1.52
2	A	501	UDP	C6-N1	2.63	1.39	1.35
3	B	502	NTO	O42-C4F	2.72	1.48	1.41
2	A	501	UDP	C4-N3	2.86	1.38	1.33
2	B	501	UDP	C4-N3	2.94	1.38	1.33
3	A	502	NTO	O5M-C5L	3.07	1.45	1.40
3	B	502	NTO	O5M-C5L	3.87	1.46	1.40
3	B	502	NTO	S1G-N1F	8.17	1.69	1.59
3	B	502	NTO	O3J-S3H	8.32	1.52	1.42
3	B	502	NTO	O3K-S3H	8.43	1.52	1.42
3	A	502	NTO	O3J-S3H	8.78	1.53	1.42
3	A	502	NTO	O3K-S3H	8.92	1.53	1.42
3	B	502	NTO	O5J-S5H	8.98	1.53	1.42
3	A	502	NTO	O1H-S1G	9.17	1.53	1.42
3	B	502	NTO	O1H-S1G	9.27	1.53	1.42
3	B	502	NTO	O5I-S5H	9.38	1.53	1.42
3	A	502	NTO	O5I-S5H	9.48	1.53	1.42
3	B	502	NTO	O1I-S1G	9.49	1.53	1.42
3	A	502	NTO	O5J-S5H	9.79	1.54	1.42
3	A	502	NTO	O1I-S1G	9.95	1.54	1.42
3	A	502	NTO	S1G-N1F	12.41	1.73	1.59
3	A	502	NTO	S5H-N5G	14.60	1.76	1.59
3	A	502	NTO	S3H-N3G	15.21	1.76	1.59
3	B	502	NTO	S3H-N3G	15.59	1.77	1.59
3	B	502	NTO	S5H-N5G	15.97	1.77	1.59

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	NTO	O3K-S3H-O3J	-4.21	109.00	120.23
3	A	502	NTO	O1I-S1G-O1H	-4.00	109.56	120.23
3	B	502	NTO	O1I-S1G-O1H	-3.70	110.35	120.23
3	B	502	NTO	O3K-S3H-O3J	-3.68	110.40	120.23
2	A	501	UDP	C5-C4-N3	-3.50	114.76	123.12
3	B	502	NTO	O5J-S5H-O5I	-3.49	110.91	120.23
2	B	501	UDP	C5-C4-N3	-3.48	114.81	123.12
3	A	502	NTO	C2C-O2D-C3O	-3.26	110.05	118.00
3	B	502	NTO	C1K-O1L-C2O	-3.25	110.07	118.00
3	B	502	NTO	C2C-O2D-C3O	-3.14	110.35	118.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	NTO	C3L-O3M-C40	-2.71	111.38	118.00
3	A	502	NTO	O5J-S5H-O5I	-2.66	113.13	120.23
3	B	502	NTO	O32-C31-C33	-2.57	101.51	106.64
3	A	502	NTO	C3L-O3M-C40	-2.47	111.97	118.00
3	A	502	NTO	C1K-O1L-C20	-2.21	112.62	118.00
3	B	502	NTO	O54-C53-C51	2.12	111.72	107.67
4	B	503	3YW	O1-C4-C5	2.13	112.94	108.82
3	B	502	NTO	O4A-C49-C4F	2.16	110.57	107.58
3	A	502	NTO	O5M-C5L-C5F	2.25	111.43	108.02
4	A	503	3YW	O4-C3-C2	2.29	115.34	110.65
3	B	502	NTO	C49-O4A-S4B	2.36	123.51	118.97
4	B	503	3YW	O4-C3-C2	2.42	115.60	110.65
3	B	502	NTO	C49-C46-C40	2.49	114.20	109.01
4	A	503	3YW	O1-C4-C5	2.57	113.80	108.82
3	A	502	NTO	O4G-C4F-O42	2.61	117.02	110.70
3	B	502	NTO	C4F-O42-C41	3.01	116.83	112.02
2	B	501	UDP	C4-N3-C2	14.95	126.97	114.13
2	A	501	UDP	C4-N3-C2	15.18	127.17	114.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	UDP	1	0
3	A	502	NTO	1	0
4	A	503	3YW	3	0
3	B	502	NTO	2	0
4	B	503	3YW	2	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	493/493 (100%)	0.25	8 (1%) 72 77	31, 54, 93, 118	0
1	B	493/493 (100%)	0.27	13 (2%) 56 64	32, 54, 95, 132	0
All	All	986/986 (100%)	0.26	21 (2%) 64 71	31, 54, 95, 132	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	295	ASP	4.9
1	B	368	ILE	4.0
1	A	1	MET	3.9
1	B	259	THR	3.8
1	B	114	ILE	3.7
1	B	295	ASP	3.7
1	B	1	MET	3.7
1	A	115	THR	3.6
1	B	260	GLY	3.3
1	A	98	ASN	3.2
1	A	127	LYS	3.1
1	B	98	ASN	2.8
1	B	100	ILE	2.7
1	B	127	LYS	2.5
1	B	113	ASP	2.5
1	A	106	LYS	2.4
1	B	375	ARG	2.3
1	B	256	PHE	2.3
1	B	106	LYS	2.2
1	A	313	LYS	2.2
1	A	444	HIS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	3YW	B	503	20/20	0.80	0.21	5.21	37,52,63,67	20
4	3YW	A	503	20/20	0.88	0.20	4.22	36,51,55,57	20
3	NTO	A	502	91/91	0.95	0.13	0.03	41,51,76,89	91
3	NTO	B	502	91/91	0.95	0.14	0.00	40,52,75,87	91
2	UDP	B	501	25/25	0.97	0.12	-0.48	46,50,56,57	0
2	UDP	A	501	25/25	0.97	0.11	-1.00	45,50,55,57	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.