



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:14 pm GMT

PDB ID : 1X8B
Title : Structure of human Wee1A kinase: kinase domain complexed with inhibitor PD0407824
Authors : Squire, C.J.; Dickson, J.M.; Ivanovic, I.; Baker, E.N.
Deposited on : 2004-08-17
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5868 (1.84-1.80)
Clashscore	112137	6856 (1.84-1.80)
Ramachandran outliers	110173	6780 (1.84-1.80)
Sidechain outliers	110143	6780 (1.84-1.80)
RSRZ outliers	101464	5947 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>5%</div> <div>67%</div> <div>18%</div> <div>• • 10%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2212 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Wee1-like protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	259	Total	C	N	O	S	0	0	0
			2041	1292	364	373	12			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	287	GLY	-	CLONING ARTIFACT	UNP P30291
A	288	ALA	-	CLONING ARTIFACT	UNP P30291
A	289	MET	-	CLONING ARTIFACT	UNP P30291
A	290	GLY	-	CLONING ARTIFACT	UNP P30291

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is 9-HYDROXY-4-PHENYLPYRROLO[3,4-C]CARBAZOLE-1,3(2H,6H)-DIONE (three-letter code: 824) (formula: C₂₀H₁₂N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			25	20	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	144	Total	O	0	0
			144	144		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Wee1-like protein kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	69.78Å 69.78Å 157.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.14 – 1.81 47.07 – 1.81	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.14-1.81) 95.3 (47.07-1.81)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.75 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.218 , 0.237 0.245 , 0.263	Depositor DCC
R_{free} test set	1745 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.637	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2212	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 824, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	1/2084 (0.0%)	1.08	10/2813 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	416	TYR	CD1-CE1	5.04	1.47	1.39

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	527	ARG	NE-CZ-NH1	17.29	128.95	120.30
1	A	527	ARG	NE-CZ-NH2	-9.82	115.39	120.30
1	A	403	LEU	CB-CG-CD2	7.37	123.54	111.00
1	A	321	ASP	CB-CG-OD2	7.07	124.67	118.30
1	A	503	LEU	CB-CG-CD2	6.15	121.45	111.00
1	A	339	ASP	CB-CG-OD2	6.03	123.73	118.30
1	A	369	ASP	CB-CG-OD2	5.63	123.36	118.30
1	A	521	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	426	ASP	CB-CG-OD2	5.42	123.18	118.30
1	A	495	LEU	CB-CG-CD1	5.30	120.00	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2041	0	2022	75	0
2	A	2	0	0	0	0
3	A	25	0	11	0	0
4	A	144	0	0	9	0
All	All	2212	0	2033	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:470:ILE:HD11	1:A:496:PRO:HD3	1.25	1.09
1:A:361:ARG:HD2	4:A:732:HOH:O	1.60	1.00
1:A:291:MET:HE3	1:A:293:SER:H	1.34	0.90
1:A:531:LEU:HD13	1:A:532:PRO:HD2	1.54	0.89
1:A:470:ILE:HD11	1:A:496:PRO:CD	2.03	0.88
1:A:470:ILE:HD12	1:A:470:ILE:O	1.76	0.85
1:A:477:GLU:OE2	1:A:483:LEU:HD22	1.78	0.84
1:A:491:ASN:OD1	1:A:493:THR:HG22	1.77	0.84
1:A:393:ARG:HH11	1:A:393:ARG:HG3	1.43	0.82
1:A:356:HIS:O	4:A:732:HOH:O	2.00	0.79
1:A:357:SER:HA	1:A:361:ARG:HH12	1.48	0.78
1:A:508:VAL:CG2	1:A:531:LEU:HD11	2.13	0.78
1:A:393:ARG:NH2	4:A:740:HOH:O	2.16	0.78
1:A:477:GLU:OE1	1:A:483:LEU:CD1	2.32	0.77
1:A:477:GLU:CD	1:A:483:LEU:HD13	2.06	0.76
1:A:422:LEU:HD13	1:A:469:ARG:HD2	1.66	0.76
1:A:335:ALA:HA	1:A:340:GLU:OE2	1.84	0.75
1:A:566:HIS:ND1	1:A:568:VAL:HG12	2.02	0.74
1:A:477:GLU:OE1	1:A:483:LEU:HD13	1.88	0.74
1:A:477:GLU:OE1	1:A:477:GLU:HA	1.89	0.73
1:A:336:GLY:N	1:A:340:GLU:OE2	2.19	0.72
1:A:424:HIS:HD2	1:A:426:ASP:H	1.37	0.71
1:A:293:SER:O	1:A:297:THR:HG22	1.91	0.70
1:A:508:VAL:HG22	1:A:531:LEU:HD11	1.73	0.69
1:A:477:GLU:CD	1:A:483:LEU:CD1	2.61	0.69
1:A:359:VAL:O	4:A:732:HOH:O	2.09	0.69
1:A:470:ILE:CD1	1:A:496:PRO:HD3	2.15	0.68
1:A:410:VAL:O	1:A:414:LEU:HD13	1.93	0.67

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:477:GLU:OE2	1:A:483:LEU:HD13	1.94	0.67
1:A:477:GLU:OE2	1:A:478:GLY:O	2.13	0.66
1:A:293:SER:HB3	1:A:296:THR:H	1.61	0.65
1:A:361:ARG:NH1	4:A:737:HOH:O	2.30	0.64
1:A:291:MET:HG2	1:A:292:LYS:N	2.12	0.64
1:A:547:LYS:HE2	4:A:739:HOH:O	1.97	0.64
1:A:508:VAL:HG21	1:A:531:LEU:HD11	1.79	0.64
1:A:414:LEU:HD12	1:A:461:ILE:HD11	1.80	0.64
1:A:368:GLU:O	1:A:369:ASP:HB2	1.96	0.63
1:A:485:ASN:HB2	1:A:527:ARG:HG3	1.80	0.63
1:A:477:GLU:OE1	1:A:483:LEU:HD11	1.99	0.62
1:A:393:ARG:HH11	1:A:393:ARG:CG	2.12	0.61
1:A:355:GLN:OE1	1:A:355:GLN:N	2.34	0.60
1:A:357:SER:HA	1:A:361:ARG:NH1	2.17	0.60
1:A:370:ASP:CG	1:A:370:ASP:O	2.40	0.58
1:A:545:LEU:HD22	1:A:568:VAL:HG11	1.86	0.57
1:A:355:GLN:CD	1:A:416:TYR:OH	2.43	0.57
1:A:531:LEU:HD13	1:A:532:PRO:CD	2.33	0.56
1:A:477:GLU:OE1	1:A:477:GLU:CA	2.54	0.54
1:A:477:GLU:OE2	1:A:483:LEU:CD2	2.54	0.53
1:A:414:LEU:HD12	1:A:461:ILE:CD1	2.40	0.51
1:A:475:VAL:HG13	4:A:709:HOH:O	2.10	0.50
1:A:477:GLU:CD	1:A:483:LEU:CD2	2.80	0.50
1:A:291:MET:CG	1:A:292:LYS:N	2.71	0.50
1:A:356:HIS:HB2	1:A:416:TYR:CE2	2.46	0.50
1:A:477:GLU:CD	1:A:483:LEU:HD22	2.31	0.49
1:A:414:LEU:CD1	1:A:461:ILE:HD11	2.42	0.48
1:A:370:ASP:OD1	1:A:370:ASP:O	2.32	0.47
1:A:393:ARG:NH1	1:A:393:ARG:CG	2.74	0.47
1:A:404:LYS:HE2	1:A:568:VAL:HG22	1.97	0.46
1:A:545:LEU:HD22	1:A:568:VAL:CG1	2.46	0.46
1:A:332:LYS:HE2	4:A:734:HOH:O	2.16	0.45
1:A:470:ILE:HD11	1:A:496:PRO:CG	2.45	0.45
1:A:493:THR:HG23	1:A:494:HIS:CD2	2.51	0.45
1:A:466:HIS:HD2	4:A:738:HOH:O	2.01	0.44
1:A:414:LEU:N	1:A:414:LEU:HD12	2.32	0.44
1:A:540:GLN:O	1:A:544:GLU:HG3	2.18	0.43
1:A:291:MET:O	1:A:292:LYS:HB2	2.17	0.43
1:A:423:VAL:CG1	1:A:470:ILE:HD13	2.49	0.43
1:A:297:THR:O	1:A:319:ARG:HD2	2.18	0.43
1:A:424:HIS:HE1	1:A:462:GLY:O	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:424:HIS:CD2	1:A:426:ASP:H	2.27	0.41
1:A:355:GLN:HG2	1:A:416:TYR:OH	2.20	0.41
1:A:291:MET:CG	1:A:292:LYS:H	2.34	0.41
1:A:313:VAL:HG22	1:A:328:LYS:HG3	2.01	0.41
1:A:291:MET:HE3	1:A:293:SER:N	2.18	0.40
1:A:354:GLY:N	1:A:355:GLN:OE1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	255/289 (88%)	247 (97%)	7 (3%)	1 (0%)	38 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	369	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	219/246 (89%)	205 (94%)	14 (6%)	20 7

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	293	SER
1	A	345	ARG
1	A	370	ASP
1	A	374	ILE
1	A	393	ARG
1	A	399	LYS
1	A	422	LEU
1	A	470	ILE
1	A	477	GLU
1	A	495	LEU
1	A	503	LEU
1	A	505	LEU
1	A	527	ARG
1	A	531	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	358	HIS
1	A	424	HIS
1	A	466	HIS
1	A	494	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	824	A	901	-	26,29,29	1.43	4 (15%)	32,44,44	2.52	12 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	824	A	901	-	-	0/4/4/4	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	824	O3-C19	-2.04	1.32	1.37
3	A	901	824	C20-C19	2.49	1.43	1.38
3	A	901	824	C6-C3	2.53	1.47	1.43
3	A	901	824	C17-C19	2.58	1.42	1.37

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	824	C19-C17-C15	-7.93	115.63	120.64
3	A	901	824	C4-C13-C14	-4.28	116.57	119.85
3	A	901	824	C13-C15-C16	-3.51	102.30	106.09
3	A	901	824	C11-C12-C7	-2.81	117.00	120.57
3	A	901	824	C20-C18-C16	-2.65	117.59	120.84
3	A	901	824	C10-C9-C8	-2.43	116.87	120.21
3	A	901	824	C1-C2-C3	-2.38	116.47	119.22
3	A	901	824	C13-C14-N2	-2.35	104.41	108.89
3	A	901	824	C14-N2-C16	3.04	113.34	107.09
3	A	901	824	C18-C20-C19	3.54	124.20	120.14
3	A	901	824	C15-C13-C14	4.00	110.43	106.10
3	A	901	824	C17-C15-C16	4.47	124.37	118.26

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/289 (89%)	0.37	14 (5%) 26 22	18, 25, 41, 55	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	310	PHE	8.5
1	A	291	MET	6.2
1	A	309	GLU	3.4
1	A	369	ASP	3.1
1	A	311	GLY	3.1
1	A	292	LYS	3.0
1	A	336	GLY	3.0
1	A	334	LEU	2.8
1	A	370	ASP	2.8
1	A	338	VAL	2.7
1	A	307	SER	2.7
1	A	335	ALA	2.5
1	A	339	ASP	2.3
1	A	565	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	824	A	901	25/25	0.95	0.11	-0.33	20,23,25,29	0
2	MG	A	602	1/1	0.91	0.09	-1.11	37,37,37,37	0
2	MG	A	601	1/1	0.96	0.08	-	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.