



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 04:39 am GMT

PDB ID : 1XIC
Title : MODES OF BINDING SUBSTRATES AND THEIR ANALOGUES TO THE
ENZYME D-XYLOSE ISOMERASE
Authors : Carrell, H.L.; Glusker, J.P.
Deposited on : 1994-03-07
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	388	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XLS	A	389	-	X	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3415 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called D-XYLOSE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	385	3031	1906	545	572	8	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	GLN	ARG	CONFLICT	UNP P24300

- Molecule 2 is SUGAR (D-XYLOSE (LINEAR FORM)) (three-letter code: XLS) (formula: $C_5H_{10}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	5	5	0	0

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	Mn 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	372	Total 372	O 372	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: D-XYLOSE ISOMERASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.90Å 99.70Å 102.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	9.00 – 1.60	Depositor
% Data completeness (in resolution range)	(Not available) (9.00-1.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.152 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3415	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, XLS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	10/3103 (0.3%)	2.23	105/4201 (2.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	217	GLU	CD-OE1	-9.45	1.15	1.25
1	A	217	GLU	CD-OE2	-7.92	1.17	1.25
1	A	292	ARG	CZ-NH1	5.98	1.40	1.33
1	A	253	LYS	C-O	5.87	1.34	1.23
1	A	112	ARG	CZ-NH2	5.56	1.40	1.33
1	A	137	TRP	CD2-CE2	5.45	1.47	1.41
1	A	340	ARG	CD-NE	-5.20	1.37	1.46
1	A	162	PHE	CE1-CZ	5.07	1.47	1.37
1	A	82	THR	CA-CB	5.04	1.66	1.53
1	A	139	GLY	N-CA	5.01	1.53	1.46

All (105) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	ARG	CD-NE-CZ	40.09	179.73	123.60
1	A	68	ARG	NE-CZ-NH2	24.08	132.34	120.30
1	A	76	ARG	NE-CZ-NH2	-19.28	110.66	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	340	ARG	NE-CZ-NH2	-17.71	111.45	120.30
1	A	340	ARG	NE-CZ-NH1	17.65	129.12	120.30
1	A	74	ARG	NE-CZ-NH2	-16.46	112.07	120.30
1	A	113	ARG	NE-CZ-NH1	15.26	127.93	120.30
1	A	331	ARG	NE-CZ-NH2	-15.07	112.77	120.30
1	A	266	ARG	NE-CZ-NH2	-14.87	112.87	120.30
1	A	387	ARG	CD-NE-CZ	12.88	141.63	123.60
1	A	316	ARG	NE-CZ-NH2	-12.80	113.90	120.30
1	A	117	ARG	NE-CZ-NH1	12.42	126.51	120.30
1	A	113	ARG	NE-CZ-NH2	-12.23	114.18	120.30
1	A	140	ARG	NE-CZ-NH2	-12.06	114.27	120.30
1	A	212	TYR	CB-CG-CD2	-12.04	113.77	121.00
1	A	363	ASP	CB-CG-OD2	-11.94	107.55	118.30
1	A	74	ARG	NE-CZ-NH1	11.76	126.18	120.30
1	A	284	ARG	NE-CZ-NH2	-11.76	114.42	120.30
1	A	10	ARG	NE-CZ-NH1	11.56	126.08	120.30
1	A	188	ARG	NE-CZ-NH1	10.80	125.70	120.30
1	A	266	ARG	CD-NE-CZ	10.68	138.55	123.60
1	A	124	ASP	CB-CG-OD1	10.52	127.77	118.30
1	A	292	ARG	NE-CZ-NH1	-10.48	115.06	120.30
1	A	368	ARG	NE-CZ-NH1	10.28	125.44	120.30
1	A	68	ARG	NH1-CZ-NH2	-10.09	108.31	119.40
1	A	175	ASP	CB-CG-OD1	-10.05	109.26	118.30
1	A	152	ARG	NE-CZ-NH2	-9.84	115.38	120.30
1	A	124	ASP	CB-CG-OD2	-9.64	109.62	118.30
1	A	208	ARG	NE-CZ-NH2	-9.51	115.54	120.30
1	A	316	ARG	NE-CZ-NH1	9.30	124.95	120.30
1	A	287	ASP	CB-CG-OD1	9.26	126.63	118.30
1	A	121	ARG	NE-CZ-NH2	-9.17	115.72	120.30
1	A	292	ARG	NE-CZ-NH2	9.00	124.80	120.30
1	A	76	ARG	NH1-CZ-NH2	8.96	129.25	119.40
1	A	277	SER	CB-CA-C	8.80	126.82	110.10
1	A	323	ASP	CB-CG-OD1	8.76	126.18	118.30
1	A	295	ASP	CB-CG-OD1	8.72	126.15	118.30
1	A	334	ARG	NE-CZ-NH2	-8.55	116.03	120.30
1	A	104	PHE	CB-CG-CD2	-8.51	114.85	120.80
1	A	28	ASP	CB-CG-OD1	8.46	125.92	118.30
1	A	42	ARG	CD-NE-CZ	8.38	135.33	123.60
1	A	368	ARG	NE-CZ-NH2	-8.35	116.12	120.30
1	A	254	TYR	CB-CG-CD2	-8.32	116.01	121.00
1	A	152	ARG	NE-CZ-NH1	8.27	124.44	120.30
1	A	212	TYR	CB-CG-CD1	8.16	125.89	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	23	ARG	NE-CZ-NH1	8.14	124.37	120.30
1	A	266	ARG	NE-CZ-NH1	8.10	124.35	120.30
1	A	28	ASP	CB-CG-OD2	-7.93	111.16	118.30
1	A	363	ASP	CB-CG-OD1	7.93	125.43	118.30
1	A	331	ARG	NH1-CZ-NH2	7.90	128.09	119.40
1	A	169	VAL	CA-CB-CG1	7.87	122.70	110.90
1	A	387	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	A	284	ARG	NE-CZ-NH1	7.82	124.21	120.30
1	A	24	ASP	CB-CA-C	7.81	126.02	110.40
1	A	175	ASP	OD1-CG-OD2	7.77	138.06	123.30
1	A	273	ASP	CB-CG-OD2	7.71	125.24	118.30
1	A	140	ARG	NH1-CZ-NH2	7.65	127.82	119.40
1	A	188	ARG	NE-CZ-NH2	-7.64	116.48	120.30
1	A	196	VAL	CA-CB-CG2	-7.51	99.63	110.90
1	A	321	ARG	NE-CZ-NH1	7.38	123.99	120.30
1	A	297	ASP	CB-CG-OD1	7.31	124.88	118.30
1	A	387	ARG	NE-CZ-NH2	-7.28	116.66	120.30
1	A	259	ARG	NE-CZ-NH1	7.28	123.94	120.30
1	A	10	ARG	NE-CZ-NH2	-7.22	116.69	120.30
1	A	175	ASP	CA-CB-CG	-7.08	97.82	113.40
1	A	245	ASP	CB-CG-OD1	7.01	124.61	118.30
1	A	217	GLU	OE1-CD-OE2	6.91	131.59	123.30
1	A	174	TYR	CB-CG-CD1	6.83	125.10	121.00
1	A	12	THR	O-C-N	6.74	133.48	122.70
1	A	76	ARG	CG-CD-NE	-6.72	97.70	111.80
1	A	121	ARG	NE-CZ-NH1	6.67	123.63	120.30
1	A	175	ASP	CB-CG-OD2	-6.57	112.38	118.30
1	A	150	ASP	CB-CG-OD1	6.42	124.08	118.30
1	A	169	VAL	CG1-CB-CG2	6.40	121.15	110.90
1	A	264	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	245	ASP	CB-CG-OD2	-6.15	112.77	118.30
1	A	292	ARG	CD-NE-CZ	6.13	132.19	123.60
1	A	4	GLN	CB-CG-CD	6.07	127.38	111.60
1	A	361	ASP	CB-CG-OD2	-6.04	112.87	118.30
1	A	316	ARG	CD-NE-CZ	-6.03	115.16	123.60
1	A	68	ARG	CA-CB-CG	6.01	126.62	113.40
1	A	31	ARG	NE-CZ-NH2	5.99	123.30	120.30
1	A	74	ARG	CD-NE-CZ	5.87	131.82	123.60
1	A	207	GLU	CG-CD-OE2	5.82	129.93	118.30
1	A	205	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	A	65	ASP	CB-CA-C	5.74	121.87	110.40
1	A	217	GLU	CG-CD-OE1	-5.64	107.02	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	ASP	CB-CG-OD1	5.58	123.32	118.30
1	A	212	TYR	CZ-CE2-CD2	-5.48	114.87	119.80
1	A	132	GLU	CB-CA-C	-5.39	99.62	110.40
1	A	132	GLU	N-CA-CB	-5.39	100.90	110.60
1	A	208	ARG	NE-CZ-NH1	5.37	122.98	120.30
1	A	35	ASP	N-CA-CB	-5.36	100.95	110.60
1	A	257	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	265	LEU	CB-CG-CD2	-5.29	102.00	111.00
1	A	207	GLU	CG-CD-OE1	-5.25	107.81	118.30
1	A	264	ASP	CB-CG-OD2	-5.23	113.59	118.30
1	A	160	GLU	OE1-CD-OE2	5.23	129.57	123.30
1	A	178	PHE	CB-CG-CD2	-5.22	117.15	120.80
1	A	373	GLU	CG-CD-OE1	5.18	128.67	118.30
1	A	35	ASP	CB-CG-OD2	-5.17	113.65	118.30
1	A	65	ASP	CB-CG-OD1	5.16	122.95	118.30
1	A	371	ALA	O-C-N	-5.13	114.48	122.70
1	A	8	GLU	CA-CB-CG	5.08	124.57	113.40
1	A	387	ARG	CB-CA-C	5.05	120.51	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	42	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3031	0	2900	5	2
2	A	10	0	6	3	0
3	A	2	0	0	0	0
4	A	372	0	0	3	3
All	All	3415	0	2906	7	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:389:XLS:H52	4:A:565:HOH:O	2.03	0.57
1:A:354:ARG:HB3	1:A:359:GLU:HG3	1.89	0.55
1:A:289:LYS:HE2	4:A:727:HOH:O	2.09	0.52
1:A:186:GLU:OE1	1:A:255:ASP:HB3	2.13	0.48
2:A:389:XLS:H4	2:A:389:XLS:H1	1.77	0.43
1:A:54:HIS:NE2	2:A:389:XLS:C1	2.82	0.42
1:A:180:ILE:HG13	4:A:739:HOH:O	2.20	0.42

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:253:LYS:CE	4:A:737:HOH:O[4_566]	1.55	0.65
1:A:253:LYS:NZ	4:A:737:HOH:O[4_566]	1.89	0.31
4:A:603:HOH:O	4:A:667:HOH:O[3_656]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	383/388 (99%)	370 (97%)	12 (3%)	1 (0%)	44 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	302/304 (99%)	291 (96%)	11 (4%)	40 14

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	8	GLU
1	A	42	ARG
1	A	66	SER
1	A	76	ARG
1	A	91	THR
1	A	132	GLU
1	A	169	VAL
1	A	175	ASP
1	A	185	ASN
1	A	340	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	GLN
1	A	234	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XLS	A	389	3	9,9,9	4.94	5 (55%)	11,11,11	7.40	9 (81%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XLS	A	389	3	-	0/10/12/12	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	389	XLS	C2-C1	-11.67	1.30	1.50
2	A	389	XLS	O5-C5	-5.84	1.17	1.42
2	A	389	XLS	O2-C2	-5.67	1.32	1.43
2	A	389	XLS	C3-C2	-2.92	1.48	1.53
2	A	389	XLS	C5-C4	-2.04	1.46	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	389	XLS	C5-C4-C3	-12.16	85.28	112.41
2	A	389	XLS	O1-C1-C2	-8.90	100.88	125.13
2	A	389	XLS	O3-C3-C2	-6.57	97.13	109.17
2	A	389	XLS	C3-C2-C1	-5.04	95.26	111.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	389	XLS	O3-C3-C4	2.53	115.08	108.82
2	A	389	XLS	O2-C2-C3	6.24	124.29	109.46
2	A	389	XLS	O5-C5-C4	6.70	125.88	111.11
2	A	389	XLS	O4-C4-C5	10.27	132.95	109.21
2	A	389	XLS	O2-C2-C1	10.43	135.06	110.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	389	XLS	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.