



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2017 – 12:54 am GMT

PDB ID : 2XJ7
Title : BTGH84 IN COMPLEX WITH 6-ACETAMIDO-6-DEOXY-CASTANOSPERMINE
Authors : He, Y.; Davies, G.J.
Deposited on : 2010-07-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	6609 (2.00-2.00)
Clashscore	112137	7775 (2.00-2.00)
Ramachandran outliers	110173	7679 (2.00-2.00)
Sidechain outliers	110143	7678 (2.00-2.00)
RSRZ outliers	101464	6696 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	716	<div> <div>4%</div> <div> <div></div> <div>77%</div> <div>11%</div> <div>11%</div> </div> </div>
1	B	716	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>12%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 10749 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called O-GLCNACASE BT_4395.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	639	Total	C	N	O	S	0	2	0
			5184	3323	870	972	19			
1	B	633	Total	C	N	O	S	0	1	0
			5139	3296	865	960	18			

- Molecule 2 is 6-ACETAMIDO-6-DEOXY-CASTANOSPERMINE (three-letter code: GC2) (formula: $C_{10}H_{18}N_2O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			16	10	2	4		
2	B	1	Total	C	N	O	0	0
			16	10	2	4		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Ca 1	0	0
3	A	1	Total 1	Ca 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	190	Total 190	O 190	0	0
4	B	202	Total 202	O 202	0	0

GLN	GLN	VAL	TYR	LEU	ARG	GLN	F709	V710	L711	T712	I713	E714	K715	LYS
							●	●	●		●			

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	51.49Å 93.98Å 98.81Å 104.13° 93.88° 103.13°	Depositor
Resolution (Å)	47.50 – 2.00 47.50 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.5 (47.50-2.00) 88.8 (47.50-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.203 , 0.239 0.209 , 0.246	Depositor DCC
R_{free} test set	5649 reflections (5.19%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10749	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GC2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/5318 (0.0%)	0.73	0/7205
1	B	0.85	1/5268 (0.0%)	0.73	0/7132
All	All	0.85	2/10586 (0.0%)	0.73	0/14337

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	32	GLU	CD-OE2	7.91	1.34	1.25
1	A	145	ALA	CA-CB	5.24	1.63	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5184	0	5089	56	0
1	B	5139	0	5057	46	0
2	A	16	0	18	0	0
2	B	16	0	18	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	190	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	202	0	0	4	0
All	All	10749	0	10182	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:LEU:HB2	1:A:104:LEU:HD21	1.30	1.07
1:A:41:LEU:HB2	1:A:104:LEU:CD2	1.99	0.91
1:B:41:LEU:HB2	1:B:104:LEU:HD21	1.59	0.85
1:B:451:LEU:HG	1:B:455:LYS:HE2	1.67	0.75
1:A:19:ILE:CD1	1:A:118:GLU:HG2	2.19	0.71
1:A:19:ILE:HD11	1:A:118:GLU:HG2	1.72	0.70
1:B:462:LYS:HG2	1:B:466:GLU:OE2	1.93	0.68
1:B:593:HIS:HD2	1:B:636:ASP:H	1.43	0.67
1:B:454:PHE:HZ	1:B:516:VAL:HG13	1.64	0.62
1:A:643:ASN:HB2	1:A:682:ALA:O	1.99	0.61
1:B:490:PRO:O	1:B:494:GLU:HG3	1.99	0.61
1:A:444:GLN:NE2	1:A:448:GLU:OE2	2.31	0.60
1:A:81:TYR:CE2	1:A:123:ASP:HB3	2.37	0.59
1:A:15:GLN:HB2	1:A:118:GLU:HB2	1.85	0.58
1:A:34:ASN:O	1:A:38:VAL:HG23	2.03	0.58
1:A:37:ALA:O	1:A:104:LEU:HD22	2.04	0.57
1:B:462:LYS:CG	1:B:466:GLU:OE2	2.52	0.57
1:B:616:ILE:HD11	1:B:633:ILE:HD11	1.87	0.57
1:A:19:ILE:HD11	1:A:118:GLU:CG	2.34	0.57
1:A:308:MET:HA	1:A:335:TYR:O	2.05	0.56
1:A:593:HIS:HD2	1:A:636:ASP:H	1.55	0.54
1:A:55:MET:HE2	1:A:90:ILE:HG13	1.90	0.54
1:B:238:ALA:HA	1:B:276:VAL:O	2.07	0.54
1:B:543:GLN:NE2	1:B:609:VAL:HG12	2.22	0.54
1:B:454:PHE:CZ	1:B:516:VAL:HG13	2.43	0.53
1:A:385:TYR:CD2	1:A:406:ALA:HB2	2.43	0.53
1:A:423[B]:MET:SD	1:A:442[B]:ASP:OD2	2.66	0.53
1:B:451:LEU:HG	1:B:455:LYS:CE	2.38	0.53
1:A:55:MET:CE	1:A:90:ILE:HG13	2.40	0.52
1:B:308:MET:HA	1:B:335:TYR:O	2.10	0.52
1:A:489:LYS:HB2	1:A:490:PRO:HD3	1.92	0.52
1:A:451:LEU:HG	1:A:564:LEU:HD12	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:GLY:N	1:A:165:PRO:HD3	2.25	0.52
1:B:536:GLN:HG3	1:B:590:TYR:CD1	2.44	0.52
1:B:715:LYS:NZ	4:B:2202:HOH:O	2.42	0.52
1:A:19:ILE:CD1	1:A:118:GLU:CG	2.88	0.52
1:A:562:LYS:HB3	1:A:563:PRO:HD3	1.93	0.51
1:B:37:ALA:O	1:B:104:LEU:HD22	2.10	0.50
1:B:133:VAL:HG13	1:B:133:VAL:O	2.11	0.50
1:B:536:GLN:CG	1:B:590:TYR:CD1	2.94	0.50
1:A:568:THR:O	1:A:572:VAL:HG22	2.13	0.48
1:B:254:GLN:O	1:B:258:LEU:HD23	2.13	0.48
1:B:473:GLU:O	1:B:477:GLU:HG3	2.13	0.48
1:A:26:GLN:HG2	1:A:56:LEU:HD12	1.96	0.47
1:B:97:GLU:HG3	4:B:2020:HOH:O	2.13	0.47
1:A:26:GLN:HE21	1:A:56:LEU:HD13	1.79	0.47
1:A:474:ARG:HA	1:A:474:ARG:HD2	1.70	0.47
1:B:444:GLN:NE2	1:B:448:GLU:OE2	2.48	0.47
1:B:170:TYR:HB2	1:B:180:TYR:CE1	2.50	0.47
1:B:239:VAL:HG12	1:B:241:PHE:CE2	2.50	0.47
1:A:616:ILE:HG13	1:A:709:PHE:HB3	1.97	0.46
1:A:644:ILE:O	1:A:681:SER:HA	2.16	0.46
1:A:324:SER:O	1:A:328:GLU:HG2	2.16	0.46
1:B:315:ILE:HD11	2:B:1000:GC2:C7	2.46	0.45
1:A:301:LEU:HD12	1:A:307:ILE:HD11	1.98	0.45
1:A:262:ASP:O	1:A:267:GLN:HG2	2.15	0.45
1:B:423:MET:HE3	1:B:423:MET:HB3	1.73	0.45
1:B:314:VAL:HG23	4:B:2112:HOH:O	2.16	0.44
1:A:484:MET:HB3	1:B:536:GLN:NE2	2.32	0.44
1:A:557:ALA:HB1	1:A:561:ILE:HB	1.99	0.44
1:A:153:TYR:O	1:A:154:GLY:C	2.55	0.44
1:A:26:GLN:NE2	1:A:56:LEU:HD13	2.33	0.44
1:B:129:TYR:O	1:B:368:GLY:HA2	2.18	0.44
1:B:557:ALA:HB1	1:B:561:ILE:HB	1.99	0.44
1:A:26:GLN:HE21	1:A:56:LEU:CD1	2.30	0.43
1:B:261:ILE:O	1:B:265:PHE:HB3	2.18	0.43
1:B:250:ASN:O	1:B:254:GLN:HG3	2.19	0.43
1:B:56:LEU:HD23	1:B:89:GLU:OE1	2.18	0.43
1:A:125:PRO:HB3	1:A:392:TRP:CE3	2.53	0.43
1:A:461:ASP:O	1:A:463:ALA:N	2.51	0.43
1:B:385:TYR:CD2	1:B:406:ALA:HB2	2.54	0.43
1:A:271:ASP:OD1	1:A:271:ASP:N	2.51	0.43
1:A:323:ILE:HG13	1:A:323:ILE:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:609:VAL:O	1:B:610:LYS:HD3	2.19	0.43
1:A:646:ILE:O	1:A:680:LEU:HB2	2.19	0.43
1:B:341:PRO:O	1:B:342:VAL:C	2.57	0.43
1:B:454:PHE:O	1:B:455:LYS:C	2.57	0.42
1:A:187:GLN:O	1:A:190:GLU:HB3	2.19	0.42
1:A:64:ASP:O	1:A:65:LYS:C	2.58	0.42
1:B:81:TYR:CZ	1:B:123:ASP:HB3	2.53	0.42
1:B:288:ASN:C	1:B:288:ASN:OD1	2.57	0.42
1:A:133:VAL:HG13	1:A:133:VAL:O	2.20	0.42
1:A:532:LYS:HD2	1:A:532:LYS:HA	1.95	0.42
1:A:516:VAL:HG22	1:A:572:VAL:HG11	2.01	0.42
1:A:453:ALA:HB1	1:A:459:ASN:O	2.20	0.41
1:B:170:TYR:CZ	1:B:181:PRO:HD3	2.55	0.41
1:A:26:GLN:HB2	1:A:51:SER:O	2.20	0.41
1:B:143:HIS:NE2	1:B:190:GLU:OE1	2.40	0.41
1:B:451:LEU:HA	1:B:451:LEU:HD12	1.85	0.41
1:A:81:TYR:CZ	1:A:123:ASP:HB3	2.55	0.41
1:B:174:PRO:HD2	4:B:2062:HOH:O	2.21	0.41
1:A:606:PRO:HG2	1:A:617:SER:HB2	2.03	0.41
1:A:451:LEU:HG	1:A:564:LEU:CD1	2.50	0.41
1:A:126:SER:HB2	1:A:394:PRO:HD2	2.03	0.40
1:A:288:ASN:HA	1:A:289:PRO:HD2	1.94	0.40
1:B:282:TYR:OH	1:B:286:TRP:CZ3	2.72	0.40
1:B:22:PRO:HB2	1:B:25:TYR:HB3	2.02	0.40
1:B:21:LEU:HD12	1:B:22:PRO:HD2	2.04	0.40
1:A:14:VAL:HG22	1:A:15:GLN:O	2.22	0.40
1:A:131:GLY:HA3	1:A:160:THR:O	2.22	0.40
1:A:166:LYS:NZ	1:A:242:ASP:OD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	627/716 (88%)	602 (96%)	25 (4%)	0	100	100
1	B	618/716 (86%)	593 (96%)	25 (4%)	0	100	100
All	All	1245/1432 (87%)	1195 (96%)	50 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/630 (89%)	556 (99%)	6 (1%)	78	82
1	B	557/630 (88%)	553 (99%)	4 (1%)	87	90
All	All	1119/1260 (89%)	1109 (99%)	10 (1%)	82	87

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	39	LYS
1	A	371	THR
1	A	461	ASP
1	A	613	ARG
1	A	616	ILE
1	B	104	LEU
1	B	423	MET
1	B	454	PHE
1	B	615	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	349	HIS
1	A	433	HIS
1	A	547	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	593	HIS
1	B	10	GLN
1	B	26	GLN
1	B	536	GLN
1	B	543	GLN
1	B	547	GLN
1	B	578	GLN
1	B	593	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GC2	A	1000	-	17,17,17	0.96	1 (5%)	18,25,25	1.42	2 (11%)
2	GC2	B	1000	-	17,17,17	1.04	1 (5%)	18,25,25	2.02	7 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GC2	A	1000	-	-	0/4/33/33	1/2/2/2
2	GC2	B	1000	-	-	0/4/33/33	1/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	GC2	C1-C2	2.07	1.55	1.52
2	B	1000	GC2	C1-C2	2.29	1.56	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1000	GC2	O3-C6-C7	-2.65	105.52	111.60
2	B	1000	GC2	O4-C3-C2	-2.60	103.81	109.39
2	B	1000	GC2	O1-C8-C10	-2.02	118.38	122.06
2	B	1000	GC2	O4-C3-C4	2.02	114.74	110.36
2	B	1000	GC2	C1-C2-N2	2.42	113.32	109.71
2	B	1000	GC2	C10-C8-N2	2.50	120.62	116.11
2	A	1000	GC2	C2-N2-C8	3.40	127.91	122.94
2	A	1000	GC2	C1-C2-N2	4.47	116.37	109.71
2	B	1000	GC2	C2-N2-C8	5.47	130.92	122.94

There are no chirality outliers.

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1000	GC2	C1-C2-C3-C4-C5-N1
2	B	1000	GC2	C1-C2-C3-C4-C5-N1

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1000	GC2	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	639/716 (89%)	-0.05	32 (5%) 30 30	25, 38, 75, 94	0
1	B	633/716 (88%)	0.03	45 (7%) 17 17	25, 40, 81, 101	0
All	All	1272/1432 (88%)	-0.01	77 (6%) 22 22	25, 39, 79, 101	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	709	PHE	7.1
1	B	691	PHE	5.6
1	B	689	VAL	5.5
1	B	638	ILE	5.5
1	B	692	VAL	5.3
1	A	633	ILE	5.2
1	A	692	VAL	5.0
1	B	684	LEU	5.0
1	B	710	VAL	5.0
1	B	24	VAL	4.9
1	A	646	ILE	4.7
1	B	614	VAL	4.4
1	A	680	LEU	4.3
1	B	53	LYS	4.3
1	B	47	GLY	4.1
1	A	710	VAL	4.0
1	B	21	LEU	4.0
1	A	709	PHE	3.9
1	B	52	LYS	3.7
1	B	711	LEU	3.7
1	B	595	MET	3.6
1	B	643	ASN	3.6
1	A	644	ILE	3.5
1	B	23	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	455	LYS	3.4
1	A	691	PHE	3.4
1	B	51	SER	3.4
1	B	48	LYS	3.4
1	B	644	ILE	3.4
1	B	637	ALA	3.3
1	A	687	ALA	3.2
1	B	686	LYS	3.2
1	A	47	GLY	3.1
1	B	690	LYS	3.1
1	B	594	LYS	3.1
1	A	49	GLN	3.1
1	A	682	ALA	3.0
1	B	617	SER	3.0
1	B	616	ILE	3.0
1	B	49	GLN	2.9
1	A	688	PRO	2.9
1	A	52	LYS	2.8
1	B	688	PRO	2.8
1	A	459	ASN	2.8
1	B	633	ILE	2.8
1	A	631	VAL	2.7
1	A	452	LYS	2.7
1	B	16	ASN	2.7
1	B	641	GLY	2.7
1	B	640	PRO	2.6
1	B	45	LEU	2.6
1	B	54	GLY	2.6
1	B	683	GLY	2.6
1	A	684	LEU	2.6
1	A	581	ASN	2.5
1	A	681	SER	2.5
1	B	451	LEU	2.5
1	B	687	ALA	2.5
1	B	25	TYR	2.4
1	B	713	ILE	2.4
1	A	616	ILE	2.4
1	B	455	LYS	2.3
1	B	46	SER	2.3
1	A	712	THR	2.2
1	A	24	VAL	2.2
1	A	715	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	594	LYS	2.2
1	A	595	MET	2.2
1	A	21	LEU	2.2
1	B	22	PRO	2.2
1	B	19	ILE	2.1
1	B	645	GLN	2.1
1	A	25	TYR	2.1
1	B	50	SER	2.1
1	A	16	ASN	2.1
1	A	645	GLN	2.1
1	A	632	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GC2	A	1000	16/16	0.94	0.10	0.63	26,28,31,33	0
2	GC2	B	1000	16/16	0.94	0.09	-0.29	26,29,32,32	0
3	CA	A	1716	1/1	0.85	0.08	-0.73	54,54,54,54	0
3	CA	B	1716	1/1	0.95	0.16	-	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.