



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:45 am GMT

PDB ID : 2XJM  
Title : CRYSTAL STRUCTURE OF STREPTOCOCCUS SUIS DPR WITH COBALT  
Authors : Haikarainen, T.; Thanassoulas, A.; Stavros, P.; Nounesis, G.; Haataja, S.; Papageorgiou, A.C.  
Deposited on : 2010-07-06  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	165	<div> <div>2%</div> <div> <div></div> <div>81%</div> <div>10%</div> <div>9%</div> </div> </div>
1	B	165	<div> <div>2%</div> <div> <div></div> <div>79%</div> <div>13%</div> <div>8%</div> </div> </div>
1	C	165	<div> <div>0%</div> <div> <div></div> <div>80%</div> <div>11%</div> <div>9%</div> </div> </div>
1	D	165	<div> <div>0%</div> <div> <div></div> <div>80%</div> <div>12%</div> <div>8%</div> </div> </div>
1	E	165	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>15%</div> <div>9%</div> </div> </div>
1	F	165	<div> <div>0%</div> <div> <div></div> <div>83%</div> <div>8%</div> <div>8%</div> </div> </div>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	165	
1	H	165	
1	I	165	
1	J	165	
1	K	165	
1	L	165	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CL	A	1174	-	-	-	X
5	EPE	E	1173	-	-	-	X
5	EPE	K	1173	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 15464 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA PROTECTION DURING STARVATION PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	150	Total	C	N	O	S	0	2	0
			1217	774	200	236	7			
1	B	152	Total	C	N	O	S	0	1	0
			1213	773	200	234	6			
1	C	150	Total	C	N	O	S	0	0	0
			1193	761	198	228	6			
1	D	152	Total	C	N	O	S	0	2	0
			1232	783	205	238	6			
1	E	150	Total	C	N	O	S	0	1	0
			1202	765	198	233	6			
1	F	151	Total	C	N	O	S	0	2	0
			1215	776	202	231	6			
1	G	150	Total	C	N	O	S	0	0	0
			1197	762	197	232	6			
1	H	151	Total	C	N	O	S	0	1	0
			1213	773	200	234	6			
1	I	151	Total	C	N	O	S	0	2	0
			1222	777	203	236	6			
1	J	151	Total	C	N	O	S	0	1	0
			1214	773	199	236	6			
1	K	152	Total	C	N	O	S	0	4	0
			1245	790	206	243	6			
1	L	151	Total	C	N	O	S	0	2	0
			1216	775	201	234	6			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Co	0	0
			1	1		
2	J	1	Total	Co	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	1	Total 1	Co 1	0	0
2	K	1	Total 1	Co 1	0	0
2	E	1	Total 1	Co 1	0	0
2	H	1	Total 1	Co 1	0	0
2	B	1	Total 1	Co 1	0	0
2	I	1	Total 1	Co 1	0	0
2	C	1	Total 1	Co 1	0	0
2	A	1	Total 1	Co 1	0	0
2	L	1	Total 1	Co 1	0	0
2	F	1	Total 1	Co 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	I	1	Total 1	Cl 1	0	0
3	A	1	Total 1	Cl 1	0	0
3	J	1	Total 1	Cl 1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	J	1	Total 1	Ca 1	0	0
4	E	1	Total 1	Ca 1	0	0
4	H	1	Total 1	Ca 1	0	0
4	B	1	Total 1	Ca 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		
4	F	2	Total	Ca	0	0
			2	2		

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	E	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	I	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	J	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
5	K	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	63	Total	O	0	0
			63	63		
6	B	93	Total	O	0	0
			93	93		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	49	Total 49	O 49	0	0
6	D	89	Total 89	O 89	0	0
6	E	44	Total 44	O 44	0	0
6	F	72	Total 72	O 72	0	0
6	G	52	Total 52	O 52	0	0
6	H	54	Total 54	O 54	0	0
6	I	67	Total 67	O 67	0	0
6	J	73	Total 73	O 73	0	0
6	K	76	Total 76	O 76	0	0
6	L	70	Total 70	O 70	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




#### • Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN





#### • Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



#### • Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN





#### • Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN






- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain F: 




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain G: 




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain H: 



- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain I: 





- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain J: 




- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain K: 



- Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.96Å 137.42Å 141.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.91 – 2.30 24.91 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (24.91-2.30) 96.7 (24.91-2.30)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.62 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.4.0078	Depositor
R, $R_{free}$	0.177 , 0.240 0.190 , 0.248	Depositor DCC
$R_{free}$ test set	4425 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	37.3	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 35.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15464	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EPE, CO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	0/1242	0.64	0/1676
1	B	0.69	0/1242	0.67	0/1677
1	C	0.61	0/1218	0.63	0/1645
1	D	0.73	1/1257 (0.1%)	0.63	0/1696
1	E	0.62	0/1227	0.62	0/1658
1	F	0.69	0/1240	0.65	0/1674
1	G	0.61	0/1222	0.63	0/1651
1	H	0.69	0/1238	0.63	0/1672
1	I	0.63	0/1247	0.61	0/1684
1	J	0.71	0/1239	0.64	0/1674
1	K	0.71	0/1274	0.64	0/1719
1	L	0.67	0/1241	0.65	0/1676
All	All	0.67	1/14887 (0.0%)	0.64	0/20102

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	120	GLU	CG-CD	5.05	1.59	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1217	0	1174	10	0
1	B	1213	0	1176	10	0
1	C	1193	0	1155	12	0
1	D	1232	0	1195	12	0
1	E	1202	0	1153	16	0
1	F	1215	0	1179	11	0
1	G	1197	0	1152	9	0
1	H	1213	0	1173	6	0
1	I	1222	0	1179	15	0
1	J	1214	0	1168	12	0
1	K	1245	0	1200	16	0
1	L	1216	0	1176	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	1	0	0	0	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	F	2	0	0	0	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
5	E	15	0	17	1	0
5	I	15	0	17	0	0
5	J	15	0	17	0	0
5	K	15	0	17	0	0
6	A	63	0	0	2	0
6	B	93	0	0	2	0
6	C	49	0	0	2	0
6	D	89	0	0	3	0
6	E	44	0	0	1	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	F	72	0	0	2	0
6	G	52	0	0	0	0
6	H	54	0	0	0	0
6	I	67	0	0	2	0
6	J	73	0	0	0	0
6	K	76	0	0	0	0
6	L	70	0	0	1	0
All	All	15464	0	14148	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 129 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:117[B]:ARG:HH21	1:I:117[B]:ARG:CG	1.68	1.06
1:C:130:GLN:NE2	1:C:152:LYS:HE3	1.78	0.99
1:I:117[B]:ARG:HG2	1:I:117[B]:ARG:HH21	0.79	0.94
1:D:130:GLN:NE2	1:D:152:LYS:HE3	1.83	0.94
1:I:117[B]:ARG:NH2	1:I:117[B]:ARG:HG2	1.61	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/165 (91%)	148 (99%)	2 (1%)	0	100	100
1	B	151/165 (92%)	149 (99%)	2 (1%)	0	100	100
1	C	148/165 (90%)	146 (99%)	2 (1%)	0	100	100
1	D	152/165 (92%)	151 (99%)	1 (1%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	149/165 (90%)	147 (99%)	2 (1%)	0	100	100
1	F	151/165 (92%)	149 (99%)	2 (1%)	0	100	100
1	G	148/165 (90%)	147 (99%)	1 (1%)	0	100	100
1	H	150/165 (91%)	148 (99%)	2 (1%)	0	100	100
1	I	151/165 (92%)	150 (99%)	1 (1%)	0	100	100
1	J	150/165 (91%)	148 (99%)	2 (1%)	0	100	100
1	K	154/165 (93%)	153 (99%)	1 (1%)	0	100	100
1	L	151/165 (92%)	150 (99%)	1 (1%)	0	100	100
All	All	1805/1980 (91%)	1786 (99%)	19 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/141 (92%)	127 (98%)	3 (2%)	56	73
1	B	130/141 (92%)	126 (97%)	4 (3%)	45	61
1	C	126/141 (89%)	123 (98%)	3 (2%)	54	72
1	D	132/141 (94%)	127 (96%)	5 (4%)	38	52
1	E	127/141 (90%)	122 (96%)	5 (4%)	37	51
1	F	128/141 (91%)	126 (98%)	2 (2%)	68	82
1	G	127/141 (90%)	119 (94%)	8 (6%)	21	28
1	H	129/141 (92%)	127 (98%)	2 (2%)	68	82
1	I	130/141 (92%)	127 (98%)	3 (2%)	56	73
1	J	129/141 (92%)	123 (95%)	6 (5%)	30	41
1	K	134/141 (95%)	131 (98%)	3 (2%)	57	74
1	L	129/141 (92%)	125 (97%)	4 (3%)	45	61
All	All	1551/1692 (92%)	1503 (97%)	48 (3%)	45	61

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	117	ARG
1	G	95	SER
1	L	44	HIS
1	G	44	HIS
1	G	69	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	44	HIS
1	G	130	GLN
1	L	99	GLN
1	F	130	GLN
1	F	163	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 23 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EPE	E	1173	-	15,15,15	0.92	1 (6%)	18,20,20	2.85	5 (27%)
5	EPE	I	1173	-	15,15,15	0.87	1 (6%)	18,20,20	2.51	8 (44%)
5	EPE	J	1173	-	15,15,15	0.79	1 (6%)	18,20,20	2.82	9 (50%)
5	EPE	K	1173	-	15,15,15	0.87	1 (6%)	18,20,20	2.61	7 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EPE	E	1173	-	-	0/9/19/19	0/1/1/1
5	EPE	I	1173	-	-	0/9/19/19	0/1/1/1
5	EPE	J	1173	-	-	0/9/19/19	0/1/1/1
5	EPE	K	1173	-	-	0/9/19/19	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	1173	EPE	C10-S	2.58	1.81	1.77
5	I	1173	EPE	C10-S	2.66	1.81	1.77
5	K	1173	EPE	C10-S	2.77	1.81	1.77
5	E	1173	EPE	C10-S	3.21	1.82	1.77

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	J	1173	EPE	C5-C6-N1	-3.01	104.54	110.63
5	I	1173	EPE	O3S-S-O2S	-2.98	104.53	111.37
5	K	1173	EPE	O2S-S-O1S	-2.90	103.81	113.86
5	J	1173	EPE	O3S-S-O2S	-2.67	105.26	111.37
5	E	1173	EPE	O2S-S-O1S	-2.39	105.58	113.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	1173	EPE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	150/165 (90%)	-0.13	3 (2%) 65 72	28, 37, 50, 58	0
1	B	152/165 (92%)	-0.05	3 (1%) 65 72	22, 30, 46, 79	0
1	C	150/165 (90%)	0.02	2 (1%) 77 81	31, 39, 55, 61	0
1	D	152/165 (92%)	-0.08	2 (1%) 77 81	22, 30, 44, 61	0
1	E	150/165 (90%)	0.02	3 (2%) 65 72	32, 42, 57, 62	0
1	F	151/165 (91%)	-0.16	2 (1%) 77 81	22, 30, 45, 65	0
1	G	150/165 (90%)	-0.04	2 (1%) 77 81	30, 39, 56, 61	0
1	H	151/165 (91%)	0.01	3 (1%) 65 72	27, 36, 53, 63	0
1	I	151/165 (91%)	-0.04	7 (4%) 33 40	25, 37, 56, 75	0
1	J	151/165 (91%)	-0.08	4 (2%) 56 63	21, 35, 50, 61	0
1	K	152/165 (92%)	-0.18	1 (0%) 87 90	24, 32, 44, 52	0
1	L	151/165 (91%)	-0.17	2 (1%) 77 81	24, 34, 48, 55	0
All	All	1811/1980 (91%)	-0.07	34 (1%) 67 73	21, 36, 53, 79	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	22	LEU	6.6
1	B	22	LEU	6.0
1	F	23	ALA	5.7
1	G	23	ALA	4.2
1	H	23	ALA	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	EPE	K	1173	15/15	0.96	0.18	2.23	22,34,44,46	0
5	EPE	E	1173	15/15	0.93	0.21	2.17	26,39,46,48	0
3	CL	A	1174	1/1	0.97	0.20	2.11	36,36,36,36	0
2	CO	C	1173	1/1	0.97	0.17	1.43	48,48,48,48	0
3	CL	J	1175	1/1	0.93	0.15	1.33	37,37,37,37	0
5	EPE	I	1173	15/15	0.96	0.15	1.23	26,36,40,41	0
2	CO	I	1174	1/1	0.97	0.15	0.87	46,46,46,46	0
4	CA	E	1175	1/1	0.62	0.24	0.58	53,53,53,53	0
2	CO	B	1173	1/1	0.98	0.14	0.36	42,42,42,42	0
2	CO	E	1174	1/1	0.99	0.15	0.33	48,48,48,48	0
2	CO	D	1173	1/1	0.97	0.15	0.29	36,36,36,36	0
2	CO	G	1173	1/1	0.97	0.15	0.21	49,49,49,49	0
2	CO	F	1173	1/1	0.98	0.14	0.08	33,33,33,33	0
2	CO	A	1173	1/1	0.98	0.14	-0.31	42,42,42,42	0
4	CA	H	1176	1/1	0.95	0.13	-0.46	45,45,45,45	0
5	EPE	J	1173	15/15	0.97	0.12	-0.54	27,34,42,43	0
2	CO	H	1173	1/1	0.97	0.12	-0.58	42,42,42,42	0
2	CO	J	1174	1/1	0.97	0.13	-0.60	50,50,50,50	0
2	CO	L	1173	1/1	0.99	0.11	-0.68	36,36,36,36	0
2	CO	K	1174	1/1	0.94	0.09	-2.24	43,43,43,43	0
4	CA	I	1176	1/1	0.82	0.30	-	41,41,41,41	0
3	CL	I	1175	1/1	0.97	0.32	-	31,31,31,31	0
4	CA	F	1174	1/1	0.98	0.10	-	15,15,15,15	0
4	CA	B	1174	1/1	0.95	0.27	-	41,41,41,41	0
4	CA	J	1176	1/1	0.93	0.16	-	54,54,54,54	0
4	CA	C	1174	1/1	0.88	0.25	-	58,58,58,58	0
4	CA	F	1175	1/1	0.87	0.18	-	45,45,45,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.