



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2017 – 01:27 PM EDT

PDB ID : 5XJB
Title : The Crystal Structure of the Minimal Core Domain of the Microtubule Depolymerizer KIF2C Complexed with ADP-Mg-BeFx
Authors : Ogawa, T.; Jiang, X.; Hirokawa, N.
Deposited on : unknown
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1001 (3.12-3.08)
Clashscore	112137	1099 (3.12-3.08)
Ramachandran outliers	110173	1057 (3.12-3.08)
Sidechain outliers	110143	1057 (3.12-3.08)
RSRZ outliers	101464	1006 (3.12-3.08)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	426	 <p>4% 70% 8% 20%</p>
1	B	426	 <p>2% 71% 7% 22%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BEF	A	602	-	-	X	X
3	BEF	B	602	-	-	X	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kinesin-like protein KIF2C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	0	0
			2663	1678	472	492	21			
1	B	333	Total	C	N	O	S	0	0	0
			2612	1650	460	481	21			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	167	MET	-	initiating methionine	UNP Q922S8
A	168	ALA	-	expression tag	UNP Q922S8
A	169	SER	-	expression tag	UNP Q922S8
A	170	MET	-	expression tag	UNP Q922S8
A	171	THR	-	expression tag	UNP Q922S8
A	172	GLY	-	expression tag	UNP Q922S8
A	173	GLY	-	expression tag	UNP Q922S8
A	174	GLN	-	expression tag	UNP Q922S8
A	175	GLN	-	expression tag	UNP Q922S8
A	176	MET	-	expression tag	UNP Q922S8
A	177	GLY	-	expression tag	UNP Q922S8
A	178	ARG	-	expression tag	UNP Q922S8
A	179	ASP	-	expression tag	UNP Q922S8
A	180	PRO	-	expression tag	UNP Q922S8
A	181	ASN	-	expression tag	UNP Q922S8
A	182	SER	-	expression tag	UNP Q922S8
A	183	SER	-	expression tag	UNP Q922S8
A	586	HIS	-	expression tag	UNP Q922S8
A	587	HIS	-	expression tag	UNP Q922S8
A	588	HIS	-	expression tag	UNP Q922S8
A	589	HIS	-	expression tag	UNP Q922S8
A	590	HIS	-	expression tag	UNP Q922S8
A	591	HIS	-	expression tag	UNP Q922S8
A	592	HIS	-	expression tag	UNP Q922S8
B	167	MET	-	initiating methionine	UNP Q922S8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	168	ALA	-	expression tag	UNP Q922S8
B	169	SER	-	expression tag	UNP Q922S8
B	170	MET	-	expression tag	UNP Q922S8
B	171	THR	-	expression tag	UNP Q922S8
B	172	GLY	-	expression tag	UNP Q922S8
B	173	GLY	-	expression tag	UNP Q922S8
B	174	GLN	-	expression tag	UNP Q922S8
B	175	GLN	-	expression tag	UNP Q922S8
B	176	MET	-	expression tag	UNP Q922S8
B	177	GLY	-	expression tag	UNP Q922S8
B	178	ARG	-	expression tag	UNP Q922S8
B	179	ASP	-	expression tag	UNP Q922S8
B	180	PRO	-	expression tag	UNP Q922S8
B	181	ASN	-	expression tag	UNP Q922S8
B	182	SER	-	expression tag	UNP Q922S8
B	183	SER	-	expression tag	UNP Q922S8
B	586	HIS	-	expression tag	UNP Q922S8
B	587	HIS	-	expression tag	UNP Q922S8
B	588	HIS	-	expression tag	UNP Q922S8
B	589	HIS	-	expression tag	UNP Q922S8
B	590	HIS	-	expression tag	UNP Q922S8
B	591	HIS	-	expression tag	UNP Q922S8
B	592	HIS	-	expression tag	UNP Q922S8

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Be	F	0	0
			4	1	3		
3	B	1	Total	Be	F	0	0
			4	1	3		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	O	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	O	0	0
			2	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	89.45Å 166.59Å 75.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.10 29.48 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-3.10) 99.9 (29.48-3.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 3.11Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.201 , 0.261 0.209 , 0.256	Depositor DCC
R_{free} test set	1061 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	72.8	Xtriage
Anisotropy	0.755	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5344	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEF, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	3/2700 (0.1%)	1.05	12/3627 (0.3%)
1	B	0.61	0/2649	0.88	5/3558 (0.1%)
All	All	0.63	3/5349 (0.1%)	0.97	17/7185 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	454	SER	CA-CB	-5.82	1.44	1.52
1	A	455	GLY	CA-C	5.66	1.60	1.51
1	A	529	ASN	C-N	5.02	1.45	1.34

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	529	ASN	CB-CA-C	-19.35	71.70	110.40
1	A	529	ASN	O-C-N	13.54	144.37	122.70
1	A	529	ASN	N-CA-C	-13.42	74.76	111.00
1	A	529	ASN	CA-C-N	-11.37	92.20	117.20
1	A	535	PHE	CB-CA-C	-10.12	90.16	110.40
1	A	535	PHE	N-CA-C	9.83	137.53	111.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	534	PRO	N-CA-CB	7.71	112.56	103.30
1	B	466	SER	N-CA-CB	-7.60	99.10	110.50
1	A	465	ARG	N-CA-C	-7.04	91.98	111.00
1	A	529	ASN	C-N-CA	6.98	139.15	121.70
1	A	534	PRO	C-N-CA	-6.21	106.18	121.70
1	A	465	ARG	CB-CA-C	6.17	122.75	110.40
1	B	534	PRO	N-CA-CB	5.74	110.18	103.30
1	B	493	GLU	CB-CA-C	-5.32	99.76	110.40
1	A	488	ASP	CB-CG-OD1	5.16	122.94	118.30
1	B	545	ARG	CG-CD-NE	5.07	122.45	111.80
1	B	545	ARG	NE-CZ-NH2	-5.03	117.79	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	535	PHE	Peptide
1	A	584	LEU	Peptide
1	B	534	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2663	0	2676	36	1
1	B	2612	0	2637	23	1
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	4	0	0	3	0
3	B	4	0	0	3	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	3	0	0	1	0
5	B	2	0	0	0	0
All	All	5344	0	5337	57	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:464:SER:N	3:B:602:BEF:F3	1.80	1.04
1:A:528:GLN:O	1:A:530:LYS:CB	2.13	0.97
1:A:528:GLN:O	1:A:529:ASN:C	2.03	0.96
1:A:463:SER:N	1:A:465:ARG:HG2	1.85	0.91
1:B:463:SER:OG	3:B:602:BEF:F3	1.75	0.91
1:A:528:GLN:O	1:A:530:LYS:N	2.12	0.81
1:A:464:SER:OG	3:A:602:BEF:F3	1.88	0.81
1:B:463:SER:OG	1:B:464:SER:N	2.20	0.71
1:B:463:SER:HG	1:B:464:SER:H	1.41	0.69
1:A:476:THR:HG23	1:A:477:LYS:H	1.59	0.68
1:A:463:SER:N	1:A:465:ARG:CG	2.56	0.66
1:A:283:CYS:SG	5:A:701:HOH:O	2.54	0.65
1:A:476:THR:HG23	1:A:477:LYS:N	2.12	0.65
1:A:464:SER:N	3:A:602:BEF:F1	2.24	0.60
1:B:464:SER:CA	3:B:602:BEF:F3	2.38	0.59
1:B:463:SER:O	1:B:464:SER:HB2	2.03	0.58
1:A:463:SER:OG	3:A:602:BEF:F1	2.00	0.54
1:A:477:LYS:C	1:A:479:ARG:H	2.10	0.53
1:A:476:THR:CG2	1:A:477:LYS:H	2.22	0.52
1:A:463:SER:N	1:A:465:ARG:HE	2.08	0.51
1:B:524:ARG:HG2	1:B:584:LEU:HB2	1.94	0.50
1:A:528:GLN:O	1:A:530:LYS:CA	2.60	0.50
1:B:369:ILE:HG22	1:B:486:LEU:HD23	1.94	0.49
1:A:533:THR:O	1:A:534:PRO:C	2.49	0.49
1:A:369:ILE:HG22	1:A:486:LEU:HD23	1.94	0.48
1:B:409:LEU:HD22	1:B:444:MET:HE3	1.96	0.48
1:B:385:ARG:HG3	1:B:386:TYR:CD1	2.49	0.48
1:A:366:SER:HB3	1:A:442:ILE:HD13	1.95	0.48
1:A:476:THR:CG2	1:A:477:LYS:N	2.77	0.47
1:B:433:TYR:CG	1:B:444:MET:HE2	2.50	0.47
1:A:463:SER:N	1:A:465:ARG:NE	2.63	0.47
1:A:249:PRO:O	1:A:250:ILE:HB	2.15	0.47
1:B:385:ARG:NE	1:B:386:TYR:CE1	2.82	0.47
1:B:248:ASP:OD1	1:B:249:PRO:O	2.34	0.46
1:A:539:LYS:O	1:A:540:LEU:C	2.51	0.46
1:B:463:SER:HG	1:B:464:SER:N	2.04	0.46
1:A:524:ARG:HG2	1:A:584:LEU:HB2	1.98	0.46
1:A:380:LEU:HD11	1:B:250:ILE:HA	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:244:LEU:O	1:B:248:ASP:HB2	2.15	0.45
1:B:241:CYS:SG	1:B:245:THR:OG1	2.51	0.45
1:A:248:ASP:OD1	1:A:249:PRO:O	2.35	0.45
1:A:249:PRO:O	1:A:250:ILE:CB	2.64	0.45
1:A:454:SER:OG	1:A:454:SER:O	2.30	0.45
1:A:439:ASP:O	1:A:443:LYS:HG2	2.17	0.44
1:A:244:LEU:O	1:A:248:ASP:HB2	2.17	0.44
1:B:452:ARG:CZ	1:B:463:SER:OG	2.65	0.44
1:B:249:PRO:O	1:B:250:ILE:HB	2.18	0.44
1:A:249:PRO:HB2	1:B:330:GLN:OE1	2.18	0.44
1:A:453:THR:HG22	1:A:454:SER:N	2.33	0.43
1:B:289:GLU:OE1	1:B:303:GLN:NE2	2.53	0.42
1:B:249:PRO:O	1:B:250:ILE:CB	2.68	0.42
1:A:289:GLU:OE1	1:A:303:GLN:NE2	2.54	0.41
1:A:437:CYS:HB3	1:A:440:ASP:CG	2.41	0.41
1:A:463:SER:N	1:A:465:ARG:CD	2.84	0.40
1:A:279:VAL:HB	1:A:312:ASP:HB3	2.02	0.40
1:A:463:SER:C	1:A:465:ARG:HG2	2.41	0.40
1:B:366:SER:HB3	1:B:442:ILE:HD12	2.02	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:301:GLU:OE2	1:A:301:GLU:OE2[2_655]	1.78	0.42
1:B:301:GLU:OE2	1:B:301:GLU:OE2[2_655]	1.88	0.32

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	331/426 (78%)	300 (91%)	30 (9%)	1 (0%)	44 79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	323/426 (76%)	304 (94%)	18 (6%)	1 (0%)	44	79
All	All	654/852 (77%)	604 (92%)	48 (7%)	2 (0%)	44	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	250	ILE
1	B	250	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/374 (77%)	283 (98%)	5 (2%)	66	88
1	B	284/374 (76%)	280 (99%)	4 (1%)	71	90
All	All	572/748 (76%)	563 (98%)	9 (2%)	68	89

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	LEU
1	A	339	THR
1	A	419	GLU
1	A	464	SER
1	A	484	PHE
1	B	286	LEU
1	B	339	THR
1	B	397	PHE
1	B	484	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	303	GLN
1	A	390	ASN
1	A	467	HIS
1	B	303	GLN
1	B	390	ASN
1	B	467	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ADP	A	601	4	25,29,29	1.22	3 (12%)	24,45,45	1.81	5 (20%)
3	BEF	A	602	-	0,3,3	0.00	-	0,3,3	0.00	-
2	ADP	B	601	4	25,29,29	1.14	3 (12%)	24,45,45	1.91	5 (20%)
3	BEF	B	602	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	601	4	-	0/12/32/32	0/3/3/3
3	BEF	A	602	-	-	0/0/0/0	0/0/0/0
2	ADP	B	601	4	-	0/12/32/32	0/3/3/3
3	BEF	B	602	-	-	0/0/0/0	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	ADP	C2-N3	2.19	1.35	1.32
2	A	601	ADP	C2-N3	2.34	1.36	1.32
2	B	601	ADP	O4'-C1'	2.63	1.44	1.41
2	B	601	ADP	C5-C4	3.13	1.47	1.40
2	A	601	ADP	O4'-C1'	3.19	1.45	1.41
2	A	601	ADP	C5-C4	3.26	1.47	1.40

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	ADP	N3-C2-N1	-6.16	123.49	128.86
2	A	601	ADP	N3-C2-N1	-5.89	123.73	128.86
2	B	601	ADP	C5'-C4'-C3'	-3.58	101.64	115.29
2	A	601	ADP	O5'-PA-O1A	-2.59	98.80	109.25
2	B	601	ADP	C4-C5-N7	-2.32	107.17	109.41
2	A	601	ADP	C5'-C4'-C3'	-2.06	107.43	115.29
2	B	601	ADP	O3B-PB-O2B	2.16	116.31	107.61
2	A	601	ADP	O3B-PB-O2B	2.17	116.37	107.61
2	A	601	ADP	N6-C6-N1	2.46	123.64	118.77
2	B	601	ADP	N6-C6-N1	2.66	124.04	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	BEF	3	0
3	B	602	BEF	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/426 (80%)	0.02	17 (4%) 30 13	30, 73, 127, 165	0
1	B	333/426 (78%)	-0.12	7 (2%) 64 43	46, 74, 122, 144	0
All	All	674/852 (79%)	-0.05	24 (3%) 43 21	30, 74, 126, 165	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	SER	4.7
1	A	528	GLN	4.2
1	A	455	GLY	4.2
1	A	456	GLN	3.7
1	A	388	ASN	3.5
1	A	532	HIS	3.4
1	A	364	ASN	3.2
1	A	533	THR	3.2
1	A	454	SER	3.1
1	A	529	ASN	3.1
1	A	534	PRO	3.0
1	A	240	GLU	3.0
1	A	497	ASP	2.8
1	B	357	ASP	2.2
1	B	363	GLN	2.2
1	A	241	CYS	2.2
1	A	418	LEU	2.1
1	A	390	ASN	2.1
1	B	551	GLU	2.1
1	B	421	SER	2.1
1	B	387	ARG	2.0
1	A	496	ALA	2.0
1	B	485	SER	2.0
1	B	451	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	BEF	B	602	4/4	0.54	0.53	5.98	104,122,124,127	0
3	BEF	A	602	4/4	0.58	0.38	3.59	102,110,114,115	0
2	ADP	A	601	27/27	0.95	0.15	-0.44	67,78,99,107	0
2	ADP	B	601	27/27	0.96	0.14	-0.71	67,84,99,119	0
4	MG	A	603	1/1	0.82	0.10	-	66,66,66,66	0
4	MG	B	603	1/1	0.73	0.13	-	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.