



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 30, 2017 – 04:28 PM EDT

PDB ID : 4XLA
Title : Tailspike protein mutant D339A of E. coli bacteriophage HK620 IN COMPLEX WITH PENTASACCHARIDE
Authors : Gohlke, U.; Broecker, N.K.; Heinemann, U.; Seckler, R.; Barbirz, S.
Deposited on : unknown
Resolution : 1.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3517 (1.50-1.46)
Clashscore	112137	3795 (1.50-1.46)
Ramachandran outliers	110173	3721 (1.50-1.46)
Sidechain outliers	110143	3719 (1.50-1.46)
RSRZ outliers	101464	3549 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	600	<div> <div style="width: 2%; height: 10px; background-color: red;"></div> <div style="width: 90%; height: 10px; background-color: green;"></div> <div style="width: 9%; height: 10px; background-color: yellow;"></div> <div style="width: 1%; height: 10px; background-color: orange;"></div> </div> <div>2% 90% 9%</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	TRS	A	1006	-	-	-	X
7	FMT	A	1008	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	NA	A	1012	-	-	-	X

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tail spike protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	600	4716	2944	817	931	24	0	22	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	339	ALA	ASP	engineered mutation	UNP Q9AYY6

- Molecule 2 is ALPHA-L-RHAMNOSE (three-letter code: RAM) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	10	6	4	0	0

- Molecule 3 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is ALPHA D-GALACTOSE (three-letter code: GLA) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			12	6	6		

- Molecule 5 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 3 1 2	0	0
7	A	1	Total C O 3 1 2	0	0
7	A	1	Total C O 3 1 2	0	0
7	A	1	Total C O 3 1 2	0	0

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	2	Total Na 2 2	0	0


- Molecule 9 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	829	Total O 829 829	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tail spike protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	74.21Å 74.21Å 174.65Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.66 – 1.47 43.66 – 1.47	Depositor EDS
% Data completeness (in resolution range)	99.3 (43.66-1.47) 99.4 (43.66-1.47)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 1.47Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.140 , 0.167 0.141 , 0.167	Depositor DCC
R_{free} test set	4760 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	13.3	Xtriage
Anisotropy	0.311	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	5628	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RAM, NAG, GLA, FMT, GLC, NA, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	5/4825 (0.1%)	0.99	12/6573 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	664	SER	CB-OG	-5.66	1.34	1.42
1	A	353	GLU	CB-CG	-5.33	1.42	1.52
1	A	667	GLU	CD-OE2	5.24	1.31	1.25
1	A	597	TYR	CE2-CZ	-5.09	1.31	1.38
1	A	322	TYR	CG-CD2	-5.04	1.32	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	ARG	NE-CZ-NH2	-9.71	115.45	120.30
1	A	390	ARG	NE-CZ-NH1	8.04	124.32	120.30
1	A	229	ARG	NE-CZ-NH1	7.98	124.29	120.30
1	A	324	ARG	NE-CZ-NH2	-7.05	116.77	120.30
1	A	327	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	A	324	ARG	NE-CZ-NH1	6.18	123.39	120.30
1	A	327	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	A	641	THR	CA-CB-CG2	-5.77	104.32	112.40
1	A	670	ASP	CB-CG-OD2	-5.77	113.11	118.30
1	A	482	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	560	PHE	CB-CG-CD2	-5.66	116.84	120.80
1	A	636	LEU	CB-CG-CD1	-5.30	101.98	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4716	0	4422	40	0
2	A	10	0	8	0	0
3	A	11	0	9	0	0
4	A	12	0	9	0	0
5	A	28	0	26	0	0
6	A	8	0	12	0	0
7	A	12	0	3	0	0
8	A	2	0	0	1	0
9	A	829	0	0	9	8
All	All	5628	0	4489	41	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:381[B]:ARG:CD	1:A:414[B]:GLU:OE2	1.76	1.30
1:A:381[B]:ARG:HD3	1:A:414[B]:GLU:OE2	1.47	1.13
1:A:381[B]:ARG:HD2	1:A:414[B]:GLU:OE2	1.42	1.12
1:A:532:SER:OG	9:A:1102:HOH:O	1.81	0.95
1:A:678[B]:THR:OG1	9:A:1103:HOH:O	1.87	0.91
1:A:381[B]:ARG:HG2	1:A:414[B]:GLU:HG3	1.52	0.91
1:A:381[A]:ARG:NH2	9:A:1105:HOH:O	2.16	0.77
1:A:414[A]:GLU:HG2	1:A:445[A]:SER:OG	1.86	0.76
1:A:381[B]:ARG:HG2	1:A:414[B]:GLU:CG	2.17	0.73
1:A:528:PRO:HD3	9:A:1276:HOH:O	1.89	0.73
8:A:1011:NA:NA	9:A:1137:HOH:O	1.67	0.65
1:A:414[B]:GLU:OE1	1:A:415[B]:ASN:ND2	2.25	0.63
1:A:381[B]:ARG:CD	1:A:414[B]:GLU:CD	2.67	0.60
1:A:409:TYR:HA	1:A:440:ASN:O	2.01	0.59
1:A:517:ASN:OD1	9:A:1106:HOH:O	2.18	0.57
1:A:667:GLU:HG2	9:A:1314:HOH:O	2.03	0.57
1:A:381[B]:ARG:HD2	1:A:414[B]:GLU:CD	2.23	0.54
1:A:381[B]:ARG:NH1	9:A:1105:HOH:O	2.07	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303[B]:ASN:ND2	9:A:1107:HOH:O	2.45	0.48
1:A:371:VAL:O	1:A:392:ALA:HA	2.14	0.48
1:A:248:ASP:OD1	1:A:248:ASP:C	2.53	0.47
1:A:440:ASN:HA	1:A:481:GLN:O	2.14	0.46
1:A:241:PHE:O	1:A:283:ARG:HA	2.15	0.46
1:A:579:GLU:HG2	1:A:581:LEU:HG	1.98	0.45
1:A:626:VAL:HG12	1:A:628[A]:VAL:HG13	1.99	0.45
1:A:381[A]:ARG:HA	1:A:414[A]:GLU:O	2.17	0.45
1:A:158:ASP:HB2	1:A:159:PRO:CD	2.47	0.44
1:A:621:GLN:HB2	1:A:628[B]:VAL:HG11	2.00	0.44
1:A:414[B]:GLU:HA	1:A:445[B]:SER:O	2.19	0.42
1:A:294:CYS:O	1:A:321:CYS:HA	2.19	0.42
1:A:600:ILE:O	1:A:628[B]:VAL:HA	2.19	0.42
1:A:297:THR:HA	1:A:324:ARG:O	2.20	0.41
1:A:465:ILE:O	1:A:502:ILE:HA	2.20	0.41
1:A:381[B]:ARG:HD3	1:A:381[B]:ARG:HH11	1.72	0.41
1:A:110:ASP:O	1:A:114:PHE:N	2.54	0.41
1:A:381[B]:ARG:NH1	1:A:412:GLN:OE1	2.51	0.41
1:A:288:PHE:HB3	1:A:291:SER:CB	2.50	0.41
1:A:537:ASN:HA	1:A:569:ASN:O	2.20	0.41
1:A:475:SER:HB2	1:A:506:ALA:HA	2.03	0.41
1:A:449:VAL:O	1:A:490:PHE:HA	2.21	0.40
1:A:293:ASN:HA	1:A:320:ASN:O	2.21	0.40

All (8) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:1161:HOH:O	9:A:1214:HOH:O[2_655]	1.32	0.88
9:A:1263:HOH:O	9:A:1788:HOH:O[6_766]	1.75	0.45
9:A:1160:HOH:O	9:A:1567:HOH:O[3_665]	1.75	0.45
9:A:1149:HOH:O	9:A:1546:HOH:O[2_655]	1.79	0.41
9:A:1108:HOH:O	9:A:1532:HOH:O[2_655]	1.97	0.23
9:A:1407:HOH:O	9:A:1479:HOH:O[3_665]	2.02	0.18
9:A:1313:HOH:O	9:A:1398:HOH:O[2_655]	2.02	0.18
9:A:1152:HOH:O	9:A:1698:HOH:O[2_655]	2.03	0.17

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	620/600 (103%)	601 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	513/491 (104%)	506 (99%)	7 (1%)	71	43

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	ASP
1	A	149	ARG
1	A	440	ASN
1	A	517	ASN
1	A	611[A]	SER
1	A	611[B]	SER
1	A	641	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	RAM	A	1001	8,3	9,10,11	1.17	1 (11%)	13,14,16	1.63	2 (15%)
3	GLC	A	1002	2,4	11,11,12	1.10	0	13,15,17	1.67	3 (23%)
4	GLA	A	1003	3,5	12,12,12	1.08	0	17,17,17	1.76	3 (17%)
5	NAG	A	1004	4	14,14,15	0.78	0	15,19,21	1.17	2 (13%)
5	NAG	A	1005	4	14,14,15	1.53	2 (14%)	16,19,21	1.44	2 (12%)
6	TRS	A	1006	-	7,7,7	1.03	0	9,9,9	1.71	3 (33%)
7	FMT	A	1007	-	0,2,2	0.00	-	0,1,1	0.00	-
7	FMT	A	1008	-	0,2,2	0.00	-	0,1,1	0.00	-
7	FMT	A	1009	-	0,2,2	0.00	-	0,1,1	0.00	-
7	FMT	A	1010	1	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RAM	A	1001	8,3	-	0/0/17/20	0/1/1/1
3	GLC	A	1002	2,4	-	0/2/19/22	0/1/1/1
4	GLA	A	1003	3,5	-	0/2/22/22	0/1/1/1
5	NAG	A	1004	4	-	0/6/23/26	0/1/1/1
5	NAG	A	1005	4	-	0/6/22/26	0/1/1/1
6	TRS	A	1006	-	-	0/9/9/9	0/0/0/0
7	FMT	A	1007	-	-	0/0/0/0	0/0/0/0
7	FMT	A	1008	-	-	0/0/0/0	0/0/0/0
7	FMT	A	1009	-	-	0/0/0/0	0/0/0/0
7	FMT	A	1010	1	-	0/0/0/0	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1005	NAG	O5-C5	-2.36	1.38	1.44
2	A	1001	RAM	O3-C3	2.37	1.48	1.43
5	A	1005	NAG	C4-C5	3.63	1.58	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1003	GLA	O3-C3-C2	-3.84	102.01	110.36
2	A	1001	RAM	O4-C4-C3	-3.63	102.45	110.36
4	A	1003	GLA	O1-C1-O5	-2.82	101.85	110.20
2	A	1001	RAM	O2-C2-C3	-2.66	104.95	110.17
3	A	1002	GLC	O3-C3-C2	-2.40	105.66	110.02
5	A	1004	NAG	O4-C4-C3	-2.01	105.98	110.36
5	A	1005	NAG	C2-N2-C7	2.02	125.88	122.94
6	A	1006	TRS	C3-C-N	2.12	112.24	107.73
4	A	1003	GLA	O5-C1-C2	2.30	113.85	110.04
6	A	1006	TRS	O3-C3-C	2.37	117.33	110.47
5	A	1004	NAG	C4-C3-C2	2.59	114.81	111.02
3	A	1002	GLC	C1-C2-C3	2.68	113.05	109.65
5	A	1005	NAG	C3-C4-C5	2.91	114.65	110.82
6	A	1006	TRS	C2-C-N	3.17	114.46	107.73
3	A	1002	GLC	C1-O5-C5	3.58	117.10	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	600/600 (100%)	-0.15	11 (1%) 69 73	10, 14, 24, 65	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	529	GLY	6.3
1	A	110	ASP	5.7
1	A	112	ASP	4.3
1	A	113	GLN	3.6
1	A	433	ALA	3.4
1	A	434	THR	3.3
1	A	114	PHE	3.0
1	A	111	PRO	2.9
1	A	530	THR	2.6
1	A	528	PRO	2.4
1	A	531	THR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
7	FMT	A	1008	3/3	0.96	0.15	4.09	26,26,30,34	0
6	TRS	A	1006	8/8	0.96	0.10	2.60	13,15,18,19	0
8	NA	A	1012	1/1	0.98	0.14	2.09	18,18,18,18	0
5	NAG	A	1005	14/15	0.97	0.07	0.44	12,14,15,15	0
5	NAG	A	1004	14/15	0.97	0.07	-0.08	13,15,27,27	0
4	GLA	A	1003	12/12	0.97	0.06	-0.36	14,14,16,17	0
2	RAM	A	1001	10/11	0.94	0.07	-0.48	17,19,20,23	0
7	FMT	A	1007	3/3	0.98	0.06	-0.79	15,15,15,18	0
8	NA	A	1011	1/1	0.91	0.14	-	25,25,25,25	0
3	GLC	A	1002	11/12	0.96	0.08	-	15,17,20,21	0
7	FMT	A	1009	3/3	0.99	0.04	-	20,20,22,23	0
7	FMT	A	1010	3/3	0.91	0.23	-	11,11,15,18	3

6.5 Other polymers [i](#)

There are no such residues in this entry.