



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:51 am GMT

PDB ID : 1XPQ
Title : Crystal structure of fms1, a polyamine oxidase from yeast
Authors : Huang, Q.; Liu, Q.; Hao, Q.
Deposited on : 2004-10-09
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2895 (2.60-2.60)
Ramachandran outliers	110173	2848 (2.60-2.60)
Sidechain outliers	110143	2848 (2.60-2.60)
RSRZ outliers	101464	2550 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	513	 55% 38% . . .
1	B	513	 49% 43% . .
1	C	513	 54% 39% . .
1	D	513	 50% 41% . . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FMS1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	498	Total	C	N	O	S	86	0	0
			3997	2526	699	750	22			
1	B	492	Total	C	N	O	S	144	0	0
			3946	2496	687	741	22			
1	C	495	Total	C	N	O	S	118	0	0
			3967	2509	690	746	22			
1	D	490	Total	C	N	O	S	144	0	0
			3929	2484	684	739	22			

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	509	LEU	-	CLONING ARTIFACT	UNP P50264
A	510	GLU	-	CLONING ARTIFACT	UNP P50264
A	511	HIS	-	CLONING ARTIFACT	UNP P50264
A	512	HIS	-	CLONING ARTIFACT	UNP P50264
A	513	HIS	-	CLONING ARTIFACT	UNP P50264
B	509	LEU	-	CLONING ARTIFACT	UNP P50264
B	510	GLU	-	CLONING ARTIFACT	UNP P50264
B	511	HIS	-	CLONING ARTIFACT	UNP P50264
B	512	HIS	-	CLONING ARTIFACT	UNP P50264
B	513	HIS	-	CLONING ARTIFACT	UNP P50264
C	509	LEU	-	CLONING ARTIFACT	UNP P50264
C	510	GLU	-	CLONING ARTIFACT	UNP P50264
C	511	HIS	-	CLONING ARTIFACT	UNP P50264
C	512	HIS	-	CLONING ARTIFACT	UNP P50264
C	513	HIS	-	CLONING ARTIFACT	UNP P50264
D	509	LEU	-	CLONING ARTIFACT	UNP P50264
D	510	GLU	-	CLONING ARTIFACT	UNP P50264
D	511	HIS	-	CLONING ARTIFACT	UNP P50264
D	512	HIS	-	CLONING ARTIFACT	UNP P50264
D	513	HIS	-	CLONING ARTIFACT	UNP P50264

- Molecule 2 is SPERMINE (three-letter code: SPM) (formula: $C_{10}H_{26}N_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	N	14	0
			14	10	4		
2	D	1	Total	C	N	14	0
			14	10	4		
2	B	1	Total	C	N	14	0
			14	10	4		
2	A	1	Total	C	N	14	0
			14	10	4		


- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	55	Total	O	0	0
			55	55		
4	B	54	Total	O	0	0
			54	54		
4	C	66	Total	O	0	0
			66	66		
4	D	50	Total	O	0	0
			50	50		




• Molecule 1: FMS1 protein



• Molecule 1: FMS1 protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	103.05Å 98.13Å 123.83Å 90.00° 103.28° 90.00°	Depositor
Resolution (Å)	35.23 – 2.60 49.84 – 2.51	Depositor EDS
% Data completeness (in resolution range)	(Not available) (35.23-2.60) 73.5 (49.84-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.88 (at 2.51Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.232 , 0.309 0.225 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	43.7	Xtriage
Anisotropy	0.200	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	16332	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SPM, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/4080	0.64	0/5515
1	B	0.39	0/4025	0.65	0/5441
1	C	0.40	0/4047	0.64	0/5471
1	D	0.38	0/4008	0.63	1/5419 (0.0%)
All	All	0.39	0/16160	0.64	1/21846 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	380	LEU	CA-CB-CG	5.92	128.90	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3997	0	3916	202	0
1	B	3946	0	3881	217	0
1	C	3967	0	3900	159	0
1	D	3929	0	3857	207	0
2	A	14	0	26	0	0
2	B	14	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	14	0	26	0	0
2	D	14	0	26	0	0
3	A	53	0	31	3	0
3	B	53	0	31	8	0
3	C	53	0	31	1	0
3	D	53	0	31	3	0
4	A	55	0	0	1	0
4	B	54	0	0	8	0
4	C	66	0	0	2	0
4	D	50	0	0	1	0
All	All	16332	0	15782	778	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 778 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:GLY:H	1:A:451:SER:HA	1.00	1.07
1:A:289:ILE:HD11	1:A:462:MET:HG3	1.35	1.05
1:A:122:MET:HE1	1:A:145:VAL:HG22	1.39	1.05
1:D:32:ILE:HD12	1:D:507:LEU:HD12	1.40	1.02
1:D:278:LEU:HA	1:D:470:GLN:HE22	1.25	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/513 (96%)	418 (85%)	53 (11%)	21 (4%)	3	4
1	B	486/513 (95%)	420 (86%)	45 (9%)	21 (4%)	3	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	489/513 (95%)	426 (87%)	49 (10%)	14 (3%)	5	8
1	D	484/513 (94%)	404 (84%)	61 (13%)	19 (4%)	3	4
All	All	1951/2052 (95%)	1668 (86%)	208 (11%)	75 (4%)	4	5

5 of 75 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	289	ILE
1	A	290	HIS
1	A	291	PHE
1	A	420	PRO
1	B	133	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/454 (97%)	417 (95%)	22 (5%)	28	53
1	B	435/454 (96%)	415 (95%)	20 (5%)	31	58
1	C	437/454 (96%)	420 (96%)	17 (4%)	37	65
1	D	433/454 (95%)	412 (95%)	21 (5%)	30	55
All	All	1744/1816 (96%)	1664 (95%)	80 (5%)	31	58

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	377	GLN
1	C	124	LYS
1	D	291	PHE
1	B	382	ASN
1	B	471	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	255	GLN
1	B	439	ASN
1	D	329	ASN
1	B	259	ASN
1	B	362	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	FAD	A	803	-	51,58,58	1.50	7 (13%)	54,89,89	2.09	10 (18%)
2	SPM	A	924	-	13,13,13	0.94	1 (7%)	12,12,12	0.56	0
3	FAD	B	804	-	51,58,58	1.49	8 (15%)	54,89,89	2.27	11 (20%)
2	SPM	B	923	-	13,13,13	0.97	1 (7%)	12,12,12	0.59	0
3	FAD	C	801	-	51,58,58	1.44	10 (19%)	54,89,89	2.51	11 (20%)
2	SPM	C	921	-	13,13,13	0.97	1 (7%)	12,12,12	0.59	0
3	FAD	D	802	-	51,58,58	1.53	7 (13%)	54,89,89	2.16	9 (16%)
2	SPM	D	922	-	13,13,13	0.98	1 (7%)	12,12,12	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	803	-	-	0/28/50/50	0/6/6/6
2	SPM	A	924	-	-	0/11/11/11	0/0/0/0
3	FAD	B	804	-	-	0/28/50/50	0/6/6/6
2	SPM	B	923	-	-	0/11/11/11	0/0/0/0
3	FAD	C	801	-	-	0/28/50/50	0/6/6/6
2	SPM	C	921	-	-	0/11/11/11	0/0/0/0
3	FAD	D	802	-	-	0/28/50/50	0/6/6/6
2	SPM	D	922	-	-	0/11/11/11	0/0/0/0

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	801	FAD	C9-C8	2.02	1.43	1.37
3	B	804	FAD	C5X-N5	2.10	1.38	1.35
2	C	921	SPM	C9-N10	2.11	1.53	1.47
2	A	924	SPM	C9-N10	2.12	1.53	1.47
2	D	922	SPM	C9-N10	2.13	1.53	1.47

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	801	FAD	O2A-PA-O5B	-6.75	76.25	108.14
3	C	801	FAD	O2A-PA-O1A	-5.50	83.80	112.28
3	B	804	FAD	C4B-O4B-C1B	-5.18	104.25	109.77
3	B	804	FAD	C4X-C4-N3	-4.59	116.95	123.48
3	C	801	FAD	C4X-C4-N3	-4.59	116.95	123.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	803	FAD	3	0
3	B	804	FAD	8	0
3	C	801	FAD	1	0
3	D	802	FAD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	498/513 (97%)	-0.52	6 (1%) 79 75	20, 38, 63, 93	22 (4%)
1	B	492/513 (95%)	-0.49	3 (0%) 89 88	18, 37, 61, 79	37 (7%)
1	C	495/513 (96%)	-0.52	3 (0%) 89 88	19, 36, 63, 87	33 (6%)
1	D	490/513 (95%)	-0.43	11 (2%) 62 56	18, 40, 68, 92	34 (6%)
All	All	1975/2052 (96%)	-0.49	23 (1%) 79 75	18, 38, 65, 93	126 (6%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	423	ASN	5.2
1	D	134	LEU	4.1
1	A	425	ALA	3.9
1	B	135	GLY	3.8
1	D	232	LYS	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	FAD	C	801	53/53	0.97	0.16	1.12	20,28,40,44	0
3	FAD	A	803	53/53	0.97	0.14	0.00	27,32,36,37	0
3	FAD	D	802	53/53	0.97	0.12	-0.17	28,34,39,39	0
3	FAD	B	804	53/53	0.96	0.13	-0.18	23,32,40,41	0
2	SPM	C	921	14/14	-	-	-	175,178,181,181	14
2	SPM	D	922	14/14	-	-	-	161,165,169,169	14
2	SPM	B	923	14/14	-	-	-	134,138,142,142	14
2	SPM	A	924	14/14	-	-	-	142,144,145,145	14

6.5 Other polymers [i](#)

There are no such residues in this entry.