



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 2, 2017 – 12:09 PM EDT

PDB ID : 1Y1S  
Title : Crystal Structure of the Uridine Phosphorylase from Salmonella Typhimurium  
in Complex with Uracil and Sulfate Ion at 2.55Å Resolution  
Authors : Gabdoulkhakov, A.G.; Dontsova, M.V.; Kachalova, G.S.; Betzel, C.; Ealick,  
S.E.; Mikhailov, A.M.  
Deposited on : unknown  
Resolution : 2.55 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4993 (2.58-2.50)
Clashscore	112137	5755 (2.58-2.50)
Ramachandran outliers	110173	5652 (2.58-2.50)
Sidechain outliers	110143	5654 (2.58-2.50)
RSRZ outliers	101464	5026 (2.58-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	<div> <div>2%</div> <div>69%</div> <div>26%</div> <div>..</div> </div>
1	B	253	<div> <div>3%</div> <div>72%</div> <div>23%</div> <div>...</div> </div>
1	C	253	<div> <div>5%</div> <div>65%</div> <div>30%</div> <div>..</div> </div>
1	D	253	<div> <div>3%</div> <div>67%</div> <div>27%</div> <div>..</div> </div>
1	E	253	<div> <div>4%</div> <div>66%</div> <div>28%</div> <div>5% .</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	253	<div><div></div><div>3%</div><div>71%</div><div>25%</div><div>..</div></div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uridine phosphorylase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			
1	C	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			
1	D	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			
1	F	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			
1	E	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			
1	B	250	Total	C	N	O	S	0	0	0
			1876	1174	330	360	12			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is URACIL (three-letter code: URA) (formula:  $C_4H_4N_2O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 8 4 2 2	0	0
3	D	1	Total C N O 8 4 2 2	0	0
3	F	1	Total C N O 8 4 2 2	0	0
3	E	1	Total C N O 8 4 2 2	0	0
3	B	1	Total C N O 8 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total 44	O 44	0	0
4	C	33	Total 33	O 33	0	0
4	D	45	Total 45	O 45	0	0
4	F	34	Total 34	O 34	0	0
4	E	25	Total 25	O 25	0	0
4	B	37	Total 37	O 37	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Uridine phosphorylase



#### • Molecule 1: Uridine phosphorylase



#### • Molecule 1: Uridine phosphorylase







• Molecule 1: Uridine phosphorylase



• Molecule 1: Uridine phosphorylase



• Molecule 1: Uridine phosphorylase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.99Å 124.12Å 133.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.84 – 2.55 28.96 – 2.51	Depositor EDS
% Data completeness (in resolution range)	96.3 (26.84-2.55) 96.6 (28.96-2.51)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.92 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.189 , 0.231 0.189 , 0.230	Depositor DCC
$R_{free}$ test set	2357 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.1	Xtriage
Anisotropy	0.894	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 55.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	11544	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: URA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.35	0/1906	0.63	0/2584
1	B	0.40	1/1906 (0.1%)	0.75	5/2584 (0.2%)
1	C	0.35	0/1906	0.65	0/2584
1	D	0.34	0/1906	0.63	0/2584
1	E	0.39	0/1906	0.70	0/2584
1	F	0.38	0/1906	0.74	2/2584 (0.1%)
All	All	0.37	1/11436 (0.0%)	0.68	7/15504 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	6232	GLU	CG-CD	5.33	1.59	1.51

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	4231	ALA	N-CA-C	9.75	137.32	111.00
1	B	6231	ALA	N-CA-C	7.89	132.30	111.00
1	F	4230	ASN	N-CA-C	6.91	129.67	111.00
1	B	6233	THR	N-CA-C	6.42	128.33	111.00
1	B	6232	GLU	CA-C-N	-6.17	103.62	117.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1876	0	1887	51	0
1	B	1876	0	1887	50	0
1	C	1876	0	1887	72	0
1	D	1876	0	1887	62	0
1	E	1876	0	1887	66	0
1	F	1876	0	1887	62	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	8	0	3	0	0
3	B	8	0	3	0	0
3	D	8	0	3	0	0
3	E	8	0	3	0	0
3	F	8	0	3	0	0
4	A	44	0	0	1	0
4	B	37	0	0	4	0
4	C	33	0	0	0	0
4	D	45	0	0	0	0
4	E	25	0	0	2	0
4	F	34	0	0	2	0
All	All	11544	0	11337	352	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 352 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:2235:LYS:HD2	1:C:2235:LYS:H	1.10	1.13
1:B:6232:GLU:OE1	1:B:6232:GLU:HA	1.65	0.95
1:B:6071:GLY:HA3	1:B:6201:THR:HG21	1.50	0.94
1:C:2235:LYS:N	1:C:2235:LYS:HD2	1.84	0.92
1:B:6201:THR:HG22	4:B:9210:HOH:O	1.72	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/253 (98%)	232 (94%)	12 (5%)	4 (2%)	11	19
1	B	248/253 (98%)	221 (89%)	23 (9%)	4 (2%)	11	19
1	C	248/253 (98%)	225 (91%)	20 (8%)	3 (1%)	15	26
1	D	248/253 (98%)	230 (93%)	12 (5%)	6 (2%)	7	10
1	E	248/253 (98%)	230 (93%)	15 (6%)	3 (1%)	15	26
1	F	248/253 (98%)	230 (93%)	17 (7%)	1 (0%)	38	57
All	All	1488/1518 (98%)	1368 (92%)	99 (7%)	21 (1%)	13	22

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	2230	ASN
1	C	2235	LYS
1	D	3005	ASP
1	E	5231	ALA
1	B	6233	THR

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/202 (98%)	187 (94%)	12 (6%)	22	39
1	B	199/202 (98%)	187 (94%)	12 (6%)	22	39
1	C	199/202 (98%)	187 (94%)	12 (6%)	22	39

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	199/202 (98%)	185 (93%)	14 (7%)	18	31
1	E	199/202 (98%)	183 (92%)	16 (8%)	14	25
1	F	199/202 (98%)	186 (94%)	13 (6%)	20	35
All	All	1194/1212 (98%)	1115 (93%)	79 (7%)	19	34

5 of 79 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	3229	PRO
1	F	4215	MET
1	B	6201	THR
1	D	3230	ASN
1	F	4127	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	3103	ASN
1	D	3236	GLN
1	B	6103	ASN
1	D	3017	GLN
1	E	5014	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	A	7001	-	4,4,4	0.37	0	6,6,6	0.14	0
3	URA	A	8001	-	6,8,8	2.98	2 (33%)	5,10,10	12.41	4 (80%)
2	SO4	B	7006	-	4,4,4	0.36	0	6,6,6	0.08	0
3	URA	B	8006	-	6,8,8	2.75	2 (33%)	5,10,10	12.35	4 (80%)
2	SO4	C	7002	-	4,4,4	0.39	0	6,6,6	0.11	0
2	SO4	D	7003	-	4,4,4	0.33	0	6,6,6	0.15	0
3	URA	D	8003	-	6,8,8	2.89	2 (33%)	5,10,10	12.28	4 (80%)
2	SO4	E	7005	-	4,4,4	0.33	0	6,6,6	0.09	0
3	URA	E	8005	-	6,8,8	3.06	2 (33%)	5,10,10	12.28	4 (80%)
2	SO4	F	7004	-	4,4,4	0.35	0	6,6,6	0.08	0
3	URA	F	8004	-	6,8,8	3.04	2 (33%)	5,10,10	12.40	4 (80%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	7001	-	-	0/0/0/0	0/0/0/0
3	URA	A	8001	-	-	0/0/0/0	0/1/1/1
2	SO4	B	7006	-	-	0/0/0/0	0/0/0/0
3	URA	B	8006	-	-	0/0/0/0	0/1/1/1
2	SO4	C	7002	-	-	0/0/0/0	0/0/0/0
2	SO4	D	7003	-	-	0/0/0/0	0/0/0/0
3	URA	D	8003	-	-	0/0/0/0	0/1/1/1
2	SO4	E	7005	-	-	0/0/0/0	0/0/0/0
3	URA	E	8005	-	-	0/0/0/0	0/1/1/1
2	SO4	F	7004	-	-	0/0/0/0	0/0/0/0
3	URA	F	8004	-	-	0/0/0/0	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	8003	URA	C6-N1	3.59	1.42	1.34
3	B	8006	URA	C6-N1	3.67	1.42	1.34
3	F	8004	URA	C6-N1	3.70	1.42	1.34
3	A	8001	URA	C6-N1	3.73	1.42	1.34
3	E	8005	URA	C6-N1	3.76	1.42	1.34

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	8001	URA	N1-C2-N3	-18.92	114.80	128.40
3	B	8006	URA	N1-C2-N3	-18.77	114.90	128.40
3	F	8004	URA	N1-C2-N3	-18.74	114.92	128.40
3	D	8003	URA	N1-C2-N3	-18.57	115.04	128.40
3	E	8005	URA	N1-C2-N3	-18.56	115.05	128.40

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/253 (98%)	-0.38	5 (2%) 65 68	6, 19, 45, 92	0
1	B	250/253 (98%)	-0.28	8 (3%) 48 52	6, 20, 43, 109	0
1	C	250/253 (98%)	-0.04	12 (4%) 31 34	6, 25, 58, 104	0
1	D	250/253 (98%)	-0.21	8 (3%) 48 52	7, 22, 52, 133	0
1	E	250/253 (98%)	-0.23	9 (3%) 43 47	7, 21, 51, 100	0
1	F	250/253 (98%)	-0.27	7 (2%) 53 57	5, 20, 45, 120	0
All	All	1500/1518 (98%)	-0.24	49 (3%) 47 51	5, 21, 52, 133	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1234	MET	12.1
1	D	3231	ALA	8.4
1	D	3230	ASN	8.4
1	B	6234	MET	7.7
1	C	2230	ASN	7.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.



## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	C	7002	5/5	0.94	0.19	0.58	48,60,62,68	0
3	URA	E	8005	8/8	0.94	0.17	0.55	29,34,39,43	0
2	SO4	B	7006	5/5	0.99	0.15	0.50	12,14,31,46	0
3	URA	B	8006	8/8	0.97	0.12	0.12	7,14,19,33	0
3	URA	F	8004	8/8	0.96	0.12	-0.15	4,6,17,19	0
2	SO4	F	7004	5/5	0.99	0.14	-0.23	16,24,27,50	0
3	URA	D	8003	8/8	0.97	0.12	-0.68	18,26,31,31	0
3	URA	A	8001	8/8	0.98	0.09	-0.96	10,15,19,24	0
2	SO4	E	7005	5/5	0.99	0.12	-0.96	28,29,39,50	0
2	SO4	D	7003	5/5	0.99	0.11	-1.70	18,26,31,40	0
2	SO4	A	7001	5/5	0.99	0.07	-1.82	6,7,23,48	0

## 6.5 Other polymers

There are no such residues in this entry.