



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2018 – 12:29 PM EST

PDB ID : 5Y77
Title : Crystal structure of Pseudomonas fluorescens Kynurenine 3-monooxygenase in complex with L-KYN (seMet derivative)
Authors : Xiang, Y.; Gao, J.J.; Zhu, D.Y.
Deposited on : 2017-08-16
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	<div> <div>4%</div> <div>81%</div> <div>13%</div> <div>..</div> </div>
1	B	473	<div> <div>8%</div> <div>74%</div> <div>20%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	KYN	A	502	-	-	X	-
3	KYN	B	502	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8388 atoms, of which 22 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kynurenine 3-monooxygenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	455	Total	C	N	O	S	Se	0	16	0
			3610	2267	668	653	7	15			
1	B	451	Total	C	N	O	S	Se	0	14	0
			3566	2239	658	649	6	14			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	expression tag	UNP Q84HF5
A	1	GLY	-	expression tag	UNP Q84HF5
A	462	ALA	-	expression tag	UNP Q84HF5
A	463	ALA	-	expression tag	UNP Q84HF5
A	464	ALA	-	expression tag	UNP Q84HF5
A	465	LEU	-	expression tag	UNP Q84HF5
A	466	GLU	-	expression tag	UNP Q84HF5
A	467	HIS	-	expression tag	UNP Q84HF5
A	468	HIS	-	expression tag	UNP Q84HF5
A	469	HIS	-	expression tag	UNP Q84HF5
A	470	HIS	-	expression tag	UNP Q84HF5
A	471	HIS	-	expression tag	UNP Q84HF5
A	472	HIS	-	expression tag	UNP Q84HF5
B	0	MSE	-	expression tag	UNP Q84HF5
B	1	GLY	-	expression tag	UNP Q84HF5
B	462	ALA	-	expression tag	UNP Q84HF5
B	463	ALA	-	expression tag	UNP Q84HF5
B	464	ALA	-	expression tag	UNP Q84HF5
B	465	LEU	-	expression tag	UNP Q84HF5
B	466	GLU	-	expression tag	UNP Q84HF5
B	467	HIS	-	expression tag	UNP Q84HF5
B	468	HIS	-	expression tag	UNP Q84HF5
B	469	HIS	-	expression tag	UNP Q84HF5
B	470	HIS	-	expression tag	UNP Q84HF5
B	471	HIS	-	expression tag	UNP Q84HF5


Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	472	HIS	-	expression tag	UNP Q84HF5

-
- The chemical structure of FAD (Flavin Adenine Dinucleotide) is shown. It consists of a flavin mononucleotide (FMN) part and an adenosine diphosphate (ADP) part. The FMN part includes a ribityl chain, an isoalloxazine ring system, and a ribose sugar. The ADP part includes an adenine base, a ribose sugar, and a diphosphate group. The structure is labeled with various atoms and bonds, including nitrogen (N), oxygen (O), carbon (C), and phosphorus (P). The overall structure is a complex molecule with multiple rings and functional groups.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is (2S)-2-amino-4-(2-aminophenyl)-4-oxobutanoic acid (three-letter code: KYN) (formula: $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			26	10	11	2	3		
3	B	1	Total	C	H	N	O	0	0
			26	10	11	2	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	643	Total	O	0	0
			643	643		
4	B	411	Total	O	0	0
			411	411		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Kynurenine 3-monooxygenase



• Molecule 1: Kynurenine 3-monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	69.56Å 52.45Å 135.71Å 90.00° 104.33° 90.00°	Depositor
Resolution (Å)	34.78 – 1.60 34.78 – 1.60	Depositor EDS
% Data completeness (in resolution range)	95.0 (34.78-1.60) 95.0 (34.78-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.96 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.185 , 0.217 0.185 , 0.218	Depositor DCC
R_{free} test set	5969 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8388	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 42.61 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9904e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KYN, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	3/3721 (0.1%)	0.99	20/5028 (0.4%)
1	B	0.69	0/3671	0.85	5/4963 (0.1%)
All	All	0.75	3/7392 (0.0%)	0.92	25/9991 (0.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	176	MSE	SE-CE	-7.50	1.51	1.95
1	A	329	GLU	CG-CD	5.14	1.59	1.51
1	A	203	SER	CB-OG	-5.12	1.35	1.42

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	NE-CZ-NH1	11.64	126.12	120.30
1	B	173	ARG	NE-CZ-NH2	-10.25	115.17	120.30
1	A	173	ARG	NE-CZ-NH2	-10.10	115.25	120.30
1	A	38[A]	ARG	NE-CZ-NH2	-8.91	115.84	120.30
1	A	38[B]	ARG	NE-CZ-NH2	-8.91	115.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3610	0	3616	64	0
1	B	3566	0	3569	88	1
2	A	53	0	29	1	0
2	B	53	0	31	3	0
3	A	15	11	11	9	0
3	B	15	11	11	3	0
4	A	643	0	0	22	3
4	B	411	0	0	27	0
All	All	8366	22	7267	159	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:452:ARG:NH1	4:B:601:HOH:O	1.96	0.96
1:B:101:ASP:OD2	1:B:103:SER:N	2.03	0.90
1:B:403[B]:ARG:NH1	4:B:602:HOH:O	2.04	0.90
1:B:49:ARG:HD3	1:B:49:ARG:H	1.38	0.87
1:B:39:ARG:NH2	4:B:603:HOH:O	2.08	0.85

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1128:HOH:O	4:A:1196:HOH:O[1_565]	2.09	0.11
1:B:45:GLU:OE2	1:B:82:ARG:NH2[2_655]	2.11	0.09
4:A:977:HOH:O	4:A:1106:HOH:O[2_546]	2.15	0.05
4:A:1006:HOH:O	4:A:1184:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	469/473 (99%)	460 (98%)	9 (2%)	0	100	100
1	B	463/473 (98%)	448 (97%)	15 (3%)	0	100	100
All	All	932/946 (98%)	908 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/366 (105%)	372 (97%)	11 (3%)	48	20
1	B	378/366 (103%)	363 (96%)	15 (4%)	36	11
All	All	761/732 (104%)	735 (97%)	26 (3%)	40	15

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	22	VAL
1	B	77	GLU
1	B	412	LEU
1	B	49	ARG
1	B	53	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	32	GLN
1	A	132	ASN
1	B	34	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	501	-	51,58,58	2.21	17 (33%)	54,89,89	3.42	21 (38%)
3	KYN	A	502	-	10,15,15	1.48	2 (20%)	14,20,20	3.26	8 (57%)
2	FAD	B	501	-	51,58,58	1.88	10 (19%)	54,89,89	3.60	19 (35%)
3	KYN	B	502	-	10,15,15	1.31	2 (20%)	14,20,20	2.87	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/28/50/50	0/6/6/6
3	KYN	A	502	-	-	0/8/12/12	0/1/1/1
2	FAD	B	501	-	-	0/28/50/50	0/6/6/6
3	KYN	B	502	-	-	0/8/12/12	0/1/1/1

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	FAD	C7M-C7	-4.75	1.41	1.51
2	B	501	FAD	C8M-C8	-3.80	1.43	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	C7M-C7	-3.15	1.44	1.51
2	A	501	FAD	C9A-N10	-3.05	1.34	1.38
2	A	501	FAD	O3B-C3B	-2.93	1.36	1.43

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	FAD	N3A-C2A-N1A	-14.63	116.12	128.86
2	A	501	FAD	N3A-C2A-N1A	-12.89	117.63	128.86
2	B	501	FAD	O4B-C4B-C5B	-8.08	82.12	109.40
2	A	501	FAD	O4B-C4B-C5B	-7.78	83.13	109.40
2	B	501	FAD	O4B-C4B-C3B	-7.43	90.40	105.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	1	0
3	A	502	KYN	9	0
2	B	501	FAD	3	0
3	B	502	KYN	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	442/473 (93%)	-0.07	18 (4%)	38 36	11, 20, 46, 79	0
1	B	438/473 (92%)	0.22	36 (8%)	12 12	14, 27, 63, 75	0
All	All	880/946 (93%)	0.07	54 (6%)	22 21	11, 24, 57, 79	0

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	461	CYS	11.7
1	A	460	LEU	7.6
1	B	150	VAL	6.1
1	A	245	ALA	6.1
1	B	247	GLN	6.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	KYN	B	502	15/15	0.91	0.20	3.44	14,35,48,55	0
3	KYN	A	502	15/15	0.88	0.16	0.86	16,35,48,50	0
2	FAD	B	501	53/53	0.97	0.10	0.04	12,16,25,35	0
2	FAD	A	501	53/53	0.97	0.10	-0.05	8,12,15,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.